

Waikato Healthy Rivers


Plan Change 1
Hearing 1





Timberlands Limited








Important improvement	Waikato river sampling sites									
Slight improvement Uncertain Slight deterioration Important deterioration	Taupo	Ohaaki	Ohakuri	Whakamaru	Waipapa	Narrows	Horotiu	Huntly	Mercer	Tuakau
Temperature	8	0	8	-	-	()	-	9,—3	0	0
Dissolved oxygen	-	0	-	1	8	8	8	-	8	(3)
Visual clarity	nd	8	(3)	8	8	0	0	33 3	nd	8
Turbidity	8	8	8	-	-	0	1	0 .	8	8
Arsenic	8	0	-	_	-	0	0	0	0	0
Ammonia	_	_		71 <u>—1</u> 3	<u></u>	8—0	0	0	_	_
Total nitrogen	-	(3)	(3)	(3)	8	8	8	8	(3)	(3)
Total phosphorus	0	0	-	_		0	0	0	0	0
Chlorophyll a	-	-	2011	nd	<u>S1/4</u> 101/7/	0	0	0	0	0
E. coli	_	0		8	_	_		_	·	-

nd = no data Flow-adjusted data, n ≤ 300 LOWESS span 30% Seasonal Kendall slope and test Significance: p < 5% Note: visual clarity, turbidity and arsanic are for the period since 1995, and E. coli is for the period since 1998.











Timberlands Limited



