

REVIEW ARTICLE

# Barriers to ecosystem restoration presented by soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species: implications for ecological restoration

Mlungele M. Nsikani<sup>1,2</sup> , Brian W. van Wilgen<sup>1</sup>, Mirijam Gaertner<sup>1,3,4</sup>

Impacts of invasive alien N<sub>2</sub>-fixing woody species and how they can persist as soil legacy effects after invasive species control are well appreciated, but how soil legacy effects can present barriers to restoration is poorly understood. Finding better ways to deal with these barriers to restoration is essential to improving restoration outcomes. In this study, we review 440 studies to identify barriers to restoration and potential management actions for the barriers to restoration, and provide practical application examples of the management actions. Our findings suggest that altered soil microbial communities, depleted native soil seed banks, elevated N status, secondary invasion and weedy native species dominance, and reinvasion are potential barriers to restoration. Furthermore, carbon addition, litter removal, soil microbial treatments, establishing species adapted to low N levels, prescribed burning, classical biological control, grazing, mowing, herbicide or graminicide application, manual weeding, soil N management, soil solarization, weed mats, native species reintroduction, and nurse plants are potential management actions for these barriers to restoration. However, there is little evidence suggesting that several of these barriers to restoration hinder improved restoration outcomes and this could be due to little research on them. More research is needed to assess their relative importance in hindering improved restoration outcomes. Management actions are rarely applied in combination, despite that they often address distinct barriers to restoration. Management actions should be combined into an integrated management effort to improve restoration outcomes.

**Key words:** barriers to restoration, biological invasion, legacy effects, management, N<sub>2</sub>-fixing, woody species

## Implications for Practice

- To address barriers to restoration from soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species during restoration, if available, classical biological control could be established to reduce seed production and seed banks.
- Where appropriate, prescribed burning could be applied after clearing the invasive species. However, where burning is inappropriate, litter could be manually removed. Subsequently, soil solarization or weed mats could be set up to reduce establishment of secondary invaders, weedy native species and a second generation of the invasive species. Mowing, manual weeding, grazing, herbicide or graminicide application could be used to manage any resulting vegetation.
- Subsequently, soil carbon addition and microbial treatments could be applied and native species reintroduced with or without the aid of nurse plants.

## Introduction

The widespread introduction of plant species to areas outside of their natural distribution ranges has led to some becoming invasive (Vitousek et al. 1997). Many of these invasive alien species are trees and shrubs (Richardson & Rejmánek 2011) and some are nitrogen-fixing such as Australian acacias

(Richardson et al. 2011). Invasive N<sub>2</sub>-fixing woody species often transform ecosystems by altering ecosystem processes and displacing native species (Vilà et al. 2011). The negative impacts of invasive N<sub>2</sub>-fixing woody species on the soil include altered soil chemistry (Ehrenfeld 2003), establishment of large seed banks (D'Antonio & Meyerson 2002; Richardson & Kluge 2008), deposition of novel allelochemicals (Callaway & Ride-nour 2004), and altered soil microbial community composition and function (Inderjit & Van Der Putten 2010). Mechanisms underlying such impacts are well documented (e.g. Levine et al. 2003; Vilà et al. 2011).

Worldwide efforts are underway to clear invasive species and restore historical ecosystems (D'Antonio & Meyerson 2002;

Author contributions: MMN, MG conceived and designed the review; MMN conducted the review; all authors wrote and edited the manuscript.

<sup>1</sup>Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Private Bag X1, Matieland 7602, Stellenbosch, South Africa

<sup>2</sup>Address correspondence to M. M. Nsikani, email mllue06@gmail.com

<sup>3</sup>Invasive Species Unit, Environmental Resource Management Department, City of Cape Town, Westlake Conservation Office, Ou Kaapse Weg, Tokai, 7966, Cape Town, South Africa

<sup>4</sup>Nürtingen-Geislingen University of Applied Sciences (HFU), Schelmenwasen 4-8, 72622, Nürtingen, Germany

© 2018 Society for Ecological Restoration

doi: 10.1111/rec.12669

Supporting information at:

<http://onlinelibrary.wiley.com/doi/10.1111/rec.12669/supinfo>

Suding et al. 2004). It is often assumed that negative impacts of invasive species will diminish after clearing (Wittenberg & Cock 2005), but this is not always the case because the invasive species can leave persistent legacy effects (i.e. measurable changes to biological, chemical, or physical conditions) in the soil (Corbin & D'Antonio 2012). Ecophysiological traits of invasive N<sub>2</sub>-fixing woody species (e.g. early and high production of seeds with long dormancy periods, high growth rates, and increasing the N content of N-limited soil) contribute to the persistence of soil legacy effects after their clearing (Pyšek & Richardson 2007). Numerous legacy effects of invasive species have been identified (reviewed by Corbin & D'Antonio 2012). Restoration of historical ecosystems hence often fails after invasive species clearing, probably because soil legacy effects create barriers to restoration (Corbin & D'Antonio 2012).

A growing number of studies describe how soil legacy effects present barriers to restoration after removal of invasive N<sub>2</sub>-fixing woody species (Appendix S2) while others present potential management actions to address these barriers to restoration (Appendix S3). Efforts have been made to review potential management actions for these barriers to restoration (e.g. Perry et al. 2010), but these have only focused on management of individual barriers to restoration. We still lack a broad review of barriers to restoration presented by soil legacy effects and management actions that could potentially be used to address them. In this study, we review (1) how soil legacy effects present barriers to restoration after invasive N<sub>2</sub>-fixing woody species clearing; (2) potential management actions to address those barriers to restoration; and (3) we give practical examples of their application. To maximize the usefulness of the review to restoration practice, we discuss all soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species, not just those directly related to N<sub>2</sub>-fixation and elevated soil N.

## Methods

We searched for relevant articles on the ISI Web of Science database (<http://www.webofknowledge.com>) with no restriction on publication year, using a range of keywords (Table 1) designed to locate articles that document barriers to restoration presented by soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species and their management. Abstracts of retrieved articles were read and those relevant to this study were selected and the full paper content read.

## Results

We identified 440 articles (35 on barriers to restoration, 399 on management actions, and 6 discussing both aspects; Appendix S1) that were relevant to this study. Some of those articles were reviews (e.g. Le Maitre et al. 2011) that included studies from the "gray literature"; therefore, we achieved a reasonably good coverage of the literature on barriers to restoration and their management, not restricted to that indexed in the Web of Science. We identified that elevated N status, secondary invasion and weedy native species dominance, reinvasion, depleted

**Table 1.** Combinations of search terms designed to locate studies that document barriers to restoration presented by soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species and potential management actions to address them.

<i>Barriers to restoration and potential management actions specific to invasive N<sub>2</sub>-fixing woody species</i>	<i>Potential management actions for barriers to restoration that also apply to invasive non-N<sub>2</sub>-fixing woody species</i>	<i>Potential management actions for each identified barrier to restoration</i>
exotic* OR invasive* OR invasion* OR alien* OR invader* OR non-native* OR nonnative AND nitrogen-fix* OR nitrogen fix* OR dinitrogen-fix* OR OR dinitrogen fix* OR N-fix* OR OR N fix* OR N2-fix OR N2 fix OR OR actinorhizal* OR OR legume* OR leguminous* OR root nodule* AND impact* OR effect* OR OR legacy* OR legacies* OR legacy effect* AND native* OR indigenous* OR restoration* OR recovery* OR reestablishment* OR OR return* OR manage*	exotic* OR invasive* OR invasion* OR alien* OR invader* OR non-native* AND legac* OR residual* OR long lasting* AND restor* OR recover* OR return* OR manage*	seed bank* OR seedbank* OR microbe* OR reinvasion* OR secondary invad* OR OR ruderal* OR soil nitrogen* AND native* OR indigenous* OR restor* OR recover* OR return* OR manage* OR reestablishment*

native soil seed banks, and altered soil microbial communities have all been noted as barriers to restoration following clearing of invasive, N<sub>2</sub>-fixing woody species (Fig. 1). We further identified that carbon addition, litter removal, soil microbial treatments, establishing species adapted to low N levels, prescribed burning, classical biological control, and grazing are methods that have been used to manage these barriers to restoration (Fig. 2). Furthermore, mowing, herbicide or graminicide application, manual weeding, soil N management, soil solarization, weed mats, native species reintroduction, and nurse plants are methods that have been used to manage these barriers to restoration (Fig. 2).

## Barriers to Restoration

The subject of barriers to restoration presented by soil legacy effects of invasive N<sub>2</sub>-fixing woody species has not been




Figure 1. Percentage of cases ( $n = 53$ ) identified from 41 studies describing barriers to restoration presented by soil legacy effects of invasive alien  $N_2$ -fixing woody species included in this review. ASMC, altered soil microbial communities; DNSB, depleted native species' soil seed banks; ESNS, elevated soil N status; RI, reinvasion; SIWN, secondary invaders and weedy native species.

extensively reported in the literature as shown by the small number (41) of publications selected for this review. Many important concepts are described by relatively few studies, with a strong bias towards certain regions—e.g. the United States (Appendix S2). While more, and as yet unidentified, barriers to restoration may exist, we believe that the few available studies have allowed us to develop a fairly robust picture of how legacy effects of invasive  $N_2$ -fixing woody species could present barriers to restoration. Each of the barriers to restoration is described in the following subsections.

**Elevated Soil N Status.** Invasive  $N_2$ -fixing woody species generally increase soil N through  $N_2$ -fixation and production of N-rich litter (Vitousek & Walker 1989; Malcolm et al. 2008). After clearing, elevated N content and availability, and altered N mineralization, can persist and increase further with decomposition of invader litter and roots that remain. Alterations to soil N status are often measured through soil sample analysis by comparing non-invaded, invaded, and/or previously invaded sites (Yelenik et al. 2004; Von Holle et al. 2013). Native species in some habitats are adapted to low soil N—e.g. pine-oak systems in the United States; hence, persistent elevated soil N status can directly hinder restoration by negatively affecting their germination, growth, diversity, and/or indirectly hinder restoration by giving a competitive advantage to weedy species adapted to high N availability (Rice et al. 2004).

**Secondary Invasion and/or Weedy Native Species Dominance.** Seeds of alien or native nitrophilous species can be present in the soil seed bank of restoration sites or disperse to such sites from surrounding areas (Yelenik et al. 2004; Pearson et al. 2016). These nitrophilous species are often more competitive than native restoration species under high N conditions (Pearson et al. 2016). Such species often take advantage of the conditions created by removing invasive  $N_2$ -fixing woody species to establish (Pearson et al. 2016). Moreover, the elevated soil N status created by the invasive  $N_2$ -fixing

woody species facilitates such species' growth and dominance (Vitousek & Walker 1989; Le Maitre et al. 2011). Secondary invaders and weedy native species have been observed to hinder restoration by limiting the growth of native restoration species (Maron & Connors 1996; Yelenik et al. 2004; Marchante et al. 2009; Von Holle et al. 2013).

**Reinvasion.** Invasive  $N_2$ -fixing woody species often produce copious amounts of seed that can persist in the seed bank for extended periods (Oneto et al. 2010). This often leads to germination and reinvasion after clearing. Furthermore, clearing the invasive species often leaves roots or stumps of the plants, which can resprout in some cases (Shortt & Vamosi 2012; Souza-Alonso et al. 2013). Reinvasion of restoration sites (through persistent seed and/or vegetative propagules) allows the invasive species to once again dominate the ecosystem, leading to failed restoration (Holmes & Cowling 1997).

**Depleted Native Soil Seed Banks.** Invasive  $N_2$ -fixing woody species often compete with and exclude native species (Vilà et al. 2011). The exclusion of native species can lead to depleted native soil seed banks—due to native species becoming greatly reduced in numbers, and with survivors producing less seed as they either do not reach maturity or do not flower under the canopy of the invader. After clearing the invasive species, the depleted native soil seed banks often become a barrier to reestablishment of native communities (Malcolm et al. 2008; Le Maitre et al. 2011).

**Altered Soil Microbial Communities.** Invasive  $N_2$ -fixing woody species can alter the soil microbial community composition, diversity, and function through several mechanisms such as deposition of allelochemicals and introduction of novel microbes (Inderjit & Van Der Putten 2010). The soil mycorrhizal community and symbioses of native species are often disrupted and such changes can persist after clearing the invasive species and limit the germination and/or growth of native species (Corbin & D'Antonio 2012; Boudiaf et al. 2013).

#### Management of Barriers to Restoration

These barriers to restoration can be addressed using a range of appropriate management actions (Table 2). Potential management actions for different barriers to restoration are described and illustrated using selected examples below.

**Elevated Soil N Status.** Prescribed burning can be used to remove the invasive species' litter to prevent it from contributing to the soil N pool in the long term (Mitchell et al. 2000). The slash can be spread over the restoration site and burnt, instead of being stacked before burning (DiTomaso et al. 2006). Prescribed burning will initially cause a strong pulse of released N previously immobilized in the litter (Fenn et al. 1998). A significant portion of the released N will be volatilized (Riggan et al. 1994; Marchante et al. 2009), whereas released  $NH_4^+$  will probably be nitrified after burning and result in leached  $NO_3^-$  (Dunn et al. 1979). Repeated burning can deplete soil N through




Figure 2. Percentage of cases ( $n = 476$ ) identified from 405 studies describing potential management actions to address barriers to restoration presented by soil legacy effects of invasive alien  $N_2$ -fixing woody species included in this review. CA, carbon addition; CBC, classical biological control; GZN, grazing; MWG, mowing; HGA, herbicide or graminicide application; LNS, establish species adapted to low N levels; LR, litter removal; MW, manual weeding; NP, nurse plants; NSR, native species reintroduction; PB, prescribed burning; SMT, soil microbial treatments; SNM, soil N management; SS, soil solarization; WM, weed mats.

**Table 2.** Potential management actions to address barriers to restoration presented by soil legacy effects of invasive alien  $N_2$ -fixing woody species.

Management action	Barrier to Restoration				
	Elevated soil N status	Secondary invaders and weedy native species	Reinvasion	Depleted native soil seed banks	Altered soil microbial communities
Soil carbon addition	X				
Litter removal	X				X
Soil microbial treatments					X
Establishing species adapted to low N availability	X				
Prescribed burning	X	X	X		
Classical biological control			X		
Grazing		X	X		
Mowing		X	X		
Herbicide or graminicide application		X	X		
Manual weeding		X	X		
Soil N management		X			
Soil solarization		X	X		
Weed mats		X	X		
Native species reintroduction				X	
Nurse plants				X	

repeated volatilization (Prober et al. 2005). For example, in grasslands of the United States repeated burning reduced total N by up to 40% in the top 10 cm of the soil (Haubensak et al. 2004). There is a need to consider the effects of fire on restoration sites because the application of fire may be inappropriate for some habitat types—e.g. riparian zones (Richardson & Kluge 2008). In areas where the use of fire is inappropriate, manual techniques such as raking can be used to remove the invasive species' litter (Elgersma et al. 2011). However, such techniques are labor intensive and logistically limited to small areas.

Labile carbon sources such as sawdust and sucrose can be added or leached into the soil to immobilize soil N (Blumenthal

et al. 2003; Prober & Thiele 2005). Briefly, the added carbon increases soil microbial biomass and activity—by serving as a substrate for heterotrophic soil microbes, which in turn increases microbial N uptake and lowers soil N availability (Baer et al. 2003). Furthermore, carbon addition in anaerobic soils can increase the activity of denitrifying bacteria and increase N loss through denitrification (Perry et al. 2010). For example, in a coastal sage scrub in the United States, addition of organic mulch resulted in a significant decrease in soil N availability (Zink & Allen 1998). Success of carbon addition depends on whether or not soil microbes are C-limited, method and duration of application, environmental conditions, habitat type, and

target species (Rice et al. 2004; Prober & Thiele 2005), and its long-term effects are often difficult to predict (Török et al. 2014). Furthermore, the success of carbon addition may be hindered because it is labor intensive and expensive to apply at large scale (Perry et al. 2010).

Many species adapted to low N availability can lower soil N because they produce high C:N litter, which slows N cycling, and they have lower minimum N requirements, which allow them to continue to grow and capture more limited available N (Perry et al. 2010). Therefore, establishing species adapted to low N availability in the site designated for restoration can ultimately lower soil N (Perry et al. 2010). For example, the establishment of native *Themeda australis* in temperate grassy woodlands significantly reduced soil N availability (Prober et al. 2005). Native restoration species in areas invaded by N<sub>2</sub>-fixing woody species are often adapted to low N availability; hence, waiting for them to establish from their soil seed banks, seeding, or transplanting them (i.e. if soil seed banks are depleted) in restoration sites may be appropriate. Furthermore, use of nurse plants adapted to low N availability during native species reintroduction could potentially lower soil N. However, establishment of species adapted to low N availability may not significantly lower soil N over short time scales or when certain N levels have been reached (Perry et al. 2010). Repeated prescribed burning can favor species adapted to low N availability (Perry et al. 2010). For example, in an oak savanna in the United States, repeated prescribed burning led to a shift from oak to grass dominance and a subsequent decline in soil N levels due to low tissue N concentrations in grasses (Dijkstra et al. 2006).

#### Secondary Invasion and Weedy Native Species Dominance.

Establishment of secondary invaders and weedy native species can be reduced through soil solarization (i.e. heating the soil surface by covering with a plastic sheet) or setting up weed mats (i.e. woven plastic mats that allow passage of water but prevent emergence of seedlings) in restoration sites. For example, weed mats set up in a coastal dune cleared of *Lupinus arboreus* reduced the establishment of secondary invaders (Pickart et al. 1998). However, the cost and logistical challenges restrict the use of soil solarization or weeds mats to small areas (Pickart et al. 1998; Richardson & Kluge 2008).

Secondary invaders and weedy native species that establish can be manually weeded, mowed, selectively grazed (Maron & Jefferies 2001; Gooden et al. 2009; Milchunas et al. 2011), or controlled through herbicide or graminicide application (Sztár et al. 2016). For example, mowing followed by biomass removal during restoration of coastal prairie grasslands reduced the abundance of secondary invaders (Maron & Jefferies 2001). If grazing or mowing are used to control secondary invaders and weedy native species, season, intensity, and duration of application need to be considered to obtain satisfactory results and avoid negative ecological consequences (Milchunas et al. 2011; Dee et al. 2016). Appropriate herbicides or graminicides should be selected and proper timing of their application, toxicity, residence times, and specificity should be carefully considered (Hobbs & Humphries 1995; Oneto et al. 2010). The technique used to apply the herbicide or graminicide would depend on the

size of the restoration site—e.g. broadcast foliar treatment for large areas and drizzle technique for small areas (Oneto et al. 2010).

Invasive alien and weedy native species often prefer high N availability (D'Antonio & Meyerson 2002; Pearson et al. 2016); hence, their dominance can decline if elevated soil N is addressed with soil N management, as described above (Kulmatiski 2011). For example, reduction of N availability after the death of *L. arboreus* significantly reduced the abundance of secondary invaders (Alpert & Maron 2000).

**Reinvasion.** To address reinvasion by N<sub>2</sub>-fixing, invasive alien plants, classical biological control can be set up when available to reduce seed production and seed banks over time (Buckley et al. 2004). For example, long-term classical biological control of invasive *Acacia* spp. in South Africa using nine insect species and a fungus has led to significant reductions in their seed production, seed banks, and distribution (Moran & Hoffmann 2012). However, It should be noted that biological control is not a quick-fix solution to reduce persistent seed banks (Moran & Hoffmann 2012). Furthermore, predation rate of biological control agents should be high (>90%) for classical biological control to be effective (Noble & Weiss 1989). Biological control agents often take time to establish but once established they can be efficient in reducing fecundity of invasive species by destroying their flowers, buds, or pods (Holmes & Cowling 1997; Buckley et al. 2004).

Numerous invasive species accumulate persistent soil seed banks characterized by seeds that need a heat pulse to break dormancy (Richardson & Kluge 2008). Therefore, persistent soil seed banks can be reduced by triggering mass germination through prescribed burning using low-intensity fires after clearing the invasive species (Holmes & Cowling 1997). Use of low-intensity fires is recommended because high-intensity fires tend to destroy native seeds and seedlings that may be present on restoration sites (Richardson & Kluge 2008). Furthermore, burning will kill a significant part of the invasive species' seed on the soil surface and litter (DiTomaso et al. 2006). Seedlings that germinate can be manually weeded (Fill et al. 2017), treated with herbicides (Krupek et al. 2016), and mowed or selectively grazed (Richardson & Kluge 2008) to avoid the development of a second generation of dense invasive species. Multiple treatments may be required to achieve desired effects (Mandle et al. 2011). For example, a combination of prescribed burning and herbicide application after invasive *Acacia mearnsii* control in South Africa significantly reduced its soil seed banks (Campbell et al. 1999).

Soil solarization and weed mats could be viable alternatives to prescribed burning in sites where use of fire is inappropriate. For example, soil solarization treatments substantially reduced the number of buried seeds of *A. saligna*, *A. murrayana*, and *A. sclerosperma* in Israel (Cohen et al. 2008). Weed mats set up in a coastal dune in the United States reduced reinvasion of *L. arboreus* (Pickart et al. 1998). However, the use of soil solarization and weed mats is logistically and financially limited to small areas (Pickart et al. 1998; Richardson & Kluge 2008).

**Depleted Native Soil Seed Banks.** Native species reintroduction through seed, vegetative propagules, transfer of seed containing plant material, or native topsoil translocation is a viable option to manage depleted native soil seed banks (Holmes & Cowling 1997; Baasch et al. 2012; Ferreira & Vieira 2017). For example, seeding with native species during restoration of coastal sandplain grassland in the United States increased native species diversity (Neill et al. 2015). There is need for careful planning and clearly defined restoration goals before conducting native species reintroduction (Honnay et al. 2002; Sztár et al. 2016). For example, some restoration programs might focus on rehabilitation of functional groups or clusters of focal species, whereas others might focus on particular endangered species (Palmer et al. 1997). Practicing restoration ecologists should consider the native species to be reintroduced, donor sites to be used, timing, order, and methods of reintroduction, seed mixtures, and seeding rates for each species.

To the extent possible, timing, order, and methods of reintroduction should be informed by ecological community theory, based on known patterns of community structure (Zedler 2000). This might involve, for example, mimicking natural successional processes by introducing early-successional or mid-successional species first, to help create suitable conditions for later-successional species introduced later (Lithgow et al. 2013). Alternatively, it could involve introducing rare species first to make sure they are not excluded by more common, rapidly establishing species (Palmer et al. 1997). Seed mixtures should be site specific and carefully selected, but diverse seed mixtures are often preferred because they offer insurance that some species will establish (Kiehl et al. 2010). Seeding rates are difficult to gauge, being species and site specific. Therefore, if available, practicing restoration ecologists should select seeding rates according to reference sites (Holmes & Richardson 1999).

Seeds should be harvested from nearby areas to match the genetic composition of native species that occupied the restoration site before invasion (Schaefer 2011). Furthermore, seeds should be harvested from multiple source populations to increase genetic variability (Ödman et al. 2012). Consideration should be made on the seed ecology of the native species to be reintroduced so that seeds of best quality are collected at the right time and stored appropriately to maintain their viability (Holmes & Richardson 1999). To improve germination, pre-treatment of native species' seeds that require certain conditions to break dormancy should be conducted before sowing (Neill et al. 2015). Seeds should be sown using an appropriate method (e.g. mechanical broadcast seeding) at a time that coincides with ideal conditions for germination and survival (Carrick & Krüger 2007). Care should be taken to cover the seeds with substrate to reduce seed predation and prevent them from blowing away (Pausas et al. 2004).

If available, seed containing plant material (e.g. hay and litter) can be harvested from multiple native sites, transferred, and spread on the site designated for restoration as a seed source (Kiehl et al. 2006). Some species such as late successional and endangered species can be transplanted to the site designated for restoration after being grown in pots to improve their chances of establishment (Manchester et al. 1999). If native topsoil

becomes available due the native site being destroyed, it can be collected (depth of 10 cm), transferred and thinly spread on the site designated for restoration as a seed source (Rokich et al. 2000). The topsoil should not be stockpiled, but spread as soon as possible to maintain viability of seeds in the soil (Rokich et al. 2000). Disturbance during topsoil translocation may be enough to break seed dormancy, but the soil can also be treated with smoke water solution to break the dormancy of some seeds (Rokich et al. 2002). However, not all seeds will germinate (Klimeš et al. 2010). Furthermore, topsoil translocation can damage species that reproduce through vegetative means (Craig & Buckley 2013).

Nurse plants can be planted before or together with the native species to be reintroduced to create safe sites with favorable microclimates for germination and survival of the reintroduced native species (Padilla & Pugnaire 2006). Nurse plants can shade the reintroduced native species, reduce solar radiation and soil temperatures, protect from herbivore damage, and/or improve the water status of the target seedlings by reducing evaporation (Aerts et al. 2007). For example, use of shrubs as nurse plants enhanced growth of native conifers during reforestation in Spain by improving seedling water status, and therefore reducing seedling mortality by drought. (Castro et al. 2002). Species that can be used as nurse plants should have a facilitative effect on the native restoration species (Gómez-Ruiz et al. 2013). It is advisable to use short-lived species as nurse plants, so that they exist for a limited period and eventually give way to the native restoration species (Gómez-Aparicio 2009). Also, nurse plants planted at too high densities may compete with and exclude target species (Paz Esquivias et al. 2015).

**Altered Soil Microbial Communities.** Practicing restoration ecologists should consider determining the extent of changes to microbial communities in the site designated for restoration (e.g. using the analysis of phospholipid fatty acids method to compare microbial communities to reference sites; Frostegård et al. 2011), to inform the selection of appropriate management actions to aid recovery of microbial communities. Soil microbial treatments can be utilized to aid the recovery of microbial communities (Richter & Stutz 2002). For example, application of fungicides, bactericides, and microbial and fungal inoculants had a positive impact on the recovery of soil microbial communities and various native plant species in an experimental study in the United States (Perkins & Hatfield 2016). However, the success of efforts to restore soil microbial communities has been inconsistent, context-dependent, and logistically limited to small areas (Perkins & Hatfield 2016). Restoration of native species in the site designated for restoration could promote arbuscular mycorrhizal fungi proliferation and diversity (Tanner & Gange 2013). Removal of invasive species' litter in the site designated for restoration (through prescribed burning or manually) could be an easier way to rectify altered soil microbial communities (Elgersma et al. 2011). However, changes to microbial communities after litter removal are often minimal and unpredictable (Elgersma et al. 2011).




Figure 3. How to combine potential management actions to address barriers to restoration presented by soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species into an integrated management effort to improve restoration outcomes.

## The Way Forward

Soil legacy effects can present several barriers to restoration after clearing invasive N<sub>2</sub>-fixing woody species (Fig. 1). The unique N<sub>2</sub>-fixing ability of these invasive species contributes directly to elevated N status, secondary invasion and weedy native species dominance, and altered soil microbial communities. Only a few studies in our review provided evidence that elevated N status, depleted native soil seed banks, and altered soil microbial communities are common barriers to restoration, but this does not necessarily indicate that these barriers to restoration are unimportant. Instead, we believe that the low number of reported cases for some barriers to restoration is a result of these barriers to restoration not being investigated during restoration programs. An investigation of these barriers, however, is crucial to gain a better understanding of the mechanisms of barriers

to restoration following invasive species control and of what actions are most needed to improve restoration outcomes.

Management of barriers to restoration is often necessary to improve restoration outcomes and a wide array of potential approaches is available (Fig. 2). We concede that management actions included in this review cannot be uniformly applied in every site designated for restoration because their selection is often context-dependent (e.g. habitat type, site conditions, and scale). However, we believe that the low numbers of reported cases for many of the management approaches is a result of practicing restoration ecologists not applying them even when necessary and appropriate for managing barriers to restoration.

There is a need for restoration ecologists to investigate all potential barriers to restoration during restoration programs. Furthermore, given that they are appropriate in their context,

we encourage restoration ecologists to apply the management actions included in this review. Also, we encourage the publication of results (even if they are negative) obtained when investigating all potential barriers to restoration and applying management actions included in this review.

Management actions were often applied individually. However, individual application of potential management actions is often ineffective because it does not address all of the factors giving rise to the target barrier to restoration, does not address all barriers to restoration present in the site designated for restoration, and/or has unintended consequences. We believe that restoration outcomes can be improved by combining potential management actions into an integrated management effort (Fig. 3).

Practicing restoration ecologists need to determine the type and extent of barriers to restoration that exist in the restoration site to inform the selection of management actions. If biological control organisms are available, classical biological control can be set up to reduce seed production and seed banks over time. If burning is appropriate, prescribed burning can be used to manage the litter. If burning is inappropriate, litter can be manually removed. Soil solarization or weed mats can be set up to reduce the establishment of secondary invaders, weedy native species, and/or a second generation of the target invader. Regardless of whether burning is applied, a combination of mowing, manual weeding, herbicides, or grazing can then be used to manage the resulting vegetation. Soil carbon addition and microbial treatments can be applied and then native species can be reintroduced, with or without the aid of nurse plants. Different combinations of approaches will be effective in different contexts (Fig. 3). For example, a combination of prescribed burning, herbicide application, and native restoration species seeding improved restoration outcomes during restoration of prairies and oak savannas in the United States (Stanley et al. 2011). There is need for further research on the efficacy of different integrated combinations of management approaches to address barriers to restoration presented by soil legacy effects of alien  $N_2$ -fixing woody species.

## Acknowledgments

Funding for this work was provided by the DST-NRF Centre of Excellence for Invasion Biology and Working for Water Program through their collaborative research project on “Integrated Management of invasive alien species in South Africa”. B.W.vW. thanks the National Research Foundation (grant 87550) for funding. We thank Mark Paschke and three anonymous reviewers for useful comments that helped to improve the manuscript.

## LITERATURE CITED

- Aerts R, Negussie A, Maes W, November E, Hermy M, Muys B (2007) Restoration of dry Afrotropical forest using pioneer shrubs as nurse-plants for *Olea europaea* ssp. *cuspidata*. *Restoration Ecology* 15:129–138
- Alpert P, Maron JL (2000) Carbon addition as a countermeasure against biological invasion by plants. *Biological Invasions* 2:33–40
- Baasch A, Kirmer A, Tischew S (2012) Nine years of vegetation development in a postmining site: effects of spontaneous and assisted site recovery. *Journal of Applied Ecology* 49:251–260
- Baer SG, Blair JM, Collins SL, Knapp AK (2003) Soil resources regulate productivity and diversity in newly established tallgrass prairie. *Ecology* 84:724–735
- Blumenthal DM, Jordan NR, Russelle MP (2003) Soil carbon addition controls weeds and facilitates prairie restoration. *Ecological Applications* 13:605–615
- Boudiaf I, Baudoin E, Sanguin H, Beddier A, Thioulouse J, Galiana A, Prin Y, Le Roux C, Lebrun M, Duponnois R (2013) The exotic legume tree species, *Acacia mearnsii*, alters microbial soil functionalities and the early development of a native tree species, *Quercus suber*, in North Africa. *Soil Biology and Biochemistry* 65:172–179
- Buckley YM, Rees M, Paynter Q, Lonsdale M (2004) Modelling integrated weed management of an invasive shrub in tropical Australia. *Journal of Applied Ecology* 41:547–560
- Callaway RM, Ridenour WM (2004) Novel weapons: invasive success and the evolution of increased competitive ability. *Frontiers in Ecology and the Environment* 2:436–443
- Campbell PL, Naudé DC, Kluge RL (1999) Development of integrated control strategies for wattle. II. Control of unwanted wattle. *South African Journal of Plant and Soil* 16:31–37
- Carrick PJ, Krüger R (2007) Restoring degraded landscapes in lowland Namaqualand: lessons from the mining experience and from regional ecological dynamics. *Journal of Arid Environments* 70:767–781
- Castro J, Zamora R, Hódar JA, Gómez JM (2002) Use of shrubs as nurse plants: a new technique for reforestation in Mediterranean Mountains. *Restoration Ecology* 10:297–305
- Cohen O, Riov J, Katan J, Gamliel A, Bar P (2008) Reducing persistent seed banks of invasive plants by soil solarization—the case of *Acacia saligna*. *Weed Science* 56:860–865
- Corbin JD, D’Antonio CM (2012) Gone but not forgotten? Invasive plants’ legacies on community and ecosystem properties. *Invasive Plant Science and Management* 5:117–124
- Craig M, Buckley GP (2013) Responses of two woodland geophytes to disturbance caused by soil translocation. *Plant Ecology* 214:1091–1103
- D’Antonio C, Meyerson LA (2002) Exotic plant species as problems and solutions in ecological restoration: a synthesis. *Restoration Ecology* 10:703–713
- Dee JR, Thomas SM, Thompson SD, Palmer MW (2016) Long-term late season mowing maintains diversity in southern U.S. tall grass prairie invaded by *Bothriochloa ischaemum*. *Applied Vegetation Science* 19:442–453
- Dijkstra FA, Wrage K, Hobbie SE, Reich PB (2006) Tree patches show greater N losses but maintain higher soil N availability than grassland patches in a frequently burned oak savanna. *Ecosystems* 9:441–452
- DiTomaso JM, Brooks ML, Allen EB, Minnich R, Rice PM, Kyser GB (2006) Control of invasive weeds with prescribed burning. *Weed Technology* 20:535–548
- Dunn PH, DeBano LF, Eberlein GE (1979) Effects of burning on chaparral soils: II. Soil microbes and nitrogen mineralization. *Soil Science Society of America Journal* 43:509–514
- Ehrenfeld JG (2003) Effects of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* 6:503–523
- Elgersma KJ, Ehrenfeld JG, Yu S, Vor T (2011) Legacy effects overwhelm the short-term effects of exotic plant invasion and restoration on soil microbial community structure, enzyme activities, and nitrogen cycling. *Oecologia* 167:733–745
- Fenn ME, Poth MA, Aber JD, Baron JS, Bormann BT, Johnson DW, Lemly AD, McNulty SG, Ryan DF, Stottlemeyer R (1998) Nitrogen excess in north American ecosystems: predisposing factors, ecosystem responses, and management strategies. *Ecological Applications* 8:706–733
- Ferreira MC, Vieira DL (2017) Topsoil for restoration: resprouting of root fragments and germination of pioneers trigger tropical dry forest regeneration. *Ecological Engineering* 103:1–12

- Fill JM, Forsyth GG, Kritzing-Klopper S, Le Maitre DC, van Wilgen BW (2017) An assessment of the effectiveness of a long-term ecosystem restoration project in a fynbos shrubland catchment in South Africa. *Journal of Environmental Management* 185:1–10
- Frostegård Å, Tunlid A, Bååth E (2011) Use and misuse of PLFA measurements in soils. *Soil Biology and Biochemistry* 43:1621–1625
- Gómez-Aparicio L (2009) The role of plant interactions in the restoration of degraded ecosystems: a meta-analysis across life-forms and ecosystems. *Journal of Ecology* 97:1202–1214
- Gómez-Ruiz P, Lindig-Cisneros R, Vargas-Ríos O (2013) Facilitation among plants: a strategy for the ecological restoration of the high-andean forest (Bogotá, DC-Colombia). *Ecological Engineering* 57:267–275
- Gooden B, French K, Turner PJ (2009) Invasion and management of a woody plant, *Lantana camara* L., alters vegetation diversity within wet sclerophyll forest in southeastern Australia. *Forest Ecology and Management* 257:960–967
- Haubensak KA, D'Antonio CM, Alexander J (2004) Effects of nitrogen-fixing shrubs in Washington and coastal California. *Weed Technology* 18:1475–1479
- Hobbs RJ, Humphries SE (1995) An integrated approach to the ecology and management of plant invasions. *Conservation Biology* 9:761–770
- Holmes PM, Cowling RM (1997) The effects of invasion by *Acacia saligna* on the guild structure and regeneration capabilities of South African fynbos shrublands. *Journal of Applied Ecology* 34:317–332
- Holmes PM, Richardson DM (1999) Protocols for restoration based on recruitment dynamics, community structure, and ecosystem function: perspectives from south African fynbos. *Restoration Ecology* 7:215–230
- Honnay O, Bossuyt B, Verheyen K, Butaye J, Jacquemyn H, Hermy M (2002) Ecological perspectives for the restoration of plant communities in European temperate forests. *Biodiversity and Conservation* 11:213–242
- Inderjit, Van Der Putten WH (2010) Impacts of soil microbial communities on exotic plant invasions. *Trends in Ecology and Evolution* 25:512–519
- Kiehl K, Thormann A, Pfadenhauer J (2006) Evaluation of initial restoration measures during the restoration of calcareous grasslands on former arable fields. *Restoration Ecology* 14:148–156
- Kiehl K, Kirmer A, Donath TW, Rasran L, Hölzel N (2010) Species introduction in restoration projects—evaluation of different techniques for the establishment of semi-natural grasslands in central and northwestern Europe. *Basic and Applied Ecology* 11:285–299
- Klimeš L, Jongepierová I, Doležal J, Klimešová J (2010) Restoration of a species-rich meadow on arable land by transferring meadow blocks. *Applied Vegetation Science* 13:403–411
- Krupek A, Gaertner M, Holmes PM, Esler KJ (2016) Assessment of post-burn removal methods for *Acacia saligna* in cape flats sand fynbos, with consideration of indigenous plant recovery. *South African Journal of Botany* 105:211–217
- Kulmatiski A (2011) Changing soils to manage plant communities: activated carbon as a restoration tool in ex-arable fields. *Restoration Ecology* 19:102–110
- Le Maitre DC, Gaertner M, Marchante E, Ens E-J, Holmes PM, Pauchard A, et al. (2011) Impacts of invasive Australian acacias: implications for management and restoration. *Diversity and Distributions* 17:1015–1029
- Levine JM, Vilà M, D'Antonio CM, Dukes JS, Grigulis K, Lavelle S (2003) Mechanisms underlying the impacts of exotic plant invasions. *Proceedings of the Royal Society of London B: Biological Sciences* 270:775–781
- Lithgow D, Martínez ML, Gallego-Fernández JB, Hesp PA, Flores P, Gachuz S, Rodríguez-Revelo N, Jiménez-Orocio O, Mendoza-González G, Álvarez-Molina LL (2013) Linking restoration ecology with coastal dune restoration. *Geomorphology* 199:214–224
- Malcolm GM, Bush DS, Rice SK (2008) Soil nitrogen conditions approach Preinvasion levels following restoration of nitrogen-fixing black locust (*Robinia pseudoacacia*) stands in a pine-oak ecosystem. *Restoration Ecology* 16:70–78
- Manchester SJ, McNally S, Treweek JR, Sparks TH, Mountford JO (1999) The cost and practicality of techniques for the reversion of arable land to lowland wet grassland—an experimental study and review. *Journal of Environmental Management* 55:91–109
- Mandle L, Bufford JL, Schmidt IB, Daehler CC (2011) Woody exotic plant invasions and fire: reciprocal impacts and consequences for native ecosystems. *Biological Invasions* 13:1815–1827
- Marchante E, Kjølter A, Struwe S, Freitas H (2009) Soil recovery after removal of the N<sub>2</sub>-fixing invasive *Acacia longifolia*: consequences for ecosystem restoration. *Biological Invasions* 11:813–823
- Maron JL, Connors PG (1996) A native nitrogen-fixing shrub facilitates weed invasion. *Oecologia* 105:302–312
- Maron JL, Jefferies RL (2001) Restoring enriched grasslands: effects of mowing on species richness, productivity, and nitrogen retention. *Ecological Applications* 11:1088–1100
- Milchunas DG, Vandever MW, Ball LO, Hyberg S (2011) Allelopathic cover crop prior to seeding is more important than subsequent grazing/mowing in grassland establishment. *Rangeland Ecology & Management* 64:291–300
- Mitchell RJ, Auld MHD, Hughes JM, Marrs RH (2000) Estimates of nutrient removal during heathland restoration on successional sites in Dorset, southern England. *Biological Conservation* 95:233–246
- Moran VC, Hoffmann JH (2012) Conservation of the fynbos biome in the Cape Floral Region: the role of biological control in the management of invasive alien trees. *BioControl* 57:139–149
- Neill C, Wheeler MM, Loucks E, Weiler A, Von Holle B, Pelikan M, Chase T (2015) Influence of soil properties on coastal sandplain grassland establishment on former agricultural fields. *Restoration Ecology* 23:531–538
- Noble IR, Weiss PW (1989) Movement and modelling of buried seed of the invasive perennial *Chrysanthemoides monilifera* in coastal dunes and biological control. *Austral Ecology* 14:55–64
- Ödman AM, Schnoor TK, Ripa J, Olsson PA (2012) Soil disturbance as a restoration measure in dry sandy grasslands. *Biodiversity and Conservation* 21:1921–1935
- Oneto SR, Kyser GB, DiTomaso JM (2010) Efficacy of mechanical and herbicide control methods for scotch broom (*Cytisus scoparius*) and cost analysis of chemical control options. *Invasive Plant Science and Management* 3:421–428
- Padilla FM, Pugnaire FI (2006) The role of nurse plants in the restoration of degraded environments. *Frontiers in Ecology and the Environment* 4:196–202
- Palmer MA, Ambrose RF, Poff NL (1997) Ecological theory and community restoration ecology. *Restoration Ecology* 5:291–300
- Pausas JG, Bladé C, Valdecantos A, Seva JP, Fuentes D, Alloza JA, Vilagrosa A, Bautista S, Cortina J, Vallejo R (2004) Pines and oaks in the restoration of Mediterranean landscapes of Spain: new perspectives for an old practice—a review. *Plant Ecology* 171:209–220
- Paz Esquivias MP, Zunzunegui M, Barradas MCD, Álvarez-Cansino L (2015) Competitive effect of a native-invasive species on a threatened shrub in a Mediterranean dune system. *Oecologia* 177:133–146
- Pearson DE, Ortega YK, Runyon JB, Butler JL (2016) Secondary invasion: the bane of weed management. *Biological Conservation* 197:8–17
- Perkins LB, Hatfield G (2016) Can commercial soil microbial treatments remediate plant–soil feedbacks to improve restoration seedling performance? *Restoration Ecology* 24:194–201
- Perry LG, Blumenthal DM, Monaco TA, Paschke MW, Redente EF (2010) Immobilizing nitrogen to control plant invasion. *Oecologia* 163:13–24
- Pickart AJ, Theiss KC, Stauffer HB, Olsen GT (1998) Yellow bush lupine invasion in northern California coastal dunes. II. Mechanical restoration techniques. *Restoration Ecology* 6:69–74
- Prober SM, Thiele KR (2005) Restoring Australia's temperate grasslands and grassy woodlands: integrating function and diversity. *Ecological Management and Restoration* 6:16–27
- Prober SM, Thiele KR, Lunt ID, Koen TB (2005) Restoring ecological function in temperate grassy woodlands: manipulating soil nutrients, exotic annuals and native perennial grasses through carbon supplements and spring burns. *Journal of Applied Ecology* 42:1073–1085

- Pyšek P, Richardson DM (2007) Traits associated with invasiveness in alien plants: where do we stand? Pages 97–125. In: Nentwig W (ed) *Biological invasions*. Springer, New York
- Rice SK, Westerman B, Federici R (2004) Impacts of the exotic, nitrogen-fixing black locust (*Robinia pseudoacacia*) on nitrogen-cycling in a pine-oak ecosystem. *Plant Ecology* 174:97–107
- Richardson DM, Kluge RL (2008) Seed banks of invasive Australian acacia species in South Africa: role in invasiveness and options for management. *Perspectives in Plant Ecology, Evolution and Systematics* 10: 16–177
- Richardson DM, Rejmánek M (2011) Trees and shrubs as invasive alien species – a global review. *Diversity and Distributions* 17:788–809
- Richardson DM, Carruthers J, Hui C, Impson FAC, Miller JT, Robertson MP, Rouget M, Le Roux JJ, Wilson JR (2011) Human-mediated introductions of Australian acacias – a global experiment in biogeography. *Diversity and Distributions* 17:771–787
- Richter BS, Stutz JC (2002) Mycorrhizal inoculation of big sacaton: implications for grassland restoration of abandoned agricultural fields. *Restoration Ecology* 10:607–616
- Riggan PJ, Lockwood RN, Jacks PM, Colver CG, Weirich F, DeBano LF, Brass JA (1994) Effects of fire severity on nitrate mobilization in watersheds subject to chronic atmospheric deposition. *Environmental Science and Technology* 28:369–375
- Rokich DP, Dixon KW, Sivasithamparam K, Meney KA (2000) Topsoil handling and storage effects on woodland restoration in Western Australia. *Restoration Ecology* 8:196–208
- Rokich DP, Dixon KW, Sivasithamparam K, Meney KA (2002) Smoke, mulch, and seed broadcasting effects on woodland restoration in Western Australia. *Restoration Ecology* 10:185–194
- Schaefer VH (2011) Remembering our roots: a possible connection between loss of ecological memory, alien invasions and ecological restoration. *Urban Ecosystems* 14:35–44
- Shortt KB, Vamosi SM (2012) A review of the biology of the weedy Siberian peashrub, *Caragana arborescens*, with an emphasis on its potential effects in North America. *Botanical Studies* 53:1–8
- Souza-Alonso P, Lorenzo P, Rubido-Bará M, González L (2013) Effectiveness of management strategies in *Acacia dealbata* link invasion, native vegetation and soil microbial community responses. *Forest Ecology and Management* 304:464–472
- Stanley AG, Kaye TN, Dunwiddie PW (2011) Multiple treatment combinations and seed addition increase abundance and diversity of native plants in Pacific northwest prairies. *Ecological Restoration* 29:35–44
- Suding KN, Gross KL, Houseman GR (2004) Alternative states and positive feedbacks in restoration ecology. *Trends in Ecology and Evolution* 19:46–53
- Szítár K, Ónodi G, Somay L, Pándi I, Kucs P, Kröel-Dulay G (2016) Contrasting effects of land use legacies on grassland restoration in burnt pine plantations. *Biological Conservation* 201:356–362
- Tanner RA, Gange AC (2013) The impact of two non-native plant species on native flora performance: potential implications for habitat restoration. *Plant Ecology* 214:423–432
- Török K, Szítár K, Halassy M, Szabo R, Szili-Kovács T, Barath N, Paschke MW (2014) Long-term outcome of nitrogen immobilization to restore endemic sand grassland in Hungary. *Journal of Applied Ecology* 51:756–765
- Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V, Maron JL, Pergl J, Schaffner U, Sun Y, Pyšek P (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecology Letters* 14:702–708
- Vitousek PM, Walker LR (1989) Biological invasion by *Myrica faya* in Hawai'i: plant demography, nitrogen fixation, ecosystem effects. *Ecological Monographs* 59:247–265
- Vitousek PM, D'Antonio CM, Loope LL, Rejmánek M, Westbrooks R (1997) Introduced species: a significant component of human-caused global change. *New Zealand Journal of Ecology* 21:1–16
- Von Holle B, Neill C, Largay EF, Budreski KA, Ozimec B, Clark SA, Lee K (2013) Ecosystem legacy of the introduced N<sub>2</sub>-fixing tree *Robinia pseudoacacia* in a coastal forest. *Oecologia* 172:915–924
- Wittenberg R, Cock MJW (2005) Best practices for the prevention and management of invasive alien species. Pages 209–232. In: Mooney HA, Mack RN, McNeely JA, Neville LE, Schei PJ, Waage JK (eds) *Invasive alien species. A new synthesis*. Island Press, Washington, DC
- Yelenik SG, Stock WD, Richardson DM (2004) Ecosystem level impacts of invasive *Acacia saligna* in the South African fynbos. *Restoration Ecology* 12:44–51
- Zedler JB (2000) Progress in wetland restoration ecology. *Trends in Ecology and Evolution* 15:402–407
- Zink TA, Allen MF (1998) The effects of organic amendments on the restoration of a disturbed coastal sage scrub habitat. *Restoration Ecology* 6:52–58. <http://www.webofknowledge.com> (accessed 28 Feb 2017)

## Supporting Information

The following information may be found in the online version of this article:

**Appendix S1.** List of studies included in the review.

**Appendix S2.** Barriers to restoration presented by soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species.

**Appendix S3.** Available management actions for the barriers to restoration presented by soil legacy effects of invasive alien N<sub>2</sub>-fixing woody species.

Coordinating Editor: Mark Paschke

Received: 10 January, 2017; First decision: 15 February, 2017; Revised: 4 December, 2017; Accepted: 6 December, 2017; First published online: 11 January, 2018