

Lake Taupo long-term monitoring programme 2008-2009

Prepared by:
Max Gibbs, NIWA Hamilton

For:
Environment Waikato
PO Box 4010
HAMILTON EAST

July 2010

Peer reviewed by:
Bill Vant Date July 2010

Approved for release by:
Dominique Noiton Date July 2010

Disclaimer

This technical report has been prepared for the use of Waikato Regional Council as a reference document and as such does not constitute council's policy.

Council requests that if excerpts or inferences are drawn from this document for further use by individuals or organisations, due care should be taken to ensure that the appropriate context has been preserved, and is accurately reflected and referenced in any subsequent spoken or written communication.

While Waikato Regional Council has exercised all reasonable skill and care in controlling the contents of this report, council accepts no liability in contract, tort or otherwise, for any loss, damage, injury or expense (whether direct, indirect or consequential) arising out of the provision of this information or its use by you or any other party.

Lake Taupo Long-Term Monitoring Programme 2008-2009

**NIWA Client Report: HAM2010-026
March 2010**

NIWA Project: EVW10210

Lake Taupo Long-Term Monitoring Programme 2008-2009

Max Gibbs

Prepared for

**Waikato Regional Council
(Environment Waikato)**

NIWA Client Report: HAM2010-026
March 2010

NIWA Project: EVW10210

National Institute of Water & Atmospheric Research Ltd
Gate 10, Silverdale Road, Hamilton
P O Box 11115, Hamilton, New Zealand
Phone +64-7-856 7026, Fax +64-7-856 0151
www.niwa.co.nz

Contents

Executive Summary	iv
Glossary	1
1. Introduction	2
2. Methods	4
2.1 Report contents	5
2.2 Statistical evaluation	5
2.3 "TREND" definition	5
3. Results and discussion	6
3.1 Temperature and dissolved oxygen	6
3.2 VHOD rate	7
3.3 Secchi depth	10
3.4 Phytoplankton	12
3.5 Algal species abundance	15
3.6 Nutrients in the upper waters	16
3.7 Nutrient accumulation in the hypolimnion	18
3.7.1 Total mass accumulated	18
3.7.2 Net accumulation rate	19
3.7.3 Total N	21
4. Knowledge gaps	22
5. Summary	23
6. Acknowledgments	26
7. References	27
8. Appendices	29
8.1 Appendix 1: Site map, sampling strategy and methods	29
8.1.1 Methods	30
8.1.2 Data handling and less than detection limit values	32
8.1.3 Statistical methods	34
8.2 Appendix 2: The calculation of net VHOD rates	35
8.3 Appendix 3: Temperature and dissolved Oxygen data	39
8.4 Appendix 4: Nutrient data	61
8.5 Appendix 5: Phytoplankton data	87
8.6 Appendix 6: Historical data	99

Reviewed by:




Dr Piet Verburg

Approved for release by:



Dr Ngaire Phillips

Formatting checked



Executive Summary

With the expectation that the trophic status of Lake Taupo will slowly change to reflect changes in land use within the lake's catchments, a long term programme monitoring the lake's water quality was commissioned by Environment Waikato. This programme commenced in October 1994 and is conducted by NIWA with field assistance from the Department of Internal Affairs, Taupo Harbourmaster's Office.

The monitoring programme was designed to detect change through assessment of the rate of consumption of oxygen from the bottom waters of the lake (volumetric hypolimnetic oxygen depletion – VHOD) as an integration of all biological processes occurring in Lake Taupo. Additional parameters are measured to provide a more comprehensive picture of water quality. Recently it has become apparent that VHOD may be too coarse to determine trophic change in a lake the size and complexity of Lake Taupo. Consequently, more emphasis is now focused on the parameters 'phytoplankton biomass', 'water clarity', and nutrient (particularly nitrate) accumulation in the lake.

The long-term monitoring programme uses the historical mid-lake site, Site A. Monitoring of additional sites in the Kuratau Basin (Site B) and the Western Bays (Site C) between January 2002 and December 2004 determined that spatial variability of water quality across Lake Taupo is minimal and that it is valid to use the mid-lake site as representative of the open water quality of the lake. Further validation of the use of a single mid-lake monitoring site was obtained from a comparison of upper water column nutrient and chlorophyll *a* concentrations and algal enumeration between Site A and near-shore sites in Whangamata Bay (Kinloch) and Whakaipo Bay, over a two-year period from February 2007 up to June 2009. That study determined that "the near-shore water quality was very similar to the mid-lake water quality" and that "within this similarity in the measured data was much variability which may be due to short period time lags between the near-shore and mid-lake sites with respect to nutrient sources, and the zones of algal growth". This report presents the results from the 2008/09 monitoring period at the mid-lake site, Site A.

There is a long-term trend of increasing phytoplankton biomass (chlorophyll *a*) in the upper 10 m of water column over the monitoring period of $0.025 \pm 0.017 \text{ mg m}^{-3} \text{ y}^{-1}$. The winter 2008 chlorophyll *a* concentrations, at 3.0 mg m^{-3} , were the highest on record for Lake Taupo.

As the long-term data accumulates, it has become apparent that the increase in chlorophyll *a* data may not be a linear trend and could be part of a long-term cycle. The annual mean chlorophyll *a* data from 1994 to 2003 was increasing at a statistically significant rate of $0.087 \pm 0.029 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.001$, $r^2 = 0.857$, $n = 10$), but since 2000 there has been a statistically significant trend of decline at a rate of $0.033 \pm 0.031 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.05$, $r^2 = 0.423$, $n = 10$).

Highest phytoplankton biomass occurred in August 2008 when the lake had mixed and lowest biomass occurred in the upper water column in early summer 2009 when that winter biomass peak had settled from the water column onto the lake bed. Chlorophyll fluorescence profiles show that, each year, a deep chlorophyll maximum (DCM) develops just below the thermocline (40 m) during summer and has up to 70% higher biomass than the upper water column.

The 2008 winter bloom was initially dominated by the diatoms *Asterionella formosa* and *Aulacoseira granulata* with *Fragilaria crotonensis* and a green algae, *Monoraphidium* sp., becoming dominant through spring. The colonial green, *Botryococcus braunii*, and some large dinoflagellates, *Gymnodinium* sp. and *Peridinium* sp., become dominant in summer and autumn 2009. Cyanobacteria (blue-green algae) were always present in low numbers in the upper water column throughout the 2008/09 monitoring period, with *Anabaena lemmermannii* being the most common species. There was an ever present background of small (<5 µm) unicellular flagellates throughout the year.

Nutrient concentrations – dissolved reactive phosphorus, ammoniacal nitrogen, and nitrate nitrogen (DRP, NH₄-N, and NO₃-N) – in the upper water column were comparable with concentrations measured since 2003. NO₃-N concentrations were lower and NH₄-N concentrations were elevated in the upper water column since 2007. The elevated NH₄-N concentrations may indicate water column decomposition of the winter-spring bloom, or excretion from a zooplankton bloom.

The total mass of NO₃-N in the hypolimnion before winter has increased at a statistically significant rate of about 7.9 t y⁻¹ ($P < 0.001$, $r^2 = 0.41$, $n = 22$) over the last 34 years. This value is the same as the previous year but includes an increase of around 90 t of NO₃-N in the hypolimnion in autumn compared with autumn the previous year. The total mass of NO₃-N in the hypolimnion in autumn 2009 was 470 t. The net accumulation rate of NO₃-N in the hypolimnion below 70 m for the 2008/09 stratified period was 1.93 t d⁻¹, which is a 20% increase over the previous year. However, because of high variability in the data, regression analysis indicates that the trend of increase in the net hypolimnetic NO₃-N accumulation rate during the stratified period is only weakly significant at 0.028 t d⁻¹ ($P = 0.07$, $r^2 = 0.154$, $n = 22$) over the last 34 years.

Spring and summer 2008/09 water clarity was high at 22 m but was not as high as the previous summer which at 25 m in February 2008, was the highest clarity on record for Lake Taupo. These extremely clear-water phases both coincided with periods of drought (declared officially summer 2008) and thus may reflect the reduced nutrient input in surface runoff as well as a low input of sediment from erosion.

Lowest water clarity in winter did not coincide with the peak of chlorophyll production in August 2008, rather, lowest water clarity occurred in November 2008 during a wet and windy phase and clarity remained low until the weather calmed down again in February 2009.

The 2008/09 net VHOD rate at $17.50 \pm 3.64 \text{ mg O}_2 \text{ m}^{-3} \text{ d}^{-1}$ (mean \pm 95% confidence limit) was almost 3 $\text{mg O}_2 \text{ m}^{-3} \text{ d}^{-1}$ higher than the previous year which was $14.51 \pm 2.94 \text{ mg O}_2 \text{ m}^{-3} \text{ d}^{-1}$. Evaluation of the VHOD data shows that there has been a statistically significant ($P < 0.002$, $r^2 = 0.69$, $n = 11$) increase of around $1.04 \text{ mg m}^{-3} \text{ d}^{-1}$ in the VHOD rate each year since the low of 1999. The period of this regression is selected from lowest to the present value, which is the highest measured during the monitoring programme, and thus does not reflect a long term trend in Lake Taupo. The persistent increase in hypolimnetic oxygen demand over the past 11 years does, however, imply a change in the carbon load on the lake.

In the 2002 review of the long-term monitoring programme data, 3 trends in the data were identified – increasing phytoplankton biomass in the upper 10 m, increasing $\text{NO}_3\text{-N}$ mass in the hypolimnion prior to winter mixing, and an increasing range in the variability of water clarity – that were of concern with respect to the water quality of Lake Taupo. In previous reports, it was also shown that the net accumulation rate of $\text{NO}_3\text{-N}$ in the hypolimnion during the stratified period has increased over the last 34 years. These trends are still present in the data to 2009.

Glossary

BOD	Biochemical Oxygen Demand: the rate of oxygen consumption associated with biological decomposition and chemical processes in the water column.
VHOD	Volumetric Hypolimnetic Oxygen Demand: the net rate of oxygen loss associated with biological, chemical and physical processes in the hypolimnion of a lake in the absence of a temperature change
Phytoplankton	Microscopic free-floating aquatic plants (algae)
Cyanobacteria	Blue-green algae. These are potentially toxic. They can adjust their depth in the water column using small gas bladders (gas vacuoles), and some species can use (i.e., fix) atmospheric nitrogen for growth when nutrient nitrogen in the water column is depleted.
Zooplankton	Small to microscopic free-swimming aquatic animals which graze on phytoplankton or smaller zooplankton
Biomass	The living mass of the phytoplankton or zooplankton populations
Thermal stratification	Separation of a water column into two layers by temperature – warmer water on top
Thermocline	The boundary zone or temperature gradient between the two layers in a thermally stratified water column.
Epilimnion	The upper water column in a thermally stratified water column
Hypolimnion	The lower water column in a thermally stratified water column
Metalimnion	The thermocline zone – of variable thickness
Euphotic zone	The upper water column in which there is sufficient light for photosynthesis and hence phytoplankton growth.
Euphotic depth	Lower limit of phytoplankton growth where light levels are 1% of surface irradiance
Nutrients	Essential dissolved inorganic nitrogen and phosphorus compounds which can be used directly by plants for growth
Ammoniacal nitrogen	Sum of ammonium ion (NH_4^+) plus free (unionised) ammonia (NH_3). Some amines (NH_2^-) may be included as interference during analysis. Symbol, $\text{NH}_4\text{-N}$.
Nitrate nitrogen	Used in this report as the sum of nitrate (NO_3^-) plus nitrite (NO_2^-). Symbol, $\text{NO}_3\text{-N}$.
DIN	Dissolved Inorganic Nitrogen: the sum of $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$
DON	Dissolved Organic Nitrogen: the soluble nitrogen other than DIN
PN	Particulate Nitrogen: includes phytoplankton and other detritus
TN	Total Nitrogen: Sum of DIN + DON + PN
NO_x	Gaseous oxides of nitrogen, including N_2O , NO, NO_2

1. Introduction

A long term monitoring programme of Lake Taupo's water quality was initiated by Environment Waikato in October 1994 with the expectation that the trophic status of the lake would slowly change to reflect changes in land use within the lake's catchment. This programme is conducted by NIWA with field assistance from the Department of Internal Affairs, Taupo Harbourmaster's Office. Various additions and improvements to the monitoring methodology have occurred with advances in available technology but the core monitoring parameters remain unchanged (Appendix 1). This report presents data from the routine mid-lake monitoring station from August 2008 to July 2009. Additional information for water clarity, temperature, and chlorophyll *a* collected between August 2009 and the time of writing this report have also been included in the data sets in the appendices.

In two earlier reports (Gibbs 2005, 2006), data were included from two additional sites representing those historically measured in the 1974-76 assessment of lake water quality (White et al. 1980) (Fig.1) to evaluate spatial variability of water quality across the lake. Results from these two additional sites showed that, in general, there was minimal variation between the sites spatially or with season and that data collected from site A (mid lake) could be used as being representative of the main body of the lake. More recently, a comparison of upper water column nutrient and chlorophyll *a* concentrations and algal enumeration was made between Site A and near-shore sites in Whangamata Bay (Kinloch) and Whakaipo Bay (Fig. 1), over a two-year period from February 2007 up to June 2009 (Gibbs 2010). That study determined that "the near-shore water quality was very similar to the mid-lake water quality" and that "within this similarity in the measured data was much variability which may be due to short period time lags between the near-shore and mid-lake sites with respect to nutrient sources, and the zones of algal growth". This report presents data from site A only.

The monitoring programme has three components: bottom water oxygen depletion, upper water column water quality, and whole water column water quality. Bottom water oxygen depletion is estimated as the volumetric hypolimnetic oxygen depletion (VHOD) rate, which is sensitive to changes in trophic status of lakes that thermally stratify for part of the year (Burns 1995). VHOD was chosen as the parameter most likely to detect a change in the water quality of Lake Taupo. Estimates of VHOD are made from dissolved oxygen and temperature profiles measured at 2-3 week intervals during the stratified period. However, the VHOD rate will only indicate that a change has occurred and will not detect other indicators of changing water quality. For that purpose, the upper water column is sampled for nutrients, chlorophyll *a*, phytoplankton species composition and water clarity at 2-3 weekly intervals, and the whole lake water quality is estimated from full profiles which are made twice during

the stratified period. The first profile is in spring, when thermal stratification has become established and is stable. The second profile is in autumn the following year before thermal stratification begins to break down, and the thermocline deepens.




Figure 1: Site map of Lake Taupo showing location of the routine monitoring site at mid lake (A), and the two additional sites at Kuratau Basin (B) and the Western Bays (C) sampled during the three-year period 2002-04. The near-shore comparison sites at Whangamata Bay and Whakaipo Bay sampled during a two-year period 2007-09 are also shown. The generally close agreement between most data from the three deep lake sites and the near-shore sites with site A gives confidence in the past and future use of the site A data as representing the water quality of the whole lake.

2. Methods

Detailed method descriptions are given in Appendix 1. The parameters measured routinely at 2-3 weekly intervals are:

- depth-related temperature and dissolved oxygen (DO), using the RBR XR420f CTD profiler until January 2008, then using the RBR XR620f CTD profiling system. Additional parameters of conductivity and chlorophyll fluorescence, and since January 2008, PAR, recorded by the profiler sensors are available at NIWA and will only be reported as appropriate;
- water clarity by Secchi disc depth (20-cm black and white quartered);
- chlorophyll *a*, nitrate+nitrite-nitrogen ($\text{NO}_3\text{-N}$), ammoniacal-N ($\text{NH}_4\text{-N}$), dissolved organic N (DON), particulate-N (PN), total nitrogen (TN), dissolved reactive phosphorus (DRP), dissolved organic phosphorus (DOP), particulate phosphorus (PP), total phosphorus (TP), and algal species dominance in integrated-tube water samples from the top 10 m. Zooplankton net hauls from 100 m (63 μm mesh) are preserved in 4% formalin and stored pending analysis.

Since 2000, water samples have also been collected at the same time from just above the lake bed (150 m) for analysis of $\text{NO}_3\text{-N}$, $\text{NH}_4\text{-N}$, and DRP to assess nutrient accumulation rates in the hypolimnion and to assess the extent of winter mixing.

For whole water column sampling, the parameters measured at 10 m depth intervals from the surface to the bottom of the lake twice a year are:

- Conductivity, pH, temperature, DO, chlorophyll *a*, DRP, DOP, PP, TP, $\text{NO}_3\text{-N}$, $\text{NH}_4\text{-N}$, DON, PN, TN, urea nitrogen (Urea-N), total suspended solids (SS), volatile suspended solids (VSS), particulate carbon (PC), and dissolved organic carbon (DOC).

Additional parameters measured twice yearly, but not as complete profiles are:

- Algal species composition and abundance on water samples from 1, 10, 50, 100 and 140 m.

Details of data handling and the treatment of values that are near analytical detection limits are described in Appendix 1.

2.1 Report contents

This report presents the results from the 2008/09 stratified period plus the winter 2009 mixing, and refers to data in previous annual monitoring reports from 1995 to 2005 (e.g., Gibbs 2009; Gibbs et al. 2002) for inter-annual comparisons, and archived historical data since 1974 held by NIWA. The methods used are as per the 1994/95 report (Gibbs 1995) and a copy of these methods is included in Appendix 1. The calculation of the net VHOD rate, as applied to Lake Taupo data, was described in the 1996/97 report and a copy is presented in Appendix 2. Copies of temperature and dissolved oxygen, and nutrient data from the previous twelve years are included in Appendix 3 and 4 respectively. Graphical presentations of historical time-series temperature, dissolved oxygen, and Secchi disc depth data collected since the start of this monitoring programme are updated and presented in figures in the text. Phytoplankton species composition and dominance data for 2008/09 are included in Appendix 5 and discussed in the text. Graphical presentations of all available chlorophyll *a* concentration and Secchi depth water clarity data are presented where appropriate. Historical nitrate and dissolved reactive phosphorus data from spring and autumn full lake profiles are presented in Appendix 6 for reference.

2.2 Statistical evaluation

Simple statistical evaluation of data has been made using Microsoft Excel® and regression results have been reported to \pm 95% confidence limits. Statistical significance (P), where used, includes the coefficient of determination (r^2) and the number of data points used (n). For details see Statistical Methods, Appendix 1.

2.3 “TREND” definition

As in previous reports, the word “trend” is used in the context of a change between the start and the end of a time series data set where the use of a linear regression analysis shows a statistically significant difference from the null hypothesis of there being no change. Use of the word “trend” is a statistical one. It does not imply any valid extrapolation of the observed change beyond the period of the data set being examined by the linear regression.

3. Results and discussion

3.1 Temperature and dissolved oxygen

Depth profiles of temperature and dissolved oxygen (DO) were measured at the mid-lake site, Site A (Fig. 1), at about 2-3 weekly intervals, depending on the weather, throughout the year. These data, plus profiles measured during the full water quality samplings at Site A on 14 October 2008 (spring) and 15 April 2009 (autumn), are presented in Appendix 3. The time-series temperature and DO data from specific depths of 20 m (epilimnion) and 130 m (hypolimnion) collected in the monitoring programme since 1994 are presented in Figure 2.

Temperature and DO data indicate that Lake Taupo was well mixed in winter 2008 with the mixing period being from around 15 July to 4 September. This longer than usual mixing period produced a 0.6 °C cooling in the bottom waters from 11.2 °C to 10.6 °C (Fig. 2A). The lake was also well mixed in winter 2009 with the mixed period extending from 6 July to 7 September with a 0.6 °C drop in bottom water temperature from 10.9 °C to 10.3 °C. This is the coolest the bottom waters have been since 1998. In contrast, surface water temperatures in summer 2008/09 had an extended period above 18 °C, from early January through to the beginning of April 2009, and were the warmest since summer 1996 (Fig. 2A).




Figure 2A: Time-series temperature from 20 m (black line) and 130 m (pink line) depths. Winter mixing occurred where these two lines meet. The data show the lack of mixing in winter 1998 and only partial mixing in 1999. Mixing was brief in 1997 and 2005.

Despite years with incomplete mixing, the dissolved oxygen content of the hypolimnion has rarely fallen below 7.0 g m^{-3} , even close to the sediment (Fig. 2B). During winter mixing in 2008 and 2009 the bottom oxygen concentrations exceeded 10.5 g m^{-3} (Fig. 2B) confirming the high degree of mixing and also consistent with the colder bottom waters (Fig. 2A). Conversely, bottom oxygen concentrations at the end of summer 2008/09 were at around 7.0 g m^{-3} with oxygen concentrations immediately above the sediment at around 6.8 g m^{-3} .




Figure 2B: Time-series dissolved oxygen data from 20 m (black line) and 130 m (pink line) depths. Mixing and complete reoxygenation occurred where the two lines in the temperature data (Fig. 2A) meet each winter. This corresponds with the period that the two lines on the oxygen data should meet. However, where temperature data indicate incomplete mixing and the 2 lines of the oxygen data do not meet, there is incomplete reoxygenation of the hypolimnion. Date ticks are 1 January in each year.

3.2 VHOD rate

The VHOD rate was estimated between September 2008 and March 2009 based on oxygen profile data collected at site A. VHOD calculations were made using the volume-weighted mean oxygen concentration below 70 m on each sampling occasion (Fig. 3) – see Appendix 2 for more detail. The VHOD rate in 2008/09 was $17.50 \pm 3.64 \text{ mg m}^{-3} \text{ d}^{-1}$ (mean \pm 95% confidence limit) (Fig. 3). This value was $3 \text{ mg m}^{-3} \text{ d}^{-1}$ higher than the value for 2007/08 at $14.51 \pm 2.94 \text{ mg m}^{-3} \text{ d}^{-1}$ (Table 1). Overall, there is a statistically significant ($P < 0.002$, $r^2 = 0.69$, $n = 11$) trend of increase in the VHOD rate data of around $1 \text{ mg m}^{-3} \text{ d}^{-1}$ each year since 1999 (Fig. 4).




Figure 3: Volume-weighted mean dissolved oxygen (DO) concentrations below 70 m for 2008/09. The linear regression through the solid data points is the VHOD rate. ($P < 0.0001$, $r^2 = 0.94$, $n = 10$).




Figure 4: Time-series of VHOD rates since 1994-95. The low VHOD in 1997-2000 (following the 1995/96 eruptions of Mount Ruapehu) corresponds to a shift in algal dominance from diatoms to colonial greens (*Botryococcus braunii*). The regression through the solid (red) dots ($P < 0.002$, $r^2 = 0.69$, $n = 11$), only looks at the change in VHOD since 1998/99, the lowest VHOD rate in the long term monitoring programme, and the beginning of the return of diatoms as the dominant algal species.

The low VHOD in 1999 can be attributed to the effects of the 1995/96 eruption of Mount Ruapehu which deposited around two million tonnes of allophanic ash falling on the lake. While allophane is known to remove phosphate from water (see section 3.6: Nutrients in the upper waters) this event appears to have triggered a change in the winter bloom dominant algal species from diatoms to buoyant colonial green algal dominance (Table 1). The change from *Aulacoseira granulata*, a heavy diatom which sinks rapidly, to *Botryococcus braunii*, a large colonial green algae which floats in the upper water column, would have allowed the phytoplankton carbon to drift inshore rather than settle in the deeper parts of the lake. However, rather than just returning to the pre-eruption VHOD levels when the diatoms once more dominated the algal species in the winter bloom, the VHOD rate has continued to increase almost every year (Fig. 4). While the period of the regression analysis is selected from lowest to highest, and thus does not reflect a long-term trend in Lake Taupo, this sustained increase in VHOD over an 11-year period suggests a change in the export of organic carbon to the hypolimnion over this period, either from external inputs (i.e., land-use effects), or primary production within the lake, or a combination of both.

Table 1: Summary of the volumetric hypolimnetic oxygen depletion (VHOD) rates ($\text{mg O}_2 \text{ m}^{-3} \text{ d}^{-1}$) ($\pm 95\%$ confidence limit) and the dominant phytoplankton species during the preceding winter bloom.

Year	VHOD rate	Dominant phytoplankton species	Type
1994-95	8.93 (2.39)	<i>Aulacoseira granulata</i> *	Diatom
1995-96	9.07 (2.77)	<i>A. granulata</i>	Diatom
1996-97	5.12 (1.37)	<i>Botryococcus braunii</i>	Colonial green
1997-98	3.21 (2.03)	<i>B. braunii</i>	Colonial green
1998-99	2.64 (1.90)	<i>B. braunii</i>	Colonial green
1999-00	5.11 (1.14)	<i>B. braunii</i> + <i>A. granulata</i> + <i>Cyclotella stelligera</i>	C.G. – Diatom mix
2000-01	9.34 (2.9)	<i>A. granulata</i>	Diatom
2001-02	9.06 (2.7)	<i>Asterionella formosa</i>	Diatom
2002-03	13.76 (2.14)	<i>A. formosa</i> + <i>A. granulata</i>	Diatom
2003-04	11.50 (2.80)	<i>A. formosa</i> + <i>A. granulata</i>	Diatom
2004-05	11.30 (1.13)	<i>Fragilaria crotonensis</i> + <i>A. formosa</i>	Diatom
2005-06	9.56 (2.24)	<i>A. formosa</i> + <i>A. granulata</i>	Diatom
2006-07	10.73 (2.45)	<i>A. granulata</i>	Diatom
2007-08	14.51 (2.94)	<i>Fragilaria crotonensis</i> + <i>A. formosa</i>	Diatom
2008-09	17.50 (3.64)	<i>A. formosa</i> + <i>A. granulata</i>	Diatom

* Not measured in winter but measured in October 1994.

3.3 Secchi depth

Water clarity, as measured by Secchi depth, in Lake Taupo generally follows an annual cyclical pattern inversely correlating with the seasonal pattern of phytoplankton abundance. Secchi depths in the long-term record, until recently (since 2002), have mostly been within the range of 10 m to 20 m (Fig. 5) with lowest water clarity during the winter/spring growth phase and highest water clarity during summer when the phytoplankton have settled out of the epilimnion, which is depleted in plant growth nutrients at that time.

The water clarity in summer 2008/09 was higher than the long-term upper range, with Secchi depths reaching 22 m in February 2009 (Fig. 5A). Although not as high as the maximum of 25 m recorded in February 2008, which was attributed to a drought (Gibbs 2009), this extreme clear-water phase also coincided with another warm calm period with low runoff. Under these unusually dry conditions it is possible that water clarity increases due to the reduced nutrient inputs resulting from expected lower stream flow and groundwater inflow, as well as low input of sediment from catchment erosion.

Of interest, the water clarity in summer 2009/10, a wet and windy summer, only reached a maximum Secchi depth of 17 m (in February 2010: Fig. 5A), lending support to the concept that the lack of suspended inorganic matter may be a factor in the high water clarity in summer in dry years.

Mean water clarity during winter (July-September) has increased by $0.1 \pm 0.1 \text{ m y}^{-1}$ ($P = 0.05$, $r^2 = 0.27$, $n = 16$) since the beginning of monitoring in 1994 (Fig. 5B). However, because winter is the mixed period and there is high inter-sampling and inter-annual variability in the data, this increase has not been consistent over the whole period. Examining the data in five-yearly blocks shows that the mean winter water clarities for the periods 1995-99, 2000-04, and 2005-09 were 12.5 m, 13.0 m, and 13.5 m, respectively, but the minimum values for each period were 10 m, 9.5 m, and 11 m, respectively.

There was substantially less inter-annual variability in the winter mean over the last 5 years (2005-09). This may reflect a shift in the timing of minimum clarity into spring. For example, the average water clarity over the 2008 winter/spring period (July to December) was 12.6 m (Fig. 5A) and in the 2009 winter/spring period it was 13.9 m. In the 2008 winter/spring period, the lowest water clarity occurred in late November whereas in previous years the water clarity minimum has typically occurred in late August. Furthermore, water clarity remained low well into summer 2008/09 at 13.0 m on 13 January 2009 before suddenly increasing to 22.0 m on 11 February 2009 (Fig. 5A).




Figure 5: A) Time-series changes in water clarity as indicated by Secchi depth water clarity from summer 2005 through to winter 2009. Dotted line at 13.5 m is the mean winter water clarity for this period. Date ticks are 1st of each month. B) Long-term changes in water clarity show the variability across seasons and years for the present monitoring programme since 1994. Date ticks are 1st January each year. C) Combined Secchi depth record since 1932. The 1932 data point was from John S. Armstrong's notes after converting his value from feet to meters.

While a higher mean winter water clarity suggests there should have been lower algal biomass in winter than in previous years, the chlorophyll *a* data (Fig. 6) shows that algal biomass, at 3.0 mg m^{-3} on 7 August 2008, was the highest ever recorded during the current long-term monitoring programme. These time-series data also show that the lowest water clarity on 26 November 2008 corresponded with a chlorophyll *a* concentration of only 1.0 mg m^{-3} . As this was a wet and windy spring period, the low water clarity in November was probably due to inputs of inorganic particulate matter washed into the lake. This would be consistent with the very rapid clearing of the

water column in January 2009 when there was an extended dry, calm period with implied reduced mixing to keep that particulate matter in the upper water column.

3.4 Phytoplankton

The overall pattern of changes in chlorophyll *a* concentration since 1994 has been one of maximum concentrations during the winter algal bloom and minimum concentrations in summer and, as expected, there is a statistically significant inverse logarithmic relationship between chlorophyll *a* concentration and Secchi disk depth, as discussed in the 2004/05 report (Gibbs 2006).

While this pattern has not changed, it is apparent that the previously reported long-term trend of increasing mean and maximum chlorophyll *a* concentrations in the upper 10 m of the water column at the mid-lake site (e.g., Gibbs 2009), has changed. Previously, this trend was reported as a mean rate of increase and has a present value of $0.025 \pm 0.017 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P = 0.005$, $r^2 = 0.033$, $n = 237$) (Fig. 6), using all data, i.e., that trend is still present.




Figure 6: Time-series chlorophyll *a* concentrations in the upper 10 m of Lake Taupo at the mid-lake site, Site A. The solid regression line represents a statistically significant increase in the mean chlorophyll *a* concentrations of $0.025 \pm 0.017 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P = 0.005$, $r^2 = 0.03$, $n = 237$). Date ticks are 1 January in each year. Chlorophyll *a* detection limit = 0.1 mg m^{-3} .

However, while there was a statistically significant increase in the annual mean chlorophyll *a* concentrations of $0.087 \pm 0.029 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.001$, $r^2 = 0.857$, $n = 10$) from 1994 to 2003 (Fig. 7), there was a statistically significant decrease of $0.033 \pm 0.031 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.05$, $r^2 = 0.423$, $n = 10$) from 2000 to 2009 (Fig. 7).

A similar pattern is seen in the annual maximum chlorophyll *a* concentrations with a statistically significant increase of $0.19 \pm 0.086 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.001$, $r^2 = 0.765$, $n = 10$) from 1994 to 2003, but a non statistically significant decrease of $0.016 \pm 0.1 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P = 0.7$, $r^2 = 0.018$, $n = 10$) from 2000 to 2009 (Fig. 7). The lack of a statistically significant change across these annual maximum data is likely to be due to the high variability in the maximum chlorophyll *a* concentration each winter in recent years (Fig. 7). The mean of the maximum data was $2.49 \pm 0.51 \text{ mg m}^{-3}$.




Figure 7: Annual average and winter maximum chlorophyll *a* concentrations from the 10-m tube samples since 1994. Regression lines indicate significant ($P < 0.001$) trends of increase between 1994 and 2003 (broken lines) but trends of decrease from 2000 to 2009 (solid lines). These regressions use a data overlap between 2000 and 2003. Note the regression for the winter maxima decrease from 2000 to 2009 is not statistically significant ($P = 0.7$). Regression slopes are as given in the text. Date ticks are 1 January in each year.

There are several possible explanations for the change from an increase to a decrease, including natural variability and that the data are tracking part of a longer, natural, cycle of the lake. It could also reflect a recovery from the effects of the 1995/96 Mount Ruapehu eruption event. Inspection of all chlorophyll *a* data available since 1974 (Fig. 8) shows that prior to the 1995/96 eruption event, there were high winter chlorophyll *a* concentrations comparable with the recent winter maximum

concentrations. For example, the chlorophyll *a* concentrations of 2.45 mg m⁻³ and 2.48 mg m⁻³ recorded in August 1987 and 1989, respectively, are approaching the maximum recently recorded value of 3.0 mg m⁻³ in August 2008.




Figure 8: Time-series upper 10 m chlorophyll *a* concentrations in Lake Taupo from 1977 to present (pink). The 1974/75 data (blue) were collected by 30-m integrated tube sampler and are not directly comparable with the later data (See text). The linear regression line shows a statistically significant trend in the data from 1977 to present of $0.012 \pm 0.008 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.005$, $r^2 = 0.0265$, $n = 304$).

Regression analysis of all data from 1977 shows a statistically significant increase in the chlorophyll *a* concentration of $0.012 \pm 0.008 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.005$, $r^2 = 0.0265$, $n = 304$). This is about half the rate indicated by the regression through the data since 1994 ($0.025 \pm 0.017 \text{ mg m}^{-3} \text{ y}^{-1}$; $P = 0.005$, $r^2 = 0.033$, $n = 237$). Note that chlorophyll *a* data collected in 1974/75 were from a 30-m integrated tube sampler which would have included higher chlorophyll *a* concentrations from the deep chlorophyll maxima (DCM) circa 40 m (see below). Consequently, those data are not consistent with the 0–10 m chlorophyll *a* data from 1977 on and cannot be used in the regression analysis.

The monitoring programme uses chlorophyll *a* concentrations (extracted from water samples) as an indicator of algal biomass in the upper 10 m because surface layer chlorophyll *a* concentrations can be directly related to water clarity (Secchi depth). However, the use of the profiler fitted with a chlorophyll fluorescence sensor indicates that a large proportion of the algal biomass in Lake Taupo through spring and summer is associated with the base of the thermocline (40 m) as a DCM with a potential

biomass up to 70% greater than in the 0-to-10 m layer, as demonstrated in Gibbs (2007). The use of the 30-m integrated tube sampler in 1974/75 would have included part of the DCM accounting for the higher chlorophyll *a* concentrations at that time (Fig. 8). The DCM was present throughout the spring-summer phase of the 2008/09 stratified period with chlorophyll fluorescence values comparable with previous years.

Note that, in summer, severe fluorescence quenching by sunlight occurs above 10m with effects decreasing to 20 m. Consequently, chlorophyll *a* concentrations cannot be estimated from the upper 10 m of the fluorescence profile in summer and rely on a correction curve between 10 m and 20 m. Below 20 m chlorophyll *a* concentrations estimated by fluorescence are typically within 5% of chlorophyll *a* concentrations measured by extraction from discrete water samples.

3.5 Algal species abundance

In 2008, algal biovolume was often dominated by *Botryococcus braunii* until July although the numbers of colonies were very low. In August and September 2008, algal biovolume and abundance was co-dominated by the diatoms *Asterionella formosa* and *Aulacoseira granulata* with the diatom *Fragilaria crotonensis* becoming co-dominant with the green algae *Monoraphidium* sp. (formerly *Ankistrodesmus falcatus*) from November 2008 through January 2009. Cyanobacteria were never dominant although the main species present, *Anabaena lemmermannii*, was more abundant between February and April 2009 than at other times of the year (Appendix 5). *Botryococcus braunii* was present in low numbers throughout summer 2008/09. The large dinoflagellates *Gymnodinium* sp. and *Peridinium* sp. were also present on occasion through summer 2008/09 although there was an ever present background of small (<5 µm) unicellular flagellates throughout the year.

Samples taken from the DCM on 14 October 2008 comprised mostly species similar to those in the surface layer. The 15 April 2009 sample from 50 m contained a higher proportion of *Botryococcus braunii* than in the upper water column. The diatom, *Aulacoseira granulata* was also present in the DCM but was not found in the upper water column. Conversely, large dinoflagellates were abundant in the upper water column but were not found in the DCM. This is consistent with a calm autumn with less mixing allowing the non-motile species to settle.

A comparison of Site A algal data with near-shore algal data from Whangamata and Whakaipo Bays (Fig. 1) provides a more detailed evaluation of the seasonal changes in algal species in Lake Taupo (Gibbs, 2010).

3.6 Nutrients in the upper waters

Time-series plots of DRP (Fig. 9A), NH₄-N (Fig. 9B), NO₃-N (Fig. 9C) and DON (Fig. 9D), show that the 2008/09 values were mostly within the seasonal range measured over the whole period of the present monitoring programme since 1994. As previously noted (Gibbs 2006), nutrient concentrations changed abruptly at the time of the Mount Ruapehu eruptions in 1995 and slowly returned to pre-eruptions levels by 2003 (Fig. 9). Since 2003, fluctuations in the surface NO₃-N and NH₄-N (Fig. 9B & C) have mostly corresponded with winter mixed periods when nutrient enriched bottom waters were dispersed up through the lake's water column.

Of special interest are the relatively high DRP concentrations in the upper water column when the lake mixed in winter 2009. While a similar feature was observed in winter 2005 when the lake only mixed very briefly, in winter 2009 the lake mixed completely (Fig. 2A). This may be related to the low algal biomass in the lake at that time (Fig. 6) which may have resulted in unused DRP (and NO₃-N) remaining in the water column. The probable source of the DRP (and NO₃-N) would be from hypolimnetic accumulation during the stratified period followed by upwelling during winter mixing (Fig. 9).

Another feature of the nutrient data since the beginning of 2007 has been the relatively elevated NH₄-N concentrations in the upper water column (Fig. 9B). While NH₄-N usually appears in the surface water at the time of winter mixing and can be attributed to upwelling of nutrient-rich bottom water, these elevated NH₄-N concentrations throughout the year cannot be linked to an upwelling event. Consequently, it may be related to water column decomposition of senescent phytoplankton or excretion by zooplankton feeding on those algae. This would be consistent with the exceptionally high water clarity in the last few summers.

During the 2008/09 monitoring period, DON concentrations fell below the long-term minimum value of around 29 mg m⁻³ for the first time since the beginning of the monitoring programme. Previously it had been considered that the lake had a near constant amount of refractory organic material, the minimum value, which was augmented by labile DON from various sources.




Figure 9: Time series data from the top 10 m of water column in Lake Taupo for (A) dissolved reactive phosphorus (DRP), (B) ammoniacal nitrogen ($\text{NH}_4\text{-N}$), (C) nitrate + nitrite nitrogen ($\text{NO}_3\text{-N}$), and (D) dissolved organic nitrogen (DON). Broken line indicates the “minimum” DON concentration, which may be the concentration of refractory organic material in the lake. Date ticks are 1 January in each year.

3.7 Nutrient accumulation in the hypolimnion

Dissolved inorganic nutrients in water samples from 150 m depth demonstrated consistent seasonal patterns which more precisely indicate the time of complete mixing in winter (Fig. 10). Winter mixing was indicated by the sudden drop in DRP and NO₃-N concentrations that usually occurs around the beginning of August. In 2008 and 2009, according to temperature data (Fig. 2A), mixing began in mid to early July. Hypolimnetic NO₃-N concentrations before winter mixing have been consistent at around 32-35 mg m⁻³ since 2004, after declining from around 46 mg m⁻³ in 2001. Concentrations were slightly higher in autumn 2008 reaching around 38 mg m⁻³ before winter mixing. At each mixing period, NH₄-N is released into the bottom water (Fig. 10), but its maximum concentration has decreased from around 9 mg m⁻³ in 2001 to around 3-4 mg m⁻³ in winter 2008. The source of this NH₄-N is not known.




Figure 10: Time series data for DRP (blue square), NO₃-N (red diamond) and NH₄-N (black triangle) in the bottom waters (150 m depth) of Lake Taupo since winter mixing 2003.

3.7.1 Total mass accumulated

The total mass¹ of NO₃-N in the hypolimnion in autumn each year before winter mixing has ranged from about 120 t (1978) to more than 650 t (2000) (Fig. 11). While this graph is similar to those in earlier reports, this graph also includes all additional information from historical data sets held by NIWA. A copy of the historical source data used to produce the additional data points from 1988 to 1990 is included in Appendix 6. Since 1975 there has been a statistically significant ($P < 0.001$, $r^2 = 0.41$, $n = 22$) long-term trend of increase in the total mass of NO₃-N in the hypolimnion before winter mixing of around 7.9 t y⁻¹ (Fig. 11). The total mass of NO₃-N in the

¹ In earlier reports the total mass of NO₃-N in the hypolimnion each year has been referred to as the “total accumulated mass” of NO₃-N. It is the “standing stock” of NO₃-N at that time.

hypolimnion in April 2009 was around 470 t, an increase of around 90 t since 2008, with the data point falling on the regression line.




Figure 11: Long-term time series of total mass of NO₃-N in the hypolimnion of Lake Taupo in autumn before winter mixing. The regression line indicates a statistically significant trend of increase in the total mass of 7.9 t y^{-1} ($P < 0.001$, $r^2 = 0.41$, $n = 22$). Open circle data excluded from regression as time periods and depth data were not the same as used for the other data. Date ticks are 1 January in each year.

3.7.2 Net accumulation rate

While the total mass of NO₃-N in the hypolimnion is a “standing stock” before winter mixing, it is recognised that the standing stock is a function of the mass of NO₃-N in the hypolimnion at the beginning of the stratified period plus the net mass that was released from the sediments and accumulated in the hypolimnion during the stratified period. The difference between the standing stock at the beginning and end of the stratified period is the net mass of NO₃-N that was released from the sediments.

To determine the net accumulation rate of NO₃-N in the hypolimnion, the total mass data (Fig. 11) have been transformed into accumulation rate data by subtracting the spring profile data from the autumn profile data and dividing by the number of days between the spring and autumn samplings (Fig. 12).




Figure 12: Net NO₃-N accumulation rates (t d⁻¹) in the hypolimnion below 70 m. The regression line shows an increase in the net accumulation rate of 0.028 ± 0.03 t d⁻¹ ($P = 0.07$, $r^2 = 0.154$, $n = 22$). Open circle data were not included in the regression analysis (see text). Note that the Y-axis extends to -0.5 t d⁻¹ for the 2006 data point. Date ticks are 1 January in each year.

Mineralisation and release of nutrients from the sediments are driven by microbial processes that are a function of temperature and dissolved oxygen concentration. As the hypolimnion is well oxygenated and the temperature remains constant within ± 0.3 °C, the expectation would be for a reasonably consistent pattern of net accumulation rates allowing any trends to be well defined. A regression through all of these rate data (Fig. 12) shows a weakly significant trend in the data with the net accumulation rate increasing at 0.028 ± 0.03 t d⁻¹ each year ($P = 0.07$, $r^2 = 0.154$, $n = 22$) over the 34 year data record. The data for 1976 and 1987 were excluded from the regression analysis because they are for different periods than the rest of the data (see also Fig. 11). The data points are included in Figure 12 as an indication of what the net accumulation rates may have been in those two years.

The net accumulation rates of NO₃-N (Fig. 12) show a higher degree of variability between years than would be normally expected, with both the 1999 and 2006 data points falling well below the spread of the rest of the data and the trend line. The negative net accumulation rate in 2006 indicates a substantial loss of NO₃-N from the hypolimnion during the 2005-06 stratified period. Both of these data points are for years following a winter where there was incomplete mixing. This suggests that the

low values are anomalies relative to the rest of the data. The effect of incomplete mixing was discussed in a previous report (Gibbs 2007).

The net accumulation rate for the 2008/09 period was 1.93 t d^{-1} , which was 20% greater than in the previous year at 1.60 t d^{-1} , with the 2008/09 data point falling just above the trend line (Fig. 12).

3.7.3 Total N

Total nitrogen (TN) mass in Lake Taupo was estimated from the spring profile in each year. Regression analysis found that, although there was an average increase of about 12.9 t y^{-1} over the data record, there was no statistically significant trend in the long-term time-series data for TN (Fig. 13). There is obvious inter-annual variability in the mass of TN in the lake in spring around the long-term mean of 3400 t. This is despite a net annual external TN input to the lake of around 1200 t. The total mass of TN in Lake Taupo for spring 2008 was 3920 t.




Figure 13: Long-term time series estimates of the mass of total nitrogen (TN) in Lake Taupo in spring after winter mixing. Although there is an average increase of about 12.9 t y^{-1} over the data record, this apparent trend in the data is not statistically significant. The mean of the mass is around 3400 t. Date ticks are 1 January in each year.

4. Knowledge gaps

An earlier report (Gibbs 2006) commented on several knowledge gaps including in-lake processes in Lake Taupo, and process rates at the sediment-water interface. This report presents estimates of the net rate of NO₃-N accumulation in the hypolimnion during the stratified period. The low NO₃-N accumulation rate in the 2005/06 stratified period following incomplete mixing in winter 2005, and the immediate return of the net accumulation rate to the trend in the net rate data the following year, points to water column processes being as important as sediment processes for controlling NO₃-N concentrations and highlights the need to understand how the in-lake processes work.

The sum of the external inputs to the lake from the catchment via rivers minus the mass lost from the lake via the Waikato river is estimated to be around 1200 t y⁻¹ (W. Vant, Environment Waikato, pers. comm.). Despite this net input of TN, which represents around a third of the average mass of N in the lake (3400 t), there is no significant increase in the long term TN in the lake (Fig. 13). In addition, the total mass of NO₃-N in the hypolimnion just before winter mixing each year appears to reach a plateau (see shape of NO₃-N concentration curves Fig. 10). Together these data suggest that processes at the sediment-water interface and elsewhere in the hypolimnion are capable of sequestering very large amounts of N each year. However, as the net accumulation rate of NO₃-N in the hypolimnion is increasing, this suggests that the sediment processes of nitrogen burial, decomposition, mineralisation, nitrification, and denitrification are changing.

We have little or no information on any of these N transformation and sequestration process rates.

The appearance of free NH₄-N along with free DRP in the upper water column during late spring and summer in 2007, 2008 and 2009 is unusual as these nutrients would be expected to be rapidly taken up by phytoplankton growth. The source of that NH₄-N is unknown.

This report has also shown that both the mean annual water clarity and the mean annual chlorophyll *a* concentrations in the upper 10 m of water column have increased significantly since 1994. These parameters are usually inversely related and thus other factors must be influencing the observed increases. Note, the deep chlorophyll maximum is often up to 70% higher than the chlorophyll *a* concentrations in the upper 10 m of the lake water column. This represents a substantial amount of the algal biomass which is presently not being assessed.

5. Summary

- Using a linear regression through all data, the annual mean chlorophyll *a* concentration in the upper 10 m of water column in Lake Taupo, as an indicator of phytoplankton biomass, has increased at a rate of $0.025 \pm 0.017 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P = 0.005$, $r^2 = 0.033$, $n = 237$) over the 15 year monitoring period.
- It has become apparent that this increase in chlorophyll *a* data may not be a linear trend. The annual mean chlorophyll *a* data from 1994 to 2003 increased at a statistically significant rate of $0.087 \pm 0.029 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.001$, $r^2 = 0.857$, $n = 10$), but since 2000 there has been a statistically significant trend of decline at a rate of $0.033 \pm 0.031 \text{ mg m}^{-3} \text{ y}^{-1}$ ($P < 0.05$, $r^2 = 0.423$, $n = 10$). These data suggest an improvement in lake water quality.
- Peak chlorophyll *a* concentrations in winter have recently become highly variable with a value of 2.2 mg m^{-3} in 2007 compared with 3.0 mg m^{-3} in 2008, and just 1.9 mg m^{-3} in 2009.
- There is a substantial deep chlorophyll maxima (DCM) below the thermocline (40 m) in the lake during spring and summer with an estimated chlorophyll *a* concentration up to 70% higher than the chlorophyll *a* concentrations measured in the upper 10 m. The DCM was present through the 2008/09 spring and summer period.
- Algal species dominance followed a succession from the diatoms, *Asterionella formosa* and *Aulacoseira granulata*, in winter 2008 to *Fragilaria crotonensis* and the green algae, *Monoraphidium* sp. (formerly *Ankistrodesmus falcatus*), which became dominant through spring, then to the colonial green, *Botryococcus braunii*, and some large dinoflagellates, *Gymnodinium* sp., and *Peridinium* sp., which became dominant in summer and autumn 2009. Cyanobacteria (blue-green algae) were always present in low numbers in the upper water column throughout the 2008/09 monitoring period, with *Anabaena lemmermannii* being the most common species. There was an ever present background of small ($<5 \mu\text{m}$) unicellular flagellates throughout the year.
- Algae collected from the DCM in October 2008 were low in biomass and numbers and were similar in composition to the surface species. The low biomass compared with a DCM as indicated by the chlorophyll fluorescence

profile was probably due to the van Dorne sampler not collecting water from the centre of the DCM peak.

- There was a statistically significant trend of increase in the total mass of $\text{NO}_3\text{-N}$ in the hypolimnion before winter mixing of around 7.9 t yr^{-1} ($P < 0.001$, $r^2 = 0.41$, $n = 22$) which was the same as determined in the previous year but with a stronger correlation. The amount of $\text{NO}_3\text{-N}$ in the hypolimnion was around 90 t higher than the previous year.
- The net accumulation rate of $\text{NO}_3\text{-N}$ in the hypolimnion below 70 m in the last few years has been in the order of 2 t d^{-1} and regression analysis showed that there has been a weak trend of increase in that rate of 0.028 t d^{-1} each year ($P = 0.07$, $r^2 = 0.154$, $n = 22$) over the last 34 years. The net accumulation rate of $\text{NO}_3\text{-N}$ in 2008/09 was 1.93 t d^{-1} which was substantially higher than the 2007/08 rate of 1.60 t d^{-1} .
- There was a non-statistically significant increase in whole Lake TN of about 12.9 t y^{-1} . The TN content of the lake in spring 2008 was 3920 t, an increase of around 340 t since the previous year.
- The 2008/09 net VHOD rate for the period from September 2008 to March 2009 was $17.50 \pm 3.64 \text{ mg m}^{-3} \text{ d}^{-1}$ (mean \pm 95% confidence limit) which was almost $3 \text{ mg m}^{-3} \text{ d}^{-1}$ higher than the previous year at $14.51 \pm 2.94 \text{ mg m}^{-3} \text{ d}^{-1}$.
- There has been a statistically significant ($P < 0.002$, $r^2 = 0.69$, $n = 11$) increase in the VHOD rate of $1.04 \text{ mg m}^{-3} \text{ d}^{-1}$ each year since the low in 1999, suggesting a decline in lake water quality. While the period of the regression analysis is selected from lowest to highest, and thus does not reflect a long-term trend in Lake Taupo, this sustained increase in VHOD over an 11-year period implies a change in the export of organic carbon to the hypolimnion over this period, either from external inputs (i.e., land-use effects), or primary production within the lake, or a combination of both.
- Nutrient concentrations (DRP, $\text{NH}_4\text{-N}$, and $\text{NO}_3\text{-N}$) in the upper water column were generally comparable with concentrations since 2003 and are similar to historical concentrations before Mount Ruapehu erupted in 1995. However, since 2006/07 there have been elevated $\text{NH}_4\text{-N}$ but low $\text{NO}_3\text{-N}$ concentrations in the upper water column through summer and autumn. In winter 2009 there were elevated DRP and $\text{NO}_3\text{-N}$ concentrations in the upper water column.

- Bottom water temperatures gradually rose to 11.2 °C just before winter mixing at the beginning of August 2008, then fell rapidly to 10.7 °C a month later. Bottom water temperatures fell to 10.3 °C after mixing in August 2009.
- Water clarity during summer 2008/09 reached a maximum of 22 m, 3 m less than the highest recorded for Lake Taupo of 25 m on February 2008. Both of these extremely high clarity events may be attributed to extended periods of calm weather with very low surface run-off. The generally low freshwater inputs during these drought conditions would contribute less plant growth nutrients and sediment to the lake than in previous years.

In a previous annual report (Gibbs et al. 2002), three trends in the data were identified — increasing phytoplankton biomass in the upper 10 m, increasing NO₃-N mass in the lake hypolimnion prior to winter mixing, and an increasing range in the variability of water clarity — that were of concern with respect to the water quality of Lake Taupo. In the previous report, it was also shown that the net accumulation rate of NO₃-N in the hypolimnion during the stratified period has increased over the last 33 years, although the trend was not strong.

While these trends in the data are still present in the whole data set, there are indications that water quality may be beginning to improve in some areas e.g., mean annual chlorophyll *a* concentrations have been declining since between 2000 and 2003. In contrast, however, the VHOD rates have been increasing since 1999, an indication that the water quality is declining.

6. Acknowledgments

This report was made possible by the team effort of Philip King and Duncan Pearce of the Taupo Harbourmaster's Office, and Eddie Bowman (NIWA Rotorua) who have collected the data. Much of the success of this monitoring programme is attributable to the extra effort by Eddie and the team.

Water samples were processed in the NIWA chemistry laboratory and analytical results were provided by Graham Bryers, Margaret McMonagle, Cara Mackle and team. Quality control was provided by Mike Crump, Lab Manager.

Phytoplankton dominance and enumeration results were provided by Karl Safi.

7. References

- Burns, N.M. (1995). Using hypolimnetic dissolved oxygen depletion rates for monitoring lakes. *New Zealand Journal of Marine and Freshwater Research* 29: 1-11.
- Gibbs, M.M. (1995). Lake Taupo long term monitoring programme. NIWA consultancy report to Environment Waikato, report No. EVW60203/1.
- Gibbs, M.M. (2000). Lake Taupo long term monitoring programme: 1999-2000. NIWA consultancy report to Environment Waikato, report No. EVW01203.
- Gibbs, M.M.; Rutherford, J.C.; Hawes, I. (2002). Lake Taupo long term monitoring programme 2000 – 2001, with a review of accumulated data since 1994. NIWA consultancy report to Environment Waikato, report No. HAM2002-029, August.
- Gibbs, M.M. (2005). Lake Taupo long term monitoring programme 2003-2004: including two additional sites. NIWA consultancy report to Environment Waikato, report No.HAM2005-006, February.
- Gibbs, M.M. (2006). Lake Taupo long term monitoring programme 2004-2005: including two additional sites. NIWA consultancy report to Environment Waikato, report No.HAM2006-033, May.
- Gibbs, M.M. (2007). Lake Taupo long term monitoring programme 2005-2006. NIWA consultancy report to Environment Waikato, report No.HAM2007-029, March.
- Gibbs, M.M. (2009). Lake Taupo long term monitoring programme 2007-2008. NIWA consultancy report to Environment Waikato, report No.HAM2009-044, March
- Gibbs, M.M. (2010). Lake Taupo Near-Shore Water Quality Monitoring 2007-2009. Environment Waikato Technical Report 2010/02
- Helsel, D.R. & Hirsch, R.M. (1992). Statistical methods in water resources. Studies in Environmental Science no. 49, Elsevier Science Publishers, Amsterdam.

Utermöhl, von H. (1931). Neue Wege in der quantitativen Erfassung des Planktons. (Mit besondere Berücksichtigung des Ultraplanktons). Verh. Int. Verein. Theor. Angew. *Limnol.* 5: 567–595.

Vincent, W.F. (1983). Plankton production and winter mixing: contrasting effects in two oligotrophic lakes. *Journal of Ecology* 71: 1-20.

White, E.; Downes, M.; Gibbs, M.; Kemp, L.; Mackenzie, L. & Payne, G. (1980). Aspects of the physics, chemistry, and phytoplankton biology of Lake Taupo. *New Zealand Journal of Marine and Freshwater Research* 14(2): 139-148.

8. Appendices

8.1 Appendix 1: Site map, sampling strategy and methods

Lake monitoring sites were originally established using land-based markers (Fig.1). These have now been defined using GPS and corrected for curvature using WGS84 convention.

Site Map




Figure 1: Site map of Lake Taupo showing location of the routine monitoring site at mid lake (A). Two additional sites at Kuratau Basin (B) and the Western Bays (C) were sampled between January 2002 and December 2004 inclusive. Data from those sites have been retained with the Site A data presented in the appendices. Map coordinates are in NZ Map Grid with WGS84 correction. Lat. Long WGS 84 corrected co-ordinates of "Site A" are $38^{\circ} 46'.810\text{ S}$; $175^{\circ} 58'.440\text{ E}$.

The following section has been copied from Gibbs 1995, and modified after 1998.

8.1.1 Methods

The sampling site was selected in the central basin of Lake Taupo (Site Map) with a water depth of about 160 m. This site is more than 5 km from the nearest land and is exposed to both the north-south and east-west axis of the lake.

To calculate VHOD requires two measurements each year far enough apart in time for a measurable change to occur in the DO concentrations in the hypolimnion of the lake. Details of the procedure and limitations of this measurement are described by Vant (1987). For the monitoring of Lake Taupo, which mixes briefly in winter between July and August, the initial sampling time was selected to be in October, to give sufficient time for thermal stratification to establish a stable hypolimnion. The final sampling time was selected to be in April, before lake cooling causes the downward movement of the thermocline which precedes the winter mixing.

At each of these biannual samplings, a detailed profile of DO and temperature was measured. Prior to 1998, measurements were made at 1 m depth intervals through the full depth of the water column using an in situ recording Applied Microsystems STD-12 profiler fitted with a Royce DO sensor, and compared with manual measurements of DO and temperature made at 10 m depth intervals from the surface to the bottom of the lake using a Yellow Springs Instrument (YSI) model 58 dissolved oxygen meter fitted with a stirred Model 5739 probe on a 160 m cable. Subsequent to 1998, a Richard Brancker Research (RBR) model TD410 conductivity-temperature-depth (CTD) profiler fitted with a stirred YSI model 5739 DO sensor was used. In January 2002, the TD410 CTD profiler was upgraded to an RBR model XR420f freshwater CTD profiler fitted with the YSI model 5739 DO sensor and a Seapoint chlorophyll fluorescence probe. The DO sensor was calibrated regularly by NIWA, Rotorua staff and chlorophyll fluorescence was converted to chlorophyll *a* from extracted chlorophyll *a* analyses of water samples collected beside profiler.

In January 2008, the XR420f profiler was upgraded to a RBR model XR620f freshwater profiler/logger with improved sensitivity. The new profiler is fitted with a Sea Point chlorophyll fluorescence probe and a Li-Cor underwater photosynthetically active radiance (PAR) sensor to measure in situ light levels and light extinction (K_d) associated with the vertical distribution of algal biomass within the lake water column. In the new system the YSI dissolved oxygen (DO) sensor was replaced with an Oxyguard DO sensor fitted to a separate RBR logger attached to the profiling frame.

Cross-calibration between the two profilers confirmed the quality of the data and the XR420f has been retained as a back-up.

The following parameters were also measured as profiles from water samples collected using a van Dorn water sampling bottle starting at 1 m and then at 10 m intervals from 10 m to the bottom of the lake:

DO, chlorophyll *a*, dissolved reactive phosphorus (DRP), dissolved organic phosphorus (DOP), particulate phosphorus (PP), total phosphorus (TP), nitrate + nitrite nitrogen ($\text{NO}_3\text{-N}$), ammoniacal nitrogen ($\text{NH}_4\text{-N}$), dissolved organic nitrogen (DON), particulate nitrogen (PN), total nitrogen (TN), urea nitrogen (Urea-N), total suspended solids (SS), volatile suspended solids (VSS), particulate carbon (PC), dissolved organic carbon (DOC), and water colour. (* Little, if any nitrite is ever found in the Lake Taupo water column, hence the use of $\text{NO}_3\text{-N}$).

Note: TN and TP values are the summation of all other N and P components, respectively, excluding Urea-N which is part of the DON component.

Additional parameters measured but not as complete profiles were:

Water clarity (by Secchi disc depth) and algal species composition and abundance on water samples from 1, 10, 50, 100, and 140 m.

Determinations on the water samples were made with the standard methods routinely used for freshwater analysis by NIWA.

Algal species composition and abundance were obtained by settling a measured volume of sample (up to 100 mL) in Utermöhl (1931) tubes and counting on an inverted microscope. Biovolume was estimated from cell volume tables calculated from the cell dimensions of each species. Dominance was estimated from relative biovolumes with the highest biovolume assigned dominance 1 as most common and the lowest biovolume assigned the dominance 10 as rare. Professional judgement was used to relate dominance between samplings.

Since 2007, dominance is no longer used and the algal data are reported in cell counts and biovolume.

Data for the long term monitoring programme were scheduled to be collected from the mid-lake sampling station at 2 weekly intervals. The practicality of achieving this

target was limited by the weather and in reality data were generally collected at about 2-3 weekly intervals. Parameters measured were:

DO and temperature profiles at 1 m depth intervals to the bottom of the lake by RBR XR420f profiler, water clarity as Secchi disc depth, and a 10 m tube water sample was collected for measurement of chlorophyll a, NO₃-N, NH₄-N, TN, DRP, TP, and algal species dominance.

Near-bottom water samples from 150 m were collected using a van Dorn water sampling bottle and analysed for DRP, NO₃-N, and NH₄-N.

8.1.2 Data handling and less than detection limit values

All data in this report have been processed and manipulated on Excel spreadsheets. Data are rounded using the Excel protocol to an appropriate number of significant numbers based on the need for detailed knowledge tempered with the confidence in the precision and accuracy of the analytical methods used. This treatment may lead to small differences between electronic copies of the data and the values presented in this report.

The difference between the written report and the Excel spreadsheet of essentially the same data is the treatment of the less than detection limit (<DL) results. The data have in the past been written as <DL or <DL(value). For statistical analysis the excel spreadsheet replaces <DL with 0 or uses the value in brackets in place of 0. Although it is recognised that the former action will be in error, the use of the value in brackets requires some justification.

In discussion with Burns Macaskill, Graham McBride, and Mike Crump from NIWA on this issue, the following conclusions were reached:

- In general the data is reported as a series of results from analytical methods which have known limitations and precision. The raw number is reported where ever possible so that the user can draw their own conclusions about the reliability of the "last significant figure" on any result when performing data manipulations.
- The real problem arises at very low levels and the result obtained is less than the method's prescribed DL. The problem is not so much the result obtained but what to do with it which in turn raises the question 'What do we mean by detection limit'?

- In the book "Statistical methods in water resources" Helsel & Hirsch 1992 [Studies in Environmental Science 49, Elsevier], and chapter 13 "Methods for data below the reporting limit" it is pointed out that the 'detection limit' is variously known as the 'reporting limit' or the 'limit of quantitation'. If no other value is available, there are 3 main options: call it zero (which is clearly an under estimate), call it the detection limit (which is clearly an over estimate), or call it half the detection limit (which gives a 50:50 chance of an over or under estimate). The choice then is one of 'which convention do you wish to use'. In the written reports, I have treated the <DL as zero for summation purposes. This is an under estimate which I should have noted on each report page so that anyone using that data is aware of the convention used.
- An alternative approach is to say that, before the sample is analysed, the DL is the **predicted minimum** level that will be found using the stipulated method. However, once the sample is analysed the result is what was **actually measured** and may be <DL on the day of analysis. As it is an actual analytical result, that value (reported in brackets) should be reported even though it is <DL. This implies that the method DL is in reality a reporting level or level of confidence.
- The "DL" was derived for the Lake Taupo data, on each analytical occasion, from a series of blanks and 1ppb standards run with the samples. The "DL" is set as 3 times the SD of the 1 ppb standard. This is actually a limit of confidence. All samples are run in duplicate and the mean of the two results becomes the concentration reported.
- With the introduction of the Lachet FIA system, the limits of detection have been confidently lowered to the point where replicate results may often be <DL. In these instances, in the written report, the value is reported as <DL(result). In the past I have still used the <DL = 0 convention in summation for the TN and TP data. This is obviously wrong and the actual result should be used, as is done in the electronic spreadsheet.

In this report the analytical value 'on-the-day' has been used wherever possible. Data reported as <DL use the <DL = DL/2 convention. Past data have not been corrected or altered to conform to this protocol.

These technical details are incorporated in this annual report so that data users are aware of how the 'DL' or confidence limit was set and how the values <DL are treated when performing data manipulations.

There is still the question of how to deal with numbers where the result has been simply reported as <DL. The use of the DL/2 convention is probably closer to reality than the DL = 0 convention.

Helsel & Hirsch suggest an alternative method for estimating a value in the <DL range. If there is sufficient real data >DL, a probability curve can be derived and extrapolated around the DL to generate the most probable number for the <DL value.

8.1.3 Statistical methods

Copied from Gibbs (2000).

In this report we have used linear regressions and associated statistical tests to examine trends. The key result of these procedures is the coefficient of determination (r^2), which measures the amount of variability in the data that is accounted for by the regression. Another is the P-value². This can be used as a weight of evidence against the hypothesis that there was in fact no trend. This weight is strong when P is small, meaning that a trend at least as large as that measured could have occurred merely by chance—we have only a limited number of data from which to infer the strength of any trend, so our measurements always are uncertain to some degree. So if P is low enough (taken as less than 5% in this report, which is the usual practice), it is conventional to say that the measured trend is "statistically significant", and that convention is followed in this report. However, it is important (and often not realised) to note that the P-value cannot be used as an absolute weight of evidence. This is because it tends to decrease as the number of samples taken in a given period is increased. For example, when we plot monthly Secchi disc depth data from 1994–2001 (Figure 3A, Gibbs 2000) with these 93 data we obtain a statistically significant result (because $P < 0.05$)—even though the coefficient of determination was only $r^2 = 0.0445$. When we plot the minimum winter clarity over this period we then have only 7 data. In this case (Fig. 3B, Gibbs 2000) we happen to have the same measured trend slope with a much higher coefficient of determination ($r^2 = 0.464$), yet the result is not statistically significant (because $P = 0.09$). This is entirely because of the reduced number of samples in the winter minimum case. What this makes clear is that the P value is useful as a relative weight of evidence when comparing datasets of the same

² It is defined as the probability of obtaining a trend at least as extreme as was obtained if in fact there was no trend at all.

size, but it has no evidential meaning when comparing results from datasets of very different sizes.

8.2 Appendix 2: The calculation of net VHOD rates

Copied from Gibbs 1995.

Rationale

In the strictest terms, VHOD can only be calculated for a lake which has thermally stratified and the resultant thermocline provides an effective barrier against re-oxygenation of the hypolimnion. The measure of the barrier efficiency is the rate of heating of the hypolimnion following stratification as heat will be transferred across the thermocline at a similar rate to oxygen.

In Lake Taupo, the thermal inertia of the hypolimnion is so great that heating during the stratified period is typically about 0.2 °C and never more than 0.4 °C over a 200 day period. While this would seem to meet the temperature criterion, in a lake that large, oxygen can be transferred into the hypolimnion by mechanisms other than diffusion.

Wind induced mixing may increase turbulent diffusion across the thermocline as would an internal seiche on the thermocline. Both of these mechanisms would transfer heat. The penetration of the thermocline by an under-flowing density current would entrain oxygenated surface water into the hypolimnion with that flow. As the density current must be colder than the thermocline to plunge through it, there is no heat transferred with this mechanism.

In Lake Taupo the Tongariro River water is always colder than the lake surface water and for at least 9 months of the year it is also colder than the minimum lake water temperature of 10.3 °C. Thus, during most of the stratified period, the Tongariro River flows directly into the hypolimnion entraining oxygenated surface water with it. The amount of surface water entrained has been estimated to be about 10 times the river discharge. The amount of oxygen transported in this way is likely to be more than 200 tonnes per day.

Clearly this is a substantial oxygen input which invalidates the concept of the thermocline forming an oxygen barrier for purposes of calculating the VHOD. The true VHOD may only be calculated during mid summer when the Tongariro River flows deep into the epilimnion but does not penetrate the thermocline.

The data collected to date indicates that hypolimnetic oxygen depletion occurs throughout the stratified period - with or without the density current re-oxygenation - and hence the value obtained from a VHOD calculation over the whole stratified period is the net VHOD rate taking all the factors affecting the hypolimnion into account.

As the data from 1996/97 shows, the density current also advects dissolved organic nutrients with it. Hence, management strategies which affect the Tongariro River also impact on the lake. Hence it is appropriate to use the net VHOD rate for inter-annual comparisons rather than the true VHOD rate calculated only through mid summer.

Method of calculation

The following is the method used to calculate the net VHOD rate for Lake Taupo.

Requirements: Microsoft Excel spreadsheet or equivalent.

Although the thermocline in Lake Taupo is usually at about 40 m, the isothermal water column lies below 70 m. To accommodate the gradient across the thermocline, the net VHOD rate calculation only uses oxygen data from below 70 m.

To calculate the mean oxygen concentration in the water column below 70 m, the DO concentration at each 10 m depth increment is multiplied by the volume of the 10 m slice it came from. This assumes rapid horizontal mixing and minimal vertical mixing to extrapolate one DO value across the whole lake. Historical data from multiple sites would suggest that this is a reasonable assumption.

The slice volumes (hypsographic volumes) for Lake Taupo have been calculated for 10 m thick layers centred on the 5 m point of each slice i.e., 75, 85, 95, 105 m etc. The DO measurements are made at 10 m intervals i.e., 70, 80, 90, 100, 110 m etc.

The mass of oxygen in each 10 m slice is the average of the DO concentration at the top and bottom of a slice multiplied by the slice volume. i.e., for the 70 - 80 m slice the calculation is:-

$$\text{Mass}_{70-80m} = ((\text{DO}_{70m} + \text{DO}_{80m}) \div 2) \times \text{Volume}_{70-80m}$$

For each profile date:

Compute the mass for each 10 m slice between 70 m and 150 m and sum the results as the total mass of DO in the hypolimnion below 70 m. Sum the slice volumes below 70 m as the total volume of the hypolimnion below 70 m.

The volume weighted mean DO concentration is the total mass value divided by the total volume value.

Use the sequential day number or equivalent to construct a time series of volume weighted mean DO concentrations over the stratified period and use the Excel regression analysis tool to obtain the $y = ax + b$ straight line fit for these data.

As the DO data are in g m^{-3} , the value of 'a' is in $\text{g m}^{-3} \text{ d}^{-1}$. Multiply 'a' by 1000 to get the net VHOD rate in $\text{mg m}^{-3} \text{ d}^{-1}$. The negative sign indicates a loss rate.

The hypsographic volumes and upper surface areas of the 10 m slices through the whole depth of Lake Taupo are listed at the end of this section.

Lake Taupo Hypsographic Data used in the Net VHOD RATE calculation.

Slice depths (m)	Volume of slice (km^3)	Upper surface area of slice (km^2)
0 - 10	5.849359	600
10 - 20	5.599702	570
20 - 30	5.459951	550
30 - 40	5.359888	542
40 - 50	5.288266	530
50 - 60	5.150538	528
60 - 70	4.899510	502
70 - 80	4.619076	478
80 - 90	4.278738	446
90 - 100	3.847292	410
100 - 110	3.006616	360
110 - 120	1.730549	245
120 - 130	0.837468	110
130 - 140	0.394439	60
140 - 150	0.073333	22
150 -	0	0

Statistical evaluation of the VHOD rate

From the 1999-2000 monitoring report (Gibbs 2000), the VHOD rate is expressed as the calculated net VHOD rate \pm the 95% confidence limit. This gives a meaningful estimate of the range within which the VHOD rate lies and is more appropriate than the standard deviation on the data or a standard error estimate on the regression coefficient.

Julian Date or sequential day number for each day of the year excluding leap years. For Leap Years, add 1 to the sequential day number from 1 March to 31 December of that year.

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	32	60	91	121	152	182	213	244	274	305	335
2	2	33	61	92	122	153	183	214	245	275	306	336
3	3	34	62	93	123	154	184	215	246	276	307	337
4	4	35	63	94	124	155	185	216	247	277	308	338
5	5	36	64	95	125	156	186	217	248	278	309	339
6	6	37	65	96	126	157	187	218	249	279	310	340
7	7	38	66	97	127	158	188	219	250	280	311	341
8	8	39	67	98	128	159	189	220	251	281	312	342
9	9	40	68	99	129	160	190	221	252	282	313	343
10	10	41	69	100	130	161	191	222	253	283	314	344
11	11	42	70	101	131	162	192	223	254	284	315	345
12	12	43	71	102	132	163	193	224	255	285	316	346
13	13	44	72	103	133	164	194	225	256	286	317	347
14	14	45	73	104	134	165	195	226	257	287	318	348
15	15	46	74	105	135	166	196	227	258	288	319	349
16	16	47	75	106	136	167	197	228	259	289	320	350
17	17	48	76	107	137	168	198	229	260	290	321	351
18	18	49	77	108	138	169	199	230	261	291	322	352
19	19	50	78	109	139	170	200	231	262	292	323	353
20	20	51	79	110	140	171	201	232	263	293	324	354
21	21	52	80	111	141	172	202	233	264	294	325	355
22	22	53	81	112	142	173	203	234	265	295	326	356
23	23	54	82	113	143	174	204	235	266	296	327	357
24	24	55	83	114	144	175	205	236	267	297	328	358
25	25	56	84	115	145	176	206	237	268	298	329	359
26	26	57	85	116	146	177	207	238	269	299	330	360
27	27	58	86	117	147	178	208	239	270	300	331	361
28	28	59	87	118	148	179	209	240	271	301	332	362
29	29		88	119	149	180	210	241	272	302	333	363
30	30		89	120	150	181	211	242	273	303	334	364
31	31		90		151		212	243		304		365
												31

8.3 Appendix 3: Temperature and dissolved Oxygen data

Includes accumulated data since 1994.

* represents data missing or invalid.

For completeness, additional data from the Kuratau Basin (site B) and Western Bays (site C) collected for the period between January 2002 and December 2004 are included as separate sheets following the mid-lake data from site A for those years.

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 11 September 2007
2008-2009
Temperature

Date	4/09/2008	16/09/2008	14/10/2008	4/11/2008	26/11/2008	22/12/2008	13/01/2009	22/01/2009	28/01/2009	11/02/2009	25/02/2009	16/03/2009	26/03/2009	15/04/2009	7/05/2009	27/05/2009	18/06/2009	6/07/2009	13/08/2009
Depth (m)																			
0	10.97	11.34	12.59	13.37	15.45	18.84	19.67	19.84	20.88	21.42	20.46	18.71	17.96	16.60	15.05	12.97	11.60	10.93	10.43
10	10.92	11.14	12.09	12.94	15.26	17.50	19.55	19.23	20.17	21.21	20.39	18.29	17.95	16.59	15.04	12.96	11.61	10.93	10.41
20	10.85	10.99	11.93	12.62	15.17	15.77	16.97	19.12	18.45	20.04	20.37	18.25	17.94	16.59	15.04	12.96	11.61	10.92	10.41
30	10.82	10.93	11.85	12.55	12.87	13.32	13.60	13.90	13.21	13.92	14.47	16.68	13.86	16.58	15.04	12.90	11.61	10.92	10.41
40	10.79	10.91	11.75	12.35	12.07	12.27	12.19	12.11	11.90	12.09	12.84	12.43	12.13	12.53	12.55	12.62	11.60	10.91	10.38
50	10.75	10.88	11.59	11.51	11.44	11.39	11.33	11.52	11.31	11.50	11.62	11.56	11.45	11.56	11.64	11.50	11.60	10.92	10.36
60	10.72	10.79	10.90	10.83	10.93	11.06	11.08	11.04	11.05	11.19	11.18	11.22	11.19	11.12	11.17	11.06	11.60	10.92	10.36
70	10.69	10.69	10.76	10.79	10.78	10.88	10.89	10.90	10.89	10.97	10.92	10.98	10.98	10.92	11.01	10.94	11.60	10.92	10.36
80	10.66	10.68	10.71	10.72	10.76	10.81	10.82	10.87	10.84	10.86	10.87	10.88	10.89	10.92	10.93	10.90	11.59	10.91	10.35
90	10.66	10.66	10.69	10.70	10.77	10.78	10.78	10.81	10.80	10.81	10.82	10.83	10.84	10.88	10.89	10.88	11.41	10.92	10.34
100	10.65	10.65	10.68	10.68	10.82	10.75	10.76	10.80	10.78	10.77	10.79	10.81	10.81	10.86	10.86	10.86	11.09	10.92	10.34
110	10.64	10.64	10.66	10.67	10.78	10.73	10.75	10.78	10.74	10.76	10.77	10.80	10.79	10.84	10.86	10.85	11.00	10.91	10.33
120	10.63	10.64	10.64	10.65	10.78	10.71	10.73	10.77	10.74	10.75	10.76	10.79	10.78	10.82	10.84	10.84	10.98	10.91	10.33
130	10.63	10.63	10.60	10.63	10.79	10.70	10.72	10.74	10.73	10.73	10.75	10.77	10.77	10.79	10.82	10.82	10.95	10.91	10.33
140	10.63	10.62	10.59	10.63	10.81	10.70	10.72	10.73	10.73	10.74	10.74	10.77	10.76	10.78	10.80	10.81	10.94	10.90	10.33
150	10.62	10.62	10.59	10.63	10.80	10.70	10.71	10.74	10.72	10.73	10.74	10.76	10.76	10.78	10.80	10.81	10.89	10.90	10.30

Dissolved Oxygen (g m⁻³)

Depth (m)																			
0	10.03	9.84	10.29	*	10.09	9.29	8.67	9.24	8.52	8.48	*	9.26	9.44	9.33	10.05	10.13	10.47	8.91	9.83
10	10.85	10.65	10.29	*	10.08	9.72	9.21	8.89	8.45	8.34	*	9.16	10.06	10.11	10.15	10.25	10.73	9.88	10.72
20	10.90	11.05	10.50	*	10.00	9.39	8.88	8.68	8.47	8.19	*	9.40	10.55	10.76	10.15	10.13	10.59	11.06	11.48
30	11.12	10.91	10.46	*	9.79	9.81	9.02	8.53	8.54	8.20	*	9.12	10.34	10.83	10.15	10.17	10.57	11.31	11.57
40	10.76	10.82	10.34	*	9.23	9.69	8.96	8.46	8.06	8.36	*	8.24	9.86	10.39	9.15	9.78	10.56	11.28	11.39
50	10.88	10.63	10.05	*	9.10	9.05	8.49	8.06	7.98	7.92	*	7.97	9.25	9.58	8.91	9.47	10.49	11.29	11.39
60	10.74	10.55	9.89	*	8.54	8.77	8.25	7.91	7.81	7.80	*	7.62	8.97	9.06	8.67	8.73	10.40	11.03	11.20
70	10.52	10.25	9.86	*	8.60	8.53	8.10	7.64	7.74	7.71	*	7.55	8.94	8.84	8.51	8.60	10.43	11.05	11.16
80	10.48	10.20	9.81	*	8.43	8.47	7.98	7.46	7.66	7.64	*	7.44	8.54	8.21	7.79	8.25	10.43	10.83	10.86
90	10.34	10.13	9.85	*	8.44	8.21	7.92	7.38	7.56	7.60	*	7.37	8.45	8.24	7.79	8.24	10.25	10.87	10.97
100	10.28	10.10	10.03	*	8.20	8.22	7.78	7.25	7.53	7.44	*	7.26	8.24	8.07	7.65	8.10	8.65	10.68	10.87
110	9.79	10.00	10.13	*	8.31	7.99	7.67	7.22	7.47	7.31	*	7.20	8.26	8.12	7.62	8.06	8.53	10.72	10.90
120	9.62	9.97	10.09	*	8.04	7.91	7.63	7.17	7.32	7.26	*	7.01	7.94	8.02	7.63	7.79	8.17	10.55	10.86
130	9.42	9.75	9.83	*	8.09	7.70	7.48	7.21	7.24	7.04	*	7.03	7.93	8.15	7.59	7.83	8.11	10.55	10.71
140	9.37	9.52	9.76	*	7.88	7.59	7.40	7.24	7.08	6.92	*	6.68	7.08	8.01	7.74	7.49	7.99	10.48	10.80
150	9.17	9.24	9.85	*	7.85	7.48	7.25	7.03	6.90	6.72	*	6.59	6.91	7.55	7.35	7.30	7.97	10.30	10.77

Secchi depth

(m)	13.0	14.5	12.2	12.0	10.0	12.0	13.0	14.8	18.0	22.0	20.0	15.6	18.5	18.0	16.0	15.0	16.0	15.0	12.0
-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 11 September 2007
2007-2008
Temperature

Date	11/9/2007	9/10/2007	30/10/2007	15/11/2007	4/12/2007	20/12/2007	17/01/2008	31/01/2008	14/02/2008	27/02/2008	13/03/2008	26/03/2008	17/04/2008	7/05/2008	22/05/2008	5/06/2008	19/06/2008	1/07/2008	15/07/2008	7/08/2008	20/08/2008
Depth (m)																					
0	11.00	12.33	12.84	13.47	16.64	17.38	21.23	19.79	19.87	19.28	18.83	19.26	17.88	15.67	14.65	13.60	12.89	11.97	11.42	11.06	10.70
10	10.99	11.69	11.83	13.19	16.20	17.15	19.96	19.62	19.81	19.26	18.75	19.24	17.87	15.67	14.65	13.60	12.90	12.03	11.41	10.98	10.70
20	10.98	11.67	11.76	12.92	14.48	14.76	17.21	17.59	19.65	19.24	18.75	18.92	17.85	15.67	14.65	13.59	12.90	12.03	11.40	10.98	10.69
30	10.99	11.44	11.70	12.86	12.58	13.19	13.64	13.82	16.07	14.08	16.20	16.92	15.58	15.67	14.65	13.60	12.90	12.01	11.40	10.98	10.69
40	10.99	11.42	11.64	12.78	12.02	12.18	12.26	12.31	12.63	12.24	12.54	12.44	12.38	15.27	12.27	13.60	12.90	12.03	11.40	10.98	10.69
50	10.99	11.39	11.51	11.80	11.69	11.75	11.64	11.61	11.80	11.71	11.76	11.77	11.72	12.11	11.66	11.93	12.86	12.03	11.39	10.99	10.70
60	10.99	11.34	11.43	11.49	11.42	11.53	11.41	11.39	11.47	11.44	11.47	11.48	11.48	11.56	11.44	11.54	11.60	12.03	11.39	10.98	10.70
70	10.99	11.16	11.32	11.37	11.29	11.33	11.23	11.26	11.33	11.30	11.34	11.29	11.34	11.37	11.32	11.37	11.36	11.61	11.38	10.98	10.70
80	10.96	11.00	11.23	11.31	11.25	11.23	11.22	11.17	11.25	11.24	11.23	11.27	11.29	11.27	11.29	11.27	11.39	11.38	10.98	10.70	10.70
90	10.96	10.98	11.16	11.17	11.14	11.12	11.12	11.11	11.19	11.18	11.18	11.17	11.20	11.21	11.22	11.24	11.23	11.35	10.98	10.70	10.70
100	10.96	10.98	11.07	11.10	11.10	11.09	11.12	11.09	11.15	11.14	11.14	11.14	11.17	11.16	11.18	11.21	11.21	11.28	11.30	10.98	10.70
110	10.96	10.97	11.04	11.04	11.07	11.04	11.06	11.08	11.11	11.11	11.11	11.12	11.14	11.16	11.16	11.19	11.19	11.28	11.25	10.98	10.70
120	10.96	10.96	11.02	11.02	11.05	11.03	11.04	11.06	11.07	11.09	11.09	11.11	11.15	11.15	11.16	11.17	11.25	11.22	10.98	10.70	10.70
130	10.96	10.96	11.00	11.00	11.02	11.00	11.02	11.05	11.06	11.07	11.07	11.09	11.12	11.13	11.15	11.15	11.22	11.20	10.98	10.70	10.70
140	10.96	10.96	10.98	10.97	10.99	11.01	11.00	11.05	11.05	11.06	11.06	11.08	11.11	11.12	11.13	11.15	11.17	11.19	10.98	10.70	10.70
150	10.96	10.95	10.96	10.95	10.98	10.99	11.00	11.04	11.04	11.05	11.06	11.08	11.11	11.12	11.13	11.15	11.16	11.19	10.98	10.70	10.70

Dissolved Oxygen (g m⁻³)

Depth (m)	0	11.00	10.23	10.18	10.03	9.35	9.21	8.61	*	10.77	9.20	9.38	9.87	9.49	9.91	10.13	10.36	10.53	10.75	10.89	10.21	9.55
10	11.12	10.37	10.27	10.11	9.45	9.24	8.63	*	8.76	9.09	9.05	8.61	8.97	9.04	9.37	9.84	10.26	10.63	10.66	11.03	10.80	
20	10.87	10.12	10.25	10.07	9.23	9.21	8.70	*	9.00	9.32	9.24	8.85	8.46	8.97	9.18	9.72	10.14	10.32	10.51	11.04	11.16	
30	10.99	10.17	10.07	10.17	9.36	9.37	8.93	*	9.35	9.45	9.01	8.73	8.52	8.86	9.16	9.63	10.10	10.37	10.48	10.94	11.11	
40	10.84	9.92	10.02	9.97	9.09	9.09	8.69	*	9.01	8.92	8.96	8.57	8.72	8.87	8.68	9.81	10.12	10.40	10.42	10.72	11.08	
50	10.92	10.09	9.85	9.66	9.08	9.21	8.67	*	8.64	8.82	8.60	8.51	8.48	8.45	8.56	8.22	10.10	10.31	10.52	10.83	11.07	
60	11.07	9.96	9.52	9.75	9.14	8.69	8.60	8.70	8.44	8.49	8.34	8.15	8.20	8.25	8.58	8.96	9.51	10.36	10.45	10.60	11.05	
70	10.89	9.90	9.77	9.30	8.74	8.69	8.26	8.22	8.19	8.15	8.02	7.79	7.84	7.89	8.37	8.65	9.07	10.28	10.39	10.76	10.98	
80	10.90	9.59	9.58	9.12	8.76	8.38	8.03	8.05	8.16	7.88	7.92	7.52	7.71	7.90	8.30	8.53	8.91	9.60	10.34	10.74	10.96	
90	10.66	9.63	9.42	9.07	8.62	8.46	8.10	8.06	7.99	7.87	7.76	7.47	7.57	7.68	8.22	8.45	8.72	9.18	10.23	10.73	10.91	
100	10.64	9.58	9.49	9.14	8.46	8.41	7.90	7.90	7.97	7.86	7.69	7.45	7.45	7.46	8.14	8.44	8.66	9.06	9.93	10.72	10.90	
110	10.62	9.57	9.16	8.83	8.37	8.46	7.83	7.87	7.81	7.64	7.50	7.20	7.29	7.38	8.03	8.19	8.43	8.72	9.34	10.68	10.84	
120	10.66	9.52	9.27	8.95	8.42	8.08	7.95	7.52	7.82	7.39	7.45	7.20	7.29	7.38	7.94	8.16	8.32	8.55	8.94	10.67	10.83	
130	10.42	9.35	9.01	8.81	8.31	8.13	7.72	7.40	7.59	7.41	7.27	7.16	7.18	7.19	7.86	8.14	8.31	8.79	10.63	10.57		
140	10.40	9.30	9.11	8.81	8.28	7.88	7.74	7.27	7.62	7.05	7.10	7.10	7.13	7.17	7.81	7.61	8.01	8.25	8.48	10.62	10.38	
150	10.37	9.13	8.91	8.45	7.95	7.95	7.33	7.35	7.27	7.00	6.76	6.59	6.72	6.85	7.40	7.50	7.73	8.08	8.48	10.57	9.67	

Secchi depth

(m)	11	15	16	14	15	17.5	22.5	21.5	25	22	22	19	20.5	16	17	15	16.5	14	13	12.5	12.5
-----	----	----	----	----	----	------	------	------	----	----	----	----	------	----	----	----	------	----	----	------	------

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 4 September 2006
2006-2007
Temperature

Date	4/09/2006	26/09/2006	18/10/2006	1/11/2006	5/12/2006	19/12/2006	9/01/2007	25/01/2007	8/02/2007	21/02/2007	21/03/2007	3/04/2007	19/04/2007	8/05/2007	22/05/2007	14/06/2007	27/06/2007	18/07/2007	8/08/2007	23/08/2007	11/09/2007
Depth (m)																					
0	11.10	11.88	11.72	12.43	15.21	15.62	16.51	18.60	19.31	19.58	18.70	18.04	16.49	19.29	15.17	13.56	12.38	11.43	11.15	11.00	11.00
10	10.93	11.48	11.73	12.27	14.06	15.46	16.41	18.42	18.98	19.12	18.03	18.03	16.48	18.98	15.16	13.56	12.39	11.43	11.15	11.00	10.99
20	10.93	11.29	11.72	12.25	13.87	14.45	15.44	17.96	18.16	17.62	17.99	17.94	16.47	18.16	15.16	13.56	12.39	11.43	11.16	11.00	10.98
30	10.89	11.19	11.69	12.20	13.69	14.15	14.42	15.82	14.86	15.17	15.18	16.72	16.47	14.86	15.16	13.56	12.39	11.36	11.15	11.00	10.99
40	10.87	11.15	11.45	12.10	13.16	12.43	12.25	13.05	12.89	13.09	12.65	13.50	13.78	12.89	15.15	13.56	12.39	11.29	11.16	11.00	10.99
50	10.83	11.08	11.34	11.96	11.77	11.64	11.74	11.84	11.89	11.91	11.94	12.33	12.47	11.89	11.99	13.55	12.39	11.27	11.16	11.00	10.99
60	10.82	11.06	11.25	11.34	11.20	11.36	11.29	11.47	11.39	11.46	11.51	11.65	11.69	11.39	11.54	11.77	12.38	11.25	11.15	11.00	10.99
70	10.82	11.00	11.21	11.17	11.11	11.21	11.15	11.26	11.21	11.22	11.28	11.33	11.21	11.33	11.35	11.39	11.22	11.16	11.01	10.99	
80	10.82	10.94	11.16	11.06	11.06	11.10	11.09	11.14	11.15	11.16	11.20	11.20	11.15	11.21	11.22	11.28	11.17	11.16	11.01	10.96	
90	10.81	10.90	11.08	10.99	10.97	11.03	11.03	11.04	11.06	11.05	11.09	11.11	11.13	11.06	11.12	11.11	11.22	11.14	11.16	11.01	10.96
100	10.81	10.87	10.97	10.94	10.94	11.00	11.00	11.00	11.03	11.05	11.05	11.10	11.09	11.03	11.10	11.10	11.16	11.13	11.16	11.01	10.96
110	10.81	10.84	10.89	10.91	10.91	10.96	10.98	10.98	11.01	11.02	11.03	11.04	11.05	11.01	11.07	11.09	11.12	11.12	11.16	11.01	10.96
120	10.80	10.81	10.86	10.88	10.90	10.94	10.97	10.99	11.06	11.02	11.04	11.04	11.06	11.06	11.07	11.08	11.11	11.12	11.16	11.01	10.96
130	10.79	10.79	10.85	10.85	10.88	10.92	10.95	10.97	10.99	10.99	11.01	11.01	11.03	10.99	11.03	11.07	11.08	11.11	11.16	11.01	10.96
140	10.76	10.78	10.83	10.84	10.88	10.89	10.94	10.97	10.97	10.98	10.99	11.00	11.02	10.97	11.03	11.05	11.07	11.10	11.16	11.01	10.96
150	10.75	10.76	10.82	10.85	10.88	10.91	10.93	10.99	10.96	11.02	11.04	11.03	11.02	11.00	11.04	11.05	11.07	11.10	11.16	11.01	10.96

Dissolved Oxygen (g m⁻³)

Depth (m)																					
0	10.52	10.31	10.36	10.23	9.62	9.52	9.35	8.99	8.95	9.16	9.31	9.44	9.74	9.20	10.01	10.01	10.26	10.36	10.96	11.02	11.00
10	10.47	10.28	10.31	10.16	9.69	9.52	9.52	8.95	8.96	9.26	9.27	9.51	9.73	9.29	10.06	9.95	10.37	10.43	11.08	11.05	11.12
20	10.33	10.25	10.23	10.14	9.56	9.43	9.64	8.95	8.77	9.22	9.27	9.45	9.84	9.08	10.12	9.83	10.48	10.56	11.05	11.15	10.87
30	10.23	10.22	10.27	10.07	9.48	9.50	9.49	8.61	8.78	9.21	8.52	9.30	9.75	9.09	10.06	9.74	10.25	10.27	10.89	11.01	10.99
40	10.13	10.10	10.14	10.08	9.38	9.39	9.47	8.84	8.95	9.08	8.94	8.86	9.26	9.28	9.87	9.71	10.17	10.11	10.89	10.92	10.84
50	10.00	9.96	9.99	10.03	9.05	9.28	9.33	8.66	8.68	8.71	8.77	8.87	9.11	9.00	9.39	9.70	10.12	9.88	10.67	10.90	10.92
60	9.91	10.06	9.93	9.73	9.15	8.97	9.15	8.61	8.62	8.63	8.72	8.76	9.00	8.93	8.83	9.28	10.23	9.84	10.67	10.84	11.07
70	9.82	9.95	9.83	9.54	8.79	8.89	9.02	8.53	8.48	8.57	8.76	8.82	8.96	8.78	8.90	8.45	9.67	9.60	10.67	10.68	10.89
80	9.88	9.83	9.82	9.51	8.66	8.85	8.85	8.34	8.47	8.41	8.62	8.49	8.89	8.78	8.62	8.42	9.34	9.39	10.78	10.88	10.90
90	9.78	9.71	9.71	9.33	8.69	8.67	8.75	8.29	8.29	8.40	8.54	8.53	8.70	8.59	8.66	7.89	8.47	8.36	10.67	10.73	10.66
100	9.82	9.69	9.65	9.30	8.49	8.46	8.65	7.99	8.21	8.01	8.36	8.23	8.58	8.51	8.13	7.66	8.56	8.20	10.79	10.67	10.64
110	9.73	9.62	9.47	9.21	8.40	8.38	8.38	8.02	8.04	7.95	8.22	8.24	8.41	8.33	8.20	7.74	8.40	7.87	10.66	10.70	10.62
120	9.79	9.38	9.37	9.08	8.34	8.33	8.38	7.88	7.84	7.72	8.02	8.01	8.24	8.12	7.74	7.69	8.30	7.92	10.61	10.76	10.66
130	9.65	9.35	9.29	9.00	8.24	8.26	8.27	7.81	7.91	7.71	7.58	8.09	8.01	8.19	7.74	7.54	7.95	7.75	10.52	10.55	10.42
140	9.61	9.38	9.10	8.94	8.22	8.21	8.14	7.75	7.86	7.61	7.58	7.72	7.66	8.15	7.34	7.94	7.74	10.50	10.75	10.40	
150	9.65	9.13	9.02	8.69	7.96	7.82	7.89	7.45	7.25	7.35	7.25	7.32	7.50	7.18	7.39	7.58	7.55	10.46	10.46	10.54	10.37

Secchi depth

(m)	11	17.5	13	14.5	16	15.5	13.5	14.5	16	18.2	16.5	19	16	16	18.5	18	18.5	14.5	14	13	11

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 17 August 2005

2005-2006

Temperature

Date	17/08/2005	31/08/2005	14/09/2005	29/09/2005	12/10/2005	25/10/2005	10/11/2005	1/12/2005	10/01/2006	2/02/2006	1/03/2006	12/04/2006	27/04/2006	9/05/2006	30/05/2006	27/06/2006	11/07/2006	25/07/2006	4/09/2006
Depth (m)																			
0	11.17	11.74	12.42	11.91	11.92	13.40	16.10	15.09	17.40	20.20	19.50	16.71	16.31	15.70	14.21	11.94	11.51	11.15	11.10
10	10.98	11.24	11.76	11.68	11.79	12.84	14.59	14.93	17.10	20.11	19.50	16.72	16.29	15.70	14.21	11.99	11.51	11.15	10.93
20	10.97	11.10	11.22	11.67	11.76	12.17	14.27	14.22	16.85	18.15	19.25	16.72	16.29	15.70	14.21	11.99	11.50	11.15	10.93
30	10.97	11.05	11.05	11.66	11.66	11.63	12.36	13.34	14.84	15.46	16.14	16.71	16.29	15.70	14.21	11.99	11.48	11.15	10.89
40	10.97	11.00	11.01	11.60	11.47	11.47	11.66	12.32	12.21	13.40	12.93	16.48	13.96	13.40	14.20	11.99	11.48	11.15	10.87
50	10.97	10.98	10.98	11.18	11.39	11.29	11.27	11.66	11.60	11.75	11.57	12.00	12.20	11.94	14.16	11.99	11.48	11.15	10.83
60	10.97	10.97	10.99	11.02	11.37	11.17	11.15	11.26	11.21	11.35	11.35	11.53	11.56	11.36	11.54	11.39	11.47	11.15	10.82
70	10.96	10.97	10.97	10.97	11.26	11.06	11.04	11.11	11.13	11.19	11.16	11.29	11.30	11.23	11.27	11.21	11.46	11.15	10.82
80	10.97	10.96	10.97	10.97	11.13	10.99	11.00	11.06	11.06	11.11	11.14	11.19	11.19	11.14	11.19	11.16	11.45	11.15	10.82
90	10.96	10.96	10.96	10.96	11.07	10.97	10.98	11.01	11.05	11.06	11.06	11.12	11.12	11.10	11.16	11.15	11.42	11.15	10.81
100	10.96	10.95	10.96	10.95	11.01	10.97	10.97	10.98	11.04	11.04	11.05	11.08	11.08	11.09	11.12	11.14	11.23	11.15	10.81
110	10.96	10.94	10.94	10.94	10.98	10.94	10.95	10.97	11.02	11.02	11.05	11.05	11.07	11.06	11.11	11.14	11.20	11.15	10.81
120	10.96	10.94	10.93	10.93	10.98	10.94	10.94	10.97	11.00	11.02	11.05	11.03	11.06	11.06	11.09	11.13	11.19	11.15	10.80
130	10.96	10.93	10.93	10.92	10.96	10.93	10.93	10.96	10.99	11.00	11.03	11.02	11.05	11.04	11.07	11.13	11.18	11.15	10.79
140	10.95	10.93	10.91	10.91	10.96	10.93	10.94	10.96	10.99	11.00	11.02	11.04	11.03	11.07	11.12	11.18	11.15	10.76	
150	10.93	10.93	10.89	10.91	10.96	10.92	10.96	10.97	10.98	10.99	11.00	11.02	11.04	11.04	11.07	11.10	11.14	11.15	10.75

Dissolved Oxygen (g m⁻³)

Depth (m)																			
0	10.52	10.47	10.26	10.35	10.38	10.04	9.95	9.70	9.23	9.00	9.20	9.33	9.39	9.46	9.97	10.29	10.84	10.54	10.52
10	10.55	10.47	10.26	10.47	10.49	9.98	9.99	9.94	9.38	9.39	9.24	9.15	9.96	9.59	10.49	10.27	10.88	10.94	10.47
20	10.41	10.26	10.37	10.39	10.40	10.04	9.88	9.69	9.37	9.20	9.43	9.51	9.39	9.47	9.97	10.30	10.77	10.59	10.33
30	10.39	10.28	10.19	10.39	10.44	9.89	9.74	9.26	8.96	8.94	8.99	9.23	9.31	9.50	10.21	10.22	10.76	10.54	10.23
40	10.31	9.80	9.40	10.32	10.25	9.61	9.48	9.74	8.95	8.69	9.02	8.92	8.82	8.90	9.98	10.22	10.74	10.34	10.13
50	10.29	9.66	9.39	10.20	10.23	9.51	9.36	9.63	8.61	8.59	8.91	8.61	8.70	8.51	10.10	10.16	10.71	10.54	10.00
60	10.17	9.57	9.18	9.83	9.92	9.14	8.65	9.08	8.69	8.22	8.78	8.49	8.31	8.29	9.25	9.64	10.70	10.38	9.91
70	10.13	9.41	9.26	9.63	9.86	9.03	8.83	8.80	8.50	8.20	8.52	8.20	8.51	8.26	8.87	8.85	10.64	10.45	9.82
80	10.06	9.38	9.01	9.46	9.63	8.76	8.50	8.78	8.21	8.04	8.19	7.94	8.17	8.19	8.47	8.42	10.47	10.36	9.88
90	10.05	9.42	9.07	9.38	9.68	8.76	8.59	8.40	8.12	8.07	7.82	7.98	8.10	8.08	8.33	8.15	10.46	10.44	9.78
100	10.04	9.41	8.86	9.20	9.33	8.54	8.35	8.39	7.96	7.88	7.89	8.05	8.12	8.06	8.16	8.05	9.65	10.34	9.82
110	10.04	9.37	8.88	9.12	9.24	8.49	8.41	8.35	7.92	7.94	7.85	7.91	7.84	7.96	8.11	7.96	8.87	10.35	9.73
120	9.96	9.23	8.56	9.03	9.13	8.44	8.22	8.28	7.89	7.62	7.86	7.44	7.57	7.77	8.04	7.89	8.41	10.17	9.79
130	9.93	9.14	8.56	8.96	9.07	8.40	8.27	8.20	7.82	7.78	7.72	7.58	7.49	7.66	8.04	7.84	8.31	10.33	9.65
140	9.32	8.94	8.38	8.79	9.01	8.38	7.92	8.08	7.62	7.36	7.67	7.34	7.32	7.58	7.99	7.82	8.29	10.39	9.61
150	8.63	8.57	8.20	8.56	8.94	8.24	7.86	8.00	7.39	7.28	7.34	7.19	7.15	7.23	7.57	7.61	8.14	10.28	9.65

Secchi depth

(m)	13	13	13	14	14	15	17.5	19.3	19	15.5	15.3	15.8	17	17.5	18.2	15.2	13.5	12	11
-----	----	----	----	----	----	----	------	------	----	------	------	------	----	------	------	------	------	----	----

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 24 August 2004

2004-2005

Temperature

Depth (m)	24/08/2004	7/09/2004	21/10/2004	2/11/2004	22/11/2004	15/12/2004	11/01/2005	25/01/2005	9/02/2005	22/02/2005	10/03/2005	21/03/2005	14/04/2005	18/05/2005	9/06/2005	20/06/2005	20/07/2005	3/08/2005	17/08/2005	31/08/2005	14/09/2005
-----------	------------	-----------	------------	-----------	------------	------------	------------	------------	-----------	------------	------------	------------	------------	------------	-----------	------------	------------	-----------	------------	------------	------------

0	10.92	10.70	11.75	12.94	15.31	14.17	16.97	19.27	20.73	20.05	19.25	19.34	17.92	14.33	12.98	12.67	11.46	11.12	11.17	11.74	12.42
10	10.83	10.66	11.61	12.89	15.15	14.10	16.01	18.05	20.19	19.73	19.24	19.17	17.96	14.31	12.99	12.47	11.31	11.11	10.98	11.24	11.76
20	10.83	10.66	11.60	12.49	13.69	13.89	15.83	16.72	18.05	18.80	19.23	18.81	17.95	14.24	12.98	12.43	11.31	11.10	10.97	11.10	11.22
30	10.83	10.66	11.59	11.65	13.17	13.79	13.37	14.55	14.65	14.02	14.92	14.59	15.13	14.13	12.98	12.42	11.30	11.11	10.97	11.05	11.05
40	10.83	10.66	11.59	11.28	11.61	13.59	12.39	13.12	12.83	12.36	13.06	12.62	12.92	13.88	12.98	12.44	11.30	11.10	10.97	11.00	11.01
50	10.83	10.65	11.58	10.93	11.09	11.35	11.33	11.89	11.75	11.49	11.75	11.64	12.00	11.47	12.97	12.42	11.28	11.11	10.97	10.98	10.98
60	10.83	10.66	11.15	10.75	10.97	11.03	11.04	11.23	11.12	11.00	11.16	11.20	11.33	11.18	12.57	11.54	11.28	11.10	10.97	10.97	10.99
70	10.83	10.66	10.78	10.72	10.77	10.88	10.86	10.98	10.90	10.87	10.92	10.96	10.99	10.97	11.13	11.07	11.26	11.11	10.96	10.97	10.97
80	10.83	10.65	10.74	10.64	10.73	10.80	10.81	10.91	10.83	10.82	10.88	10.94	10.88	10.93	10.98	11.00	11.21	11.10	10.97	10.96	10.97
90	10.82	10.61	10.72	10.62	10.69	10.73	10.75	10.80	10.75	10.80	10.80	10.81	10.82	10.89	10.95	10.93	10.98	11.10	10.96	10.96	10.96
100	10.83	10.58	10.71	10.61	10.68	10.70	10.74	10.81	10.80	10.78	10.80	10.82	10.78	10.90	10.90	10.91	10.94	11.10	10.96	10.95	10.96
110	10.83	10.56	10.67	10.60	10.64	10.67	10.69	10.72	10.73	10.75	10.74	10.76	10.76	10.87	10.89	10.87	10.93	11.08	10.96	10.94	10.94
120	10.83	10.56	10.66	10.58	10.64	10.66	10.68	10.73	10.76	10.76	10.76	10.79	10.76	10.88	10.87	10.86	10.89	10.99	10.96	10.94	10.93
130	10.82	10.55	10.64	10.57	10.61	10.63	10.66	10.69	10.71	10.71	10.72	10.73	10.74	10.81	10.84	10.86	10.88	10.97	10.96	10.93	10.93
140	10.82	10.53	10.61	10.57	10.61	10.61	10.65	10.68	10.74	10.73	10.75	10.77	10.74	10.82	10.80	10.86	10.88	10.93	10.95	10.93	10.91
150	10.79	10.47	10.56	10.58	10.60	10.62	10.67	10.67	10.70	10.70	10.71	10.72	10.72	10.77	10.78	10.85	10.87	10.90	10.93	10.93	10.89

Dissolved Oxygen (g m⁻³)

Depth (m)	10.7	10.7	10.4	10.1	9.5	9.9	9.4	8.95	8.64	8.74	8.77	8.89	9.12	9.75	10.12	10.15	10.7	10.7	10.52	10.47	10.26
0	10.5	10.5	10.1	10.2	9.6	9.8	9.5	8.87	8.75	8.78	8.77	8.87	9.01	9.75	10.03	10.12	10.5	10.5	10.55	10.47	10.26
10	10.5	10.5	10.3	10.0	9.5	9.8	9.5	8.79	8.73	8.59	8.72	8.85	9.04	9.66	9.97	10.17	10.5	10.5	10.41	10.26	10.37
20	10.4	10.4	10.1	9.9	9.5	9.7	9.2	8.72	8.68	8.62	8.01	8.34	8.37	9.55	9.97	10.03	10.4	10.4	10.39	10.28	10.19
30	10.4	10.3	10.1	9.9	9.5	9.7	9.2	8.80	8.76	8.68	8.48	8.39	8.66	9.49	9.88	9.99	10.4	10.3	10.29	9.66	9.39
40	10.4	10.3	10.2	9.9	9.5	9.7	9.2	8.80	8.76	8.68	8.48	8.39	8.66	9.49	9.88	9.99	10.4	10.3	10.31	9.80	9.40
50	10.3	10.3	10.0	9.6	9.4	9.3	9.0	8.54	8.45	8.36	8.16	8.17	8.34	9.01	9.87	9.93	10.3	10.3	10.29	9.66	9.39
60	10.3	10.2	9.9	9.5	9.1	9.4	8.9	8.50	8.41	8.37	8.14	8.22	8.21	8.66	9.69	9.05	10.3	10.2	10.17	9.57	9.18
70	10.2	10.2	9.7	9.3	9.1	9.3	8.8	8.40	8.36	8.32	8.04	8.18	8.21	8.56	8.90	8.72	10.2	10.2	10.13	9.41	9.26
80	10.2	10.1	9.6	9.2	9.0	9.2	8.7	8.29	8.24	8.27	8.04	8.13	8.19	8.22	8.70	8.33	10.2	10.1	10.06	9.38	9.01
90	10.1	10.0	9.4	9.1	8.8	9.1	8.6	8.18	8.12	8.13	8.03	8.11	8.27	8.07	8.39	8.23	10.1	10.0	10.05	9.42	9.07
100	10.1	10.0	9.4	9.0	8.8	9.0	8.5	8.13	7.86	7.93	7.89	7.90	7.99	7.90	8.27	8.06	10.1	10.0	10.04	9.41	8.86
110	9.9	9.9	9.3	9.0	8.8	8.9	8.4	8.07	7.84	7.81	7.82	7.83	7.82	7.75	8.16	7.99	9.9	9.9	10.04	9.37	8.88
120	10.0	9.9	9.3	8.9	8.6	8.8	8.4	8.02	7.78	7.71	7.73	7.81	7.66	7.78	8.08	7.70	10.0	9.9	9.96	9.23	8.56
130	10.0	9.9	9.3	8.7	8.6	8.7	8.3	8.00	7.76	7.71	7.68	7.78	7.69	7.77	8.03	7.57	10.0	9.9	9.93	9.14	8.56
140	9.9	9.9	9.2	8.7	8.4	8.5	8.1	7.83	7.59	7.50	7.36	7.48	7.56	7.69	7.94	7.42	9.9	9.9	9.32	8.94	8.38
150	9.8	9.7	9.0	8.6	8.2	8.3	7.9	7.51	7.54	7.46	7.35	7.43	7.47	7.67	7.75	7.36	9.8	9.7	8.63	8.57	8.20

Secchi depth

(m)	12.5	12	15	16	16	19.5	20	19.5	18	21.5	18.5	20	17.2	16	14.1	13.8	13	14	13	13	13

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 14 July 2003
2003-2004
Temperature

Date	14/07/2003	31/07/2003	14/08/2003	26/08/2003	8/09/2003	7/10/2003	21/10/2003	19/11/2003	4/12/2003	18/12/2003	13/01/2004	26/02/2004	8/03/2004	31/03/2004	14/04/2004	10/05/2004	10/06/2004	13/07/2004	26/07/2004	24/08/2004	7/09/2004
Depth (m)																					
0	11.85	11.38	11.25	11.23	11.13	11.48	13.11	13.96	16.15	17.72	20.29	17.20	17.50	16.49	15.27	14.74	13.04	11.59	11.29	10.92	10.70
10	11.86	11.38	11.24	11.17	11.13	11.39	11.92	13.79	15.11	17.76	19.60	17.19	17.00	16.29	15.24	14.74	13.05	11.64	11.26	10.83	10.66
20	11.86	11.38	11.24	11.12	11.11	11.37	11.53	13.78	14.53	15.57	16.72	17.18	16.70	16.23	15.21	14.74	13.04	11.62	11.25	10.83	10.66
30	11.86	11.38	11.24	11.11	11.06	11.37	11.40	13.70	12.96	13.23	13.87	17.16	16.55	16.19	15.19	14.74	13.05	11.65	11.25	10.83	10.66
40	11.86	11.38	11.24	11.11	11.06	11.32	11.34	12.30	12.26	12.33	12.58	12.90	13.30	16.15	15.13	14.73	13.05	11.62	11.26	10.83	10.66
50	11.86	11.38	11.24	11.11	11.06	11.31	11.23	11.35	11.48	11.84	11.58	11.83	11.60	12.51	12.40	12.56	13.05	11.65	11.26	10.83	10.65
60	11.86	11.38	11.24	11.11	11.06	11.31	11.19	11.28	11.41	11.39	11.33	11.53	11.60	11.59	11.67	11.66	13.05	11.64	11.26	10.83	10.66
70	11.86	11.38	11.24	11.10	11.06	11.31	11.16	11.23	11.26	11.26	11.35	11.40	11.40	11.48	11.43	12.42	11.65	11.25	10.83	10.66	
80	11.35	11.38	11.24	11.00	11.06	11.30	11.15	11.19	11.25	11.22	11.23	11.30	11.35	11.34	11.39	11.38	11.56	11.64	11.25	10.83	10.65
90	11.31	11.38	11.24	11.09	11.06	11.29	11.13	11.16	11.20	11.17	11.22	11.25	11.27	11.30	11.32	11.35	11.51	11.66	11.25	10.82	10.61
100	11.27	11.35	11.24	11.09	11.06	11.25	11.11	11.15	11.18	11.17	11.21	11.23	11.27	11.27	11.30	11.32	11.39	11.65	11.25	10.83	10.58
110	11.24	11.34	11.23	11.09	11.06	11.21	11.10	11.12	11.17	11.15	11.19	11.20	11.24	11.26	11.28	11.30	11.35	11.65	11.26	10.83	10.56
120	11.22	11.32	11.22	11.09	11.06	11.14	11.10	11.11	11.18	11.14	11.18	11.18	11.22	11.24	11.25	11.30	11.34	11.65	11.26	10.83	10.56
130	11.21	11.27	11.22	11.08	11.06	11.11	11.08	11.09	11.14	11.13	11.17	11.18	11.20	11.22	11.23	11.28	11.33	11.49	11.26	10.82	10.55
140	11.21	11.26	11.21	11.08	11.06	11.09	11.08	11.09	11.15	11.13	11.16	11.17	11.20	11.21	11.27	11.32	11.39	11.26	10.82	10.53	
150	11.20	11.22	11.20	11.08	11.07	11.09	11.08	11.09	11.14	11.13	11.16	11.17	11.20	11.21	11.26	11.31	11.34	11.26	10.79	10.47	

Dissolved Oxygen (g m⁻³)

Depth (m)																					
0	10.3	10.6	10.5	10.5	10.5	10.1	9.9	9.5	9.1	9.2	9.3	9.4	9.2	9.5	9.7	10.2	10.5	10.6	10.7	10.7	
10	10.2	10.4	10.5	10.5	10.6	10.5	10.0	9.9	9.5	9.2	9.3	9.4	9.0	9.1	9.2	9.6	9.9	10.5	10.6	10.5	10.5
20	10.2	10.2	10.3	10.4	10.4	10.4	10.2	9.8	9.4	9.0	9.1	9.0	8.8	9.0	9.1	9.4	9.8	10.5	10.6	10.5	10.5
30	10.2	9.9	10.1	10.3	10.1	10.1	10.0	9.5	9.2	9.2	9.1	8.9	8.5	9.0	8.8	9.3	9.5	10.3	10.3	10.4	10.4
40	10.1	9.9	10.0	10.0	9.8	10.0	9.7	9.3	9.0	9.1	8.7	8.4	8.0	8.9	8.8	9.2	9.5	10.1	10.1	10.4	10.3
50	10.0	9.0	9.9	9.9	9.8	9.8	9.4	9.0	8.7	8.8	8.5	8.1	7.9	8.2	8.2	8.6	9.4	9.8	9.9	10.3	10.3
60	9.9	8.8	9.8	9.7	9.6	9.7	9.2	8.9	8.6	8.4	8.2	8.0	7.7	8.0	8.0	8.2	9.4	9.9	9.8	10.3	10.2
70	9.9	8.7	9.8	9.6	9.6	9.6	9.1	8.7	8.5	8.3	8.1	7.9	7.6	8.0	7.8	7.9	9.1	9.6	9.7	10.2	10.2
80	8.7	8.6	9.7	9.5	9.5	9.6	8.9	8.6	8.4	8.1	8.0	7.9	7.5	8.0	7.7	7.9	8.5	9.7	9.6	10.2	10.1
90	8.5	8.5	9.7	9.5	9.5	9.5	8.9	8.6	8.3	8.1	8.0	7.9	7.5	7.9	7.6	7.8	8.0	9.5	9.5	10.1	10.0
100	8.2	8.4	9.6	9.5	9.5	9.4	8.8	8.6	8.2	7.9	7.8	7.8	7.4	7.8	7.5	7.7	7.7	9.5	9.4	10.1	10.0
110	8.2	8.1	9.6	9.4	9.5	9.3	8.8	8.4	8.2	7.9	7.8	7.7	7.3	7.7	7.4	7.6	7.6	9.4	9.4	9.9	9.9
120	8.0	8.0	9.5	9.4	9.5	9.3	8.7	8.4	8.1	7.8	7.7	7.5	7.1	7.6	7.3	7.4	7.5	9.4	9.3	10.0	9.9
130	8.0	7.9	9.5	9.4	9.4	9.1	8.7	8.3	8.0	7.8	7.5	7.3	7.0	7.5	7.2	7.3	7.4	9.1	9.2	10.0	9.9
140	7.8	7.8	9.5	9.3	9.4	9.0	8.5	8.2	7.9	7.5	7.4	7.3	6.9	7.4	7.0	7.3	7.3	8.3	9.2	9.9	9.9
150	7.7	7.6	9.3	9.4	8.9	8.5	8.0	7.7	7.3	7.2	7.1	6.8	7.1	6.8	7.1	7.3	8.0	9.2	9.8	9.7	

Secchi depth

(m)	14.5	14	13.5	13	12.5	13	17	16	18.5	17.5	19	17	15	16	15	18	13.5	12	11	12.5	12
-----	------	----	------	----	------	----	----	----	------	------	----	----	----	----	----	----	------	----	----	------	----

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Additional site B (Kuratau Basin) for the period starting 14 July 2003

2003-2004

Temperature

Date	14/07/2003	31/07/2003	14/08/2003	26/08/2003	8/09/2003	7/10/2003	21/10/2003	19/11/2003	4/12/2003	18/12/2003	13/01/2004	26/02/2004	8/03/2004	31/03/2004	14/04/2004	10/05/2004	10/06/2004	13/07/2004	26/07/2004	24/08/2004	7/09/2004
Depth (m)																					
0	11.82	11.32	11.38	11.36	11.13	11.70	13.31	13.79	15.65	17.08	20.25	16.83	17.63	15.92	15.10	14.72	13.02	11.43	11.26	10.92	10.85
10	11.80	11.29	11.22	11.17	11.11	11.44	12.28	13.49	15.00	16.43	19.73	16.72	16.56	15.90	15.02	14.68	12.95	11.40	11.20	10.77	10.59
20	11.79	11.29	11.22	11.14	11.07	11.40	11.71	13.33	13.81	15.28	16.73	16.58	16.51	15.89	15.00	14.64	12.84	11.41	11.20	10.73	10.58
30	11.79	11.29	11.21	11.13	11.03	11.35	11.46	12.22	12.37	13.38	13.74	16.16	16.40	15.88	14.99	14.47	12.71	11.41	11.20	10.72	10.57
40	11.79	11.29	11.21	11.13	11.02	11.34	11.38	11.67	11.90	12.91	12.48	15.75	15.53	15.53	14.18	14.07	12.67	11.41	11.19	10.72	10.57
50	11.79	11.29	11.21	11.13	11.02	11.33	11.28	11.40	11.57	11.65	11.62	12.97	12.55	12.89	12.48	12.48	12.66	11.41	11.19	10.72	10.56
60	11.78	11.29	11.21	11.13	11.01	11.25	11.23	11.31	11.37	11.33	11.40	11.88	11.64	11.69	11.72	11.78	12.57	11.40	11.19	10.72	10.56
70	11.78	11.29	11.21	11.12	11.01	11.12	11.15	11.24	11.25	11.27	11.28	11.55	11.47	11.49	11.51	11.47	12.51	11.41	11.18	10.72	10.56
80	11.77	11.29	11.16	11.12	11.01	11.06	11.09	11.18	11.21	11.25	11.20	11.38	11.41	11.37	11.43	11.38	12.27	11.37	11.18	10.72	10.51
90	11.35	11.29	11.04	11.11	11.01	11.02	11.08	11.13	11.13	11.19	11.16	11.32	11.35	11.32	11.37	11.31	11.77	11.26	11.17	10.71	10.45
100	11.27	11.29	10.91	11.08	11.01	11.02	11.05	11.10	11.11	11.16	11.14	11.28	11.33	11.26	11.30	11.24	11.65	11.24	11.17	10.66	10.38

Dissolved Oxygen (g m⁻³)

Depth (m)																						
0	10.7	10.9	10.8	10.6	10.6	10.4	10.5	10.1	9.8	9.1	9.2	9.3	9.5	8.8	10.5	11.4	12.3	10.6	10.5	10.5	10.8	
10	10.5	11.0	10.6	10.6	10.5	10.4	10.4	10.3	9.9	9.3	9.2	9.1	9.0	9.0	9.5	10.2	10.7	10.6	10.5	10.4	10.7	
20	10.3	11.3	10.4	10.2	10.2	10.2	10.1	9.9	9.6	9.4	9.2	9.0	8.9	8.9	9.2	9.9	10.1	10.1	10.5	10.5	10.7	
30	10.2	11.2	10.1	9.9	10.1	9.9	10.0	9.6	9.3	9.1	9.0	9.0	8.7	8.8	8.9	9.4	9.7	9.8	10.3	10.4	10.6	
40	10.1	11.2	9.9	9.8	9.9	9.6	9.7	9.2	8.9	9.1	8.8	8.7	8.2	8.7	8.5	9.1	9.6	9.6	10.0	10.3	10.5	
50	10.0	10.9	9.8	9.6	9.8	9.6	9.4	9.0	8.8	8.7	8.5	8.2	7.9	8.2	7.9	8.5	9.3	9.5	9.8	10.2	10.3	
60	9.9	10.7	9.7	9.5	9.7	9.4	9.0	8.8	8.6	8.3	8.2	8.1	7.7	8.0	7.6	8.0	9.2	9.3	9.6	10.1	10.3	
70	9.9	10.4	9.7	9.5	9.7	9.3	8.9	8.7	8.6	8.3	8.1	7.9	7.6	7.8	7.3	7.7	8.9	9.2	9.6	10.1	10.2	
80	9.8	10.3	9.4	9.4	9.6	9.1	8.7	8.6	8.4	7.9	7.8	7.8	7.4	7.6	7.1	7.4	8.7	9.1	9.4	10.0	10.1	
90	9.2	10.1	9.2	9.3	9.6	9.0	8.7	8.5	8.3	7.9	7.8	7.7	7.3	7.6	7.0	7.5	8.3	8.7	9.5	9.9	10.1	
100	8.3	10.0	9.2	9.3	9.6	8.9	8.6	8.2	7.9	7.9	7.6	7.4	7.3	7.3	6.8	7.0	8.1	8.1	9.4	9.8	10.0	

Secchi depth

(m)	12	13	13	11.5	11	9.5	15	17	17	15	16	13.5	5	11	14	15.5	12	11	10	10	11

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Additional site C (Western Bays) for the period starting 14 July 2003

2003-2004

Temperature

Date	14/07/2003	31/07/2003	14/08/2003	26/08/2003	8/09/2003	7/10/2003	21/10/2003	19/11/2003	4/12/2003	18/12/2003	13/01/2004	26/02/2004	8/03/2004	31/03/2004	14/04/2004	10/05/2004	10/06/2004	13/07/2004	26/07/2004	24/08/2004	7/09/2004
Depth (m)																					
0	11.86	11.43	11.56	11.31	11.32	11.85	13.29	15.10	15.79	17.00	20.17	16.90	18.43	16.37	15.41	14.98	13.16	11.58	11.51	10.97	11.14
10	11.80	11.36	11.26	11.21	11.13	11.24	11.93	13.84	15.29	16.33	18.89	16.69	17.02	16.35	15.18	14.80	13.08	11.61	11.32	10.94	10.73
20	11.80	11.34	11.25	11.14	11.09	11.17	11.62	13.76	14.31	15.26	17.11	16.34	16.45	16.35	15.15	14.76	13.07	11.61	11.30	10.90	10.71
30	11.80	11.32	11.25	11.14	11.08	11.14	11.52	13.63	12.99	13.46	13.74	14.66	15.33	15.95	15.15	14.75	13.07	11.61	11.31	10.90	10.71
40	11.80	11.31	11.25	11.14	11.08	11.14	11.50	11.91	12.03	12.88	12.25	12.56	13.64	13.21	15.14	14.73	13.07	11.60	11.31	10.89	10.70
50	11.80	11.31	11.25	11.14	11.07	11.13	11.46	11.42	11.43	11.64	11.57	11.63	11.64	11.68	12.68	12.57	12.80	11.61	11.30	10.90	10.70
60	11.80	11.31	11.25	11.14	11.07	11.13	11.38	11.31	11.30	11.31	11.36	11.53	11.48	11.45	11.76	11.73	11.68	11.60	11.30	10.89	10.70
70	11.80	11.31	11.25	11.14	11.07	11.12	11.21	11.27	11.28	11.26	11.28	11.39	11.37	11.34	11.54	11.48	11.44	11.61	11.30	10.89	10.70
80	11.79	11.31	11.25	11.14	11.07	1.10	11.13	11.20	11.25	11.22	11.25	11.31	11.35	11.32	11.37	11.39	11.37	11.58	11.30	10.89	10.70
90	11.60	11.29	11.25	11.14	11.07	11.04	11.07	11.14	11.21	11.19	11.21	11.26	11.33	11.29	11.30	11.32	11.33	11.61	11.30	10.89	10.70
100	11.28	11.27	11.24	11.14	11.07	11.03	11.07	11.11	11.19	11.12	11.19	11.23	11.32	11.25	11.29	11.31	11.32	11.61	11.30	10.89	10.70

Dissolved Oxygen (g m⁻³)

Depth (m)	0	10.3	10.7	10.3	10.4	10.4	11.4	10.1	9.8	9.5	9.2	9.2	9.3	9.3	9.4	10.4	10.3	10.6	10.6	11.0	10.4	10.7
10	10.3	10.8	10.3	10.3	10.4	10.4	11.0	10.1	9.9	9.9	9.1	9.2	9.1	9.0	9.2	9.5	9.8	10.1	10.6	10.5	10.4	10.4
20	10.1	10.3	10.1	10.1	10.2	10.8	9.9	9.9	9.5	9.2	9.1	9.2	9.1	9.0	9.1	9.7	9.9	10.6	10.2	10.3	10.4	
30	10.1	10.0	9.9	9.9	10.0	10.1	9.6	9.6	9.3	9.1	8.8	8.6	8.6	8.9	8.9	9.4	9.7	10.3	9.9	10.2	10.4	
40	10.0	10.0	9.8	9.7	9.9	9.7	9.4	9.4	9.0	9.1	8.8	8.4	8.4	8.3	8.7	9.2	9.6	9.9	9.8	10.1	10.3	
50	9.9	9.9	9.6	9.6	9.7	9.7	9.3	9.2	8.8	8.8	8.5	8.2	8.0	8.0	8.2	8.7	9.3	9.6	9.6	10.1	10.2	
60	9.8	9.6	9.6	9.5	9.6	9.5	9.2	9.0	8.5	8.5	8.2	8.0	7.9	8.0	7.8	8.2	8.6	9.5	10.1	10.2		
70	9.8	9.5	9.5	9.4	9.5	9.4	9.1	8.8	8.5	8.3	8.1	7.9	7.8	7.9	7.5	8.0	8.2	9.4	9.5	10.0	10.1	
80	9.7	9.5	9.5	9.4	9.5	9.3	8.8	8.8	8.3	8.2	7.9	7.8	7.8	7.4	7.8	8.0	9.3	9.4	10.0	10.0		
90	9.6	9.1	9.4	9.3	9.4	9.2	8.7	8.6	8.4	7.9	7.8	7.7	7.7	7.3	7.6	7.9	9.2	9.9	10.0			
100	8.8	8.8	9.0	9.3	9.4	9.1	8.7	8.5	8.3	7.9	7.7	7.6	7.7	7.5	7.3	7.5	7.8	9.1	9.3	9.9	10.0	

Secchi depth

(m)	14	12	14.5	13	12	12.5	12	17.2	17	19	17.5	14	13	12.5	16.5	16	14	12.5	11	10	12
-----	----	----	------	----	----	------	----	------	----	----	------	----	----	------	------	----	----	------	----	----	----

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Mid-Lake site A for the period starting 1 July 2002
2002-2003
Temperature

Date	1/07/2002	17/07/2002	31/07/2002	29/08/2002	18/09/2002	9/10/2002	13/11/2002	28/11/2002	18/12/2002	30/01/2003	13/02/2003	17/03/2003	3/04/2003	28/04/2003	15/05/2003	12/06/2003	14/07/2003	31/07/2003	14/08/2003	26/08/2003	8/09/2003
Depth (m)																					
0	12.13	11.44	11.20	11.10	11.38	11.60	12.58	14.12	15.00	17.84	19.31	18.55	19.05	16.76	15.67	13.59	11.85	11.38	11.25	11.23	11.13
10	12.12	11.44	11.20	10.90	11.33	11.60	12.55	14.02	14.78	17.59	19.19	18.43	18.70	16.73	15.57	13.56	11.86	11.38	11.24	11.17	11.13
20	12.11	11.44	11.20	10.90	11.28	11.40	12.50	12.91	14.48	17.08	18.10	18.37	18.59	16.73	15.56	13.55	11.86	11.38	11.24	11.12	11.11
30	12.11	11.44	11.20	10.80	11.02	11.30	12.38	12.41	14.26	16.13	15.50	16.77	17.02	16.72	15.57	13.55	11.86	11.38	11.24	11.11	11.06
40	12.11	11.44	11.20	10.90	10.97	11.30	12.16	11.98	12.67	12.69	12.85	13.44	13.31	12.80	15.53	12.22	11.86	11.38	11.24	11.11	11.06
50	12.11	11.44	11.20	10.90	10.96	11.20	12.00	11.54	11.87	12.03	12.14	12.03	12.30	11.96	12.20	11.82	11.86	11.38	11.24	11.11	11.06
60	12.10	11.44	11.20	10.80	10.94	11.20	11.72	11.22	11.64	11.70	11.68	11.60	11.81	11.62	11.61	11.52	11.86	11.38	11.24	11.11	11.06
70	12.10	11.44	11.20	10.80	10.93	11.20	11.51	11.09	11.31	11.41	11.33	11.39	11.52	11.34	11.36	11.38	11.86	11.38	11.24	11.10	11.06
80	11.97	11.44	11.20	10.90	10.92	11.10	11.32	10.98	11.17	11.25	11.25	11.27	11.31	11.27	11.27	11.27	11.35	11.38	11.24	11.00	11.06
90	11.49	11.43	11.20	10.90	10.91	11.10	11.13	10.95	11.06	11.15	11.16	11.16	11.20	11.17	11.22	11.21	11.31	11.38	11.24	11.09	11.06
100	11.39	11.41	11.20	10.90	10.90	11.10	11.05	10.92	11.04	11.11	11.10	11.13	11.18	11.15	11.20	11.20	11.27	11.35	11.24	11.09	11.06
110	11.32	11.37	11.20	10.90	10.89	11.00	11.05	10.90	11.04	11.09	11.08	11.10	11.13	11.13	11.16	11.17	11.24	11.34	11.23	11.09	11.06
120	11.29	11.32	11.20	10.90	10.87	11.00	11.01	10.87	11.00	11.06	11.06	11.09	11.13	11.13	11.15	11.15	11.22	11.32	11.22	11.09	11.06
130	11.25	11.27	11.20	10.90	10.85	10.90	10.99	10.85	10.98	11.04	11.04	11.08	11.09	11.10	11.12	11.12	11.21	11.27	11.22	11.08	11.06
140	11.23	11.26	11.20	10.80	10.83	10.90	10.97	10.83	10.97	11.03	11.03	11.09	11.09	11.09	11.12	11.11	11.21	11.26	11.21	11.08	11.06
150	11.23	11.26	11.20	10.80	10.81	10.90	10.96	10.82	10.97	11.03	11.03	11.07	11.08	11.09	11.11	11.11	11.20	11.22	11.20	11.08	11.07

Dissolved Oxygen (g m⁻³)

Depth (m)	0	10.3	10.4	9.7	10.5	10.5	10.3	10.2	9.8	9.6	9.1	8.9	9.0	8.8	9.2	9.5	10.0	10.3	10.6	10.5	10.5
10	10.3	10.7	9.5	10.4	10.7	10.3	10.2	10.0	9.7	9.1	8.9	8.9	8.8	8.8	9.2	9.2	9.7	10.2	10.4	10.5	10.6
20	10.3	10.7	9.4	10.3	10.6	10.2	10.2	10.1	9.6	9.2	8.9	8.8	8.6	8.6	9.1	9.3	9.4	10.2	10.2	10.3	10.4
30	10.2	10.7	9.4	10.3	10.5	10.2	10.2	10.1	9.6	9.1	8.8	8.5	8.3	8.9	9.2	9.3	10.2	9.9	10.1	10.3	10.1
40	10.2	10.6	9.4	10.2	10.4	10.2	10.1	9.7	9.5	9.2	8.8	8.4	8.4	8.0	8.4	9.1	9.0	10.1	9.9	10.0	9.8
50	10.2	10.6	9.4	10.2	10.3	10.1	10.1	9.7	9.3	9.1	8.6	8.2	7.8	8.2	8.2	8.2	10.0	9.0	9.9	9.9	9.8
60	10.1	10.5	9.4	10.2	10.2	10.1	10.0	9.5	9.1	8.9	8.4	8.0	7.7	8.1	8.1	8.1	9.9	8.8	9.8	9.7	9.6
70	10.1	10.5	9.3	10.1	10.2	10.0	9.9	9.5	8.8	8.8	8.4	7.8	7.6	8.0	8.0	8.0	9.9	8.7	9.8	9.6	9.6
80	10.0	10.3	9.4	10.1	10.2	10.1	9.7	9.4	8.7	8.7	8.3	7.8	7.5	7.9	7.8	7.9	8.7	8.6	9.7	9.5	9.5
90	9.7	10.3	9.4	10.1	10.1	9.5	9.3	8.7	8.7	8.2	7.8	7.4	7.8	7.5	7.6	8.5	8.5	9.7	9.5	9.5	9.5
100	8.6	10.1	9.4	10.1	10.0	9.8	9.4	9.1	8.6	8.6	8.1	7.7	7.3	7.7	7.2	7.5	8.2	8.4	9.6	9.5	9.5
110	8.3	9.8	9.3	9.9	9.9	9.8	9.4	9.1	8.4	8.4	8.0	7.6	7.2	7.6	7.1	7.4	8.2	8.1	9.6	9.4	9.5
120	8.1	8.8	9.3	9.9	9.9	9.8	9.3	9.0	8.3	8.3	7.8	7.4	7.0	7.5	7.1	7.2	8.0	8.0	9.5	9.4	9.5
130	8.0	8.5	9.3	9.9	9.9	9.7	9.2	9.0	8.3	8.2	7.7	7.2	6.9	7.4	7.0	7.0	8.0	7.9	9.5	9.4	9.4
140	7.8	8.1	9.3	9.9	9.9	9.4	9.0	8.8	8.2	8.0	7.4	7.1	6.8	7.2	6.8	6.7	7.8	7.8	9.5	9.3	9.4
150	7.8	8.1	9.3	9.8	9.8	9.4	8.9	8.7	8.1	7.9	7.3	6.9	6.5	6.9	6.7	6.5	7.7	7.6	9.3	9.3	9.4

Secchi depth

(m)	16	15.5	12	9.5	12	15.5	18	12.7	13.5	18	19	15	13.5	14	16.5	11	14.5	14	13.5	13	12.5
-----	----	------	----	-----	----	------	----	------	------	----	----	----	------	----	------	----	------	----	------	----	------

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Additional site B (Kuratau Basin) for the period starting 1 July 2002

2002-2003

Temperature

Date	1/07/2002	17/07/2002	31/07/2002	29/08/2002	18/09/2002	9/10/2002	13/11/2002	28/11/2002	18/12/2002	30/01/2003	13/02/2003	17/03/2003	3/04/2003	28/04/2003	15/05/2003	12/06/2003	14/07/2003	31/07/2003	14/08/2003	26/08/2003	8/09/2003
Depth (m)																					
0	12.13	11.48	11.3	11	11.08	11.70	11.98	13.82	15.16	16.76	18.87	18.74	19.09	16.73	15.79	13.24	11.82	11.32	11.38	11.36	11.13
10	12.09	11.49	11.1	10.8	11.05	11.30	11.94	13.67	15.08	16.75	18.46	18.54	18.82	16.66	15.49	13.02	11.8	11.29	11.22	11.17	11.11
20	12.09	11.48	11.1	10.8	11.03	11.20	11.9	12.79	13.86	16.53	17.71	18.45	18.49	16.62	15.47	12.79	11.79	11.29	11.22	11.14	11.07
30	12.09	11.48	11.1	10.8	11.03	11.20	11.8	12.31	13.4	14.33	16.2	14.87	15.32	16.2	15.41	11.83	11.79	11.29	11.21	11.13	11.03
40	12.08	11.48	11.1	10.8	11.02	11.20	11.68	11.75	13.18	12.98	13.89	12.03	13.25	13.46	13.2	11.62	11.79	11.29	11.21	11.13	11.02
50	11.97	11.49	11.1	10.8	10.91	11.20	11.44	11.44	12.91	12.1	12.59	12.06	12	12.28	12.09	11.51	11.79	11.29	11.21	11.13	11.02
60	11.93	11.49	11.1	10.8	10.9	11.10	11.26	11.27	12.27	11.69	11.75	11.58	11.58	11.7	11.71	11.38	11.78	11.29	11.21	11.13	11.01
70	11.87	11.48	11.1	10.8	10.89	11.10	11.11	11.17	11.58	11.37	11.4	11.36	11.35	11.4	11.4	11.29	11.78	11.29	11.21	11.12	11.01
80	11.78	11.48	11.1	10.8	10.89	11.00	11	11.03	11.51	11.23	11.3	11.24	11.25	11.25	11.28	11.27	11.77	11.29	11.16	11.12	11.01
90	11.37	11.46	11.1	10.7	10.87	11.00	10.93	10.96	11.39	11.14	11.17	11.13	11.15	11.18	11.21	11.26	11.35	11.29	11.04	11.11	11.01
100	11.28	11.3	11	10.7	10.85	11.00	10.91	10.92	11.2	11.09	11.12	11.13	11.12	11.12	11.18	11.25	11.27	11.29	10.91	11.08	11.01
110					10.7	10.7		10.90													

Dissolved Oxygen (g m⁻³)

Depth (m)																						
0	10.3	10.4	9.9	10.4	10.4	10.3	9.9	9.6	9.3	9.4	8.9	8.9	9.7	9.4	10	10.7	10.9	10.8	10.6	10.6	10.6	
10	10.3	10.8	9.7	10.3	10.5	10.5	10.3	10	9.7	9.3	9.3	8.9	8.8	9.6	9.4	10	10.5	11	10.6	10.6	10.5	
20	10.2	10.6	9.6	10.3	10.5	10.3	10.3	9.9	9.5	9.2	9.3	8.8	8.5	9.5	9.3	9.6	10.3	11.3	10.4	10.2	10.2	
30	10.2	10.6	9.6	10.2	10.5	10.3	10.3	103	9.9	9.6	9.2	9.2	8.2	8.1	9.4	8.8	9.2	10.2	11.2	10.1	9.9	10.1
40	10.1	10.5	9.6	10.2	10.4	10.2	10.2	10.2	9.5	9.4	9.1	9	8.2	8	8.8	8.5	8.8	10.1	11.2	9.9	9.8	9.9
50	10.1	10.5	9.6	10.1	10.3	10.1	10.1	9.5	9.4	8.9	8.8	8	7.7	8.3	7.9	8.5	10	10.9	9.8	9.6	9.8	9.8
60	9.8	10.4	9.6	10.1	10.2	10.1	10.1	9.9	9.4	9.2	8.6	8.6	7.8	7.6	8.3	7.8	8.3	9.9	10.7	9.7	9.5	9.7
70	9.7	10.4	9.5	10	10.1	9.8	9.8	9.4	9	8.4	8.4	7.7	7.4	8.2	7.7	8.2	9.9	10.4	9.7	9.5	9.7	9.7
80	9.5	10.3	9.5	10	10.1	9.7	9.7	9	8.6	8.3	8.3	7.3	7.3	8	7.7	8.1	9.8	10.3	9.4	9.4	9.6	9.6
90	9.1	10.3	9.5	10	10	9.7	9.5	9	8.6	8.2	8	7.2	7.1	7.7	7.5	7.7	9.2	10.1	9.2	9.3	9.6	9.6
100	8.7	9.8	9.6	9.9	9.9	9.7	9.2	9	8.4	7.7	7.6	7	7	7.6	7.1	7.5	8.3	10	9.2	9.3	9.6	9.6
110			9.2	9.8		9.4																

Secchi depth

(m)	16	12.5	10.5	8	11	16	14	12.7	14	18	11	14	12.8	13.5	15.5	12	12	13	13	11.5	11
-----	----	------	------	---	----	----	----	------	----	----	----	----	------	------	------	----	----	----	----	------	----

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Additional site C (Western Bays) for the period starting 1 July 2002

2002-2003

Temperature

Date	1/07/2002	17/07/2002	31/07/2002	29/08/2002	18/09/2002	9/10/2002	13/11/2002	28/11/2002	18/12/2002	30/01/2003	13/02/2003	17/03/2003	3/04/2003	28/04/2003	15/05/2003	12/06/2003	14/07/2003	31/07/2003	14/08/2003	26/08/2003	8/09/2003
Depth (m)																					
0	12.22	11.52	11.6	11.4	11.24	12.10	12.56	13.98	15.12	17.61	19.58	19.04	18.15	17.1	15.8	13.65	11.86	11.43	11.56	11.31	11.32
10	12.15	11.5	11.2	10.9	11.23	11.30	12.5	13.45	14.21	17.49	18.95	18.45	18.58	16.82	15.54	13.62	11.8	11.36	11.26	11.21	11.13
20	12.14	11.49	11.2	10.9	11.16	11.30	12.38	12.63	13.31	17.48	17.41	18.29	18.3	16.77	15.52	13.59	11.8	11.34	11.25	11.14	11.09
30	12.14	11.49	11.2	10.8	11.06	11.20	12.33	12.42	12.73	14.31	14.19	14.81	14.61	16.76	15.51	13.59	11.8	11.32	11.25	11.14	11.08
40	12.13	11.49	11.2	10.8	11.02	11.20	11.75	12.2	11.98	12.36	12.79	12.88	12.73	13.62	13.07	13.59	11.8	11.31	11.25	11.14	11.08
50	12.13	11.49	11.2	10.8	11.02	11.20	11.28	11.98	11.53	12	11.98	11.86	12.1	12.08	12.14	13.54	11.8	11.31	11.25	11.14	11.07
60	11.92	11.49	11.2	10.8	11	11.10	11.12	11.37	11.33	11.61	11.68	11.49	11.71	11.56	11.71	13.28	11.8	11.31	11.25	11.14	11.07
70	11.55	11.49	11.2	10.8	10.99	11.10	11.08	11.21	11.15	11.29	11.3	11.35	11.37	11.35	11.4	11.8	11.8	11.31	11.25	11.14	11.07
80	11.5	11.49	11.2	10.8	10.95	11.10	11.03	11.04	11.12	11.19	11.19	11.25	11.22	11.24	11.27	11.45	11.79	11.31	11.25	11.14	11.07
90	11.47	11.49	11.2	10.8	10.94	11.00	11	10.98	11.1	11.11	11.15	11.2	11.18	11.18	11.22	11.35	11.6	11.29	11.25	11.14	11.07
100	11.45	11.49	11.2	10.8	10.92	11.00	10.97	10.96	11.08	11.08	11.13	11.2	11.15	11.15	11.17	11.23	11.28	11.27	11.24	11.14	11.07

Dissolved Oxygen (g m⁻³)

Depth (m)	0	10.4	10.5	9.7	10.3	10.5	10.4	10.2	9.9	9.6	9.1	9.5	9.9	8.9	9.4	9.3	10	10.3	10.7	10.3	10.4	10.4
10	10.4	10.8	9.5	10.2	10.7	10.4	10.3	9.7	9.6	9	9.3	9.7	8.8	9.2	9.1	9.6	10.3	10.8	10.3	10.3	10.4	10.4
20	10.4	10.8	9.5	10.2	10.7	10.4	10.3	9.9	9.7	9	9.3	9	8.8	9.2	9	9.3	10.1	10.3	10.1	10.1	10.2	10.2
30	10.3	10.7	9.4	10.1	10.6	10.4	10.2	9.9	9.6	8.7	9	8.4	8.3	9	8.8	9.1	10.1	10	9.9	9.9	9.9	10
40	10.3	10.5	9.4	10	10.5	10.3	10.1	9.7	9.5	8.7	9	8.4	8.1	8.5	8.3	9.3	10	10	9.8	9.7	9.9	9.9
50	10.2	10.5	9.4	10	10.4	10	9.9	9.7	9.2	8.6	8.7	8.1	7.9	8.2	7.8	9.2	9.9	9.9	9.6	9.6	9.7	9.7
60	10	10.5	9.4	10	10.4	10	9.7	9.6	9.1	8.5	8.5	8.1	7.9	8.2	7.8	9.9	9.8	9.6	9.6	9.5	9.6	9.6
70	9.6	10.5	9.4	9.9	10.3	9.9	9.7	9.5	9	8.4	8.4	7.9	7.8	8	7.7	9.7	9.8	9.5	9.5	9.4	9.5	9.5
80	8.8	10.5	9.3	9.9	10.2	9.9	9.5	9	8.8	8.3	8.3	7.6	7.7	8	7.5	9.4	9.7	9.5	9.5	9.4	9.5	9.5
90	8.7	10.4	9.3	9.9	10.1	9.8	9.5	9.1	8.7	8.1	8.3	7.5	7.6	7.9	7.3	9.2	9.6	9.1	9.4	9.3	9.4	9.4
100	8.6	10.2	9.3	10	10	9.6	9.3	9.1	8.7	8	8.1	7.3	7.4	7.8	7.2	9.1	8.8	8.8	9	9.3	9.4	9.4

Secchi depth

(m)	14	12.5	12	8	12	19	16	15.5	13.5	18.5	19	15	14.5	14.5	17	11	14	12	14.5	13	12
-----	----	------	----	---	----	----	----	------	------	------	----	----	------	------	----	----	----	----	------	----	----

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
2001-2002
Mid-Lake site A for the period starting 2 July 2001
Temperature

Date	2/07/01	25/07/01	13/08/01	3/09/01	25/09/01	25/10/01	12/11/01	10/12/01	20/12/01	8/01/02	22/01/02	6/03/02	4/04/02	22/04/02	5/05/02	19/06/02	1/07/02	17/07/02	31/07/02	29/08/02	18/09/02	9/10/02
Depth (m)																						
0	12.11	11.26	11.15	10.96	11.58	12.97	14.23	15.47	17.92	18.37	19.4	18.69	17.45	17.05	15.51	12.57	12.13	11.44	11.2	11.1	11.38	11.60
10	12.04	11.26	11.12	10.98	11.57	12.91	14.16	15.51	16.60	18.07	18.8	18.69	17.38	16.64	15.54	12.57	12.12	11.44	11.2	10.9	11.33	11.60
20	12.00	11.26	11.12	10.95	11.56	12.90	13.37	15.52	15.46	17.62	18.05	18.68	17.18	16.61	15.52	12.57	12.11	11.44	11.2	10.9	11.28	11.40
30	11.99	11.26	11.11	10.94	11.52	12.89	12.85	14.52	13.79	13.5	14.8	15.3	16.83	16.56	15.5	12.56	12.11	11.44	11.2	10.8	11.02	11.30
40	11.98	11.26	11.11	10.94	11.04	12.00	11.87	13.01	12.41	12.43	13.1	12.42	12.9	13.35	15.39	12.56	12.11	11.44	11.2	10.9	10.97	11.30
50	11.98	11.26	11.11	10.94	10.96	11.50	11.57	11.80	11.70	11.61	12.06	11.73	12.09	11.93	11.92	12.56	12.11	11.44	11.2	10.9	10.96	11.20
60	11.95	11.26	11.10	10.94	10.92	11.13	11.24	11.27	11.32	11.38	11.52	11.43	11.51	11.53	11.49	12.53	12.1	11.44	11.2	10.8	10.94	11.20
70	11.76	11.26	11.09	10.94	10.91	11.01	11.13	11.13	11.22	11.24	11.25	11.27	11.3	11.3	11.33	11.98	12.1	11.44	11.2	10.8	10.93	11.20
80	11.51	11.26	11.08	10.92	10.90	10.96	11.03	11.05	11.16	11.16	11.17	11.2	11.24	11.25	11.27	11.35	11.97	11.44	11.2	10.9	10.92	11.10
90	11.45	11.26	11.08	10.91	10.90	10.95	11.01	11.02	11.12	11.13	11.15	11.17	11.19	11.22	11.28	11.27	11.49	11.43	11.2	10.9	10.91	11.10
100	11.41	11.26	11.08	10.91	10.90	10.94	10.99	11.00	11.08	11.12	11.14	11.16	11.17	11.38	11.25	11.39	11.41	11.2	10.9	10.9	11.10	
110	11.39	11.26	11.08	10.91	10.90	10.92	10.97	10.99	11.07	11.1	11.13	11.13	11.14	11.18	11.27	11.24	11.32	11.37	11.2	10.9	10.89	11.00
120	11.36	11.26	11.08	10.91	10.89	10.92	10.95	10.97	11.04	11.1	11.12	11.13	11.14	11.17	11.26	11.21	11.29	11.32	11.2	10.9	10.87	11.00
130	11.35	11.26	11.07	10.90	10.89	10.91	10.94	10.96	11.04	11.09	11.1	11.13	11.13	11.15	11.24	11.2	11.25	11.27	11.2	10.9	10.85	10.90
140	11.34	11.26	11.07	10.90	10.89	10.90	10.94	10.96	11.04	11.08	11.1	11.13	11.13	11.14	11.23	11.19	11.23	11.26	11.2	10.8	10.83	10.90
150	11.33	11.26	11.07	10.90	10.89	10.90	10.94	10.96	11.03	11.08	11.1	11.12	11.13	11.14	11.19	11.9	11.23	11.26	11.2	10.8	10.81	10.90

Dissolved Oxygen (g m⁻³)

Depth (m)																						
0	9.2	10.2	9.6	10.6	10.4	9.9	9.5	9.4	9.1	9.1	9.0	8.7	8.8	9.4	10.5	10.2	10.3	10.4	9.7	10.5	10.5	10.3
10	9.1	10.5	9.6	10.7	10.4	9.9	9.8	9.5	8.9	9.0	8.9	8.7	8.9	9.3	9.5	10.2	10.3	10.7	9.5	10.4	10.7	10.3
20	9.4	9.4	9.6	10.6	10.4	10.0	9.4	9.5	9.0	9.0	9.1	8.7	8.8	9.3	9.5	10.2	10.3	10.7	9.4	10.3	10.6	10.2
30	9.8	9.2	9.6	10.6	10.4	10.1	9.4	9.1	8.8	9.0	9.1	8.4	8.7	9.2	9.4	10.2	10.2	10.7	9.4	10.3	10.5	10.2
40	9.8	9.1	9.6	10.6	10.0	9.7	8.9	9.1	8.6	8.8	9.0	8.4	8.3	8.7	9.3	10.1	10.2	10.6	9.4	10.2	10.4	10.2
50	9.6	8.9	9.6	10.6	9.9	9.5	9.0	8.7	8.6	8.7	8.7	8.2	8.2	8.3	8.6	10.1	10.2	10.6	9.4	10.2	10.3	10.1
60	9.4	8.9	9.5	10.5	9.8	9.3	8.7	8.6	8.5	8.6	8.6	8.2	8.1	8.1	8.3	10.0	10.1	10.5	9.4	10.2	10.2	10.1
70	9.5	9.0	9.4	10.4	9.7	9.3	8.8	8.7	8.5	8.6	8.5	8.2	8.0	8.0	8.2	9.6	10.1	10.5	9.3	10.1	10.2	10.0
80	7.7	8.9	9.4	10.4	9.7	9.2	8.6	8.4	8.5	8.6	8.4	8.1	7.9	7.9	8.2	8.5	10.0	10.3	9.4	10.1	10.2	10.1
90	7.8	8.9	9.4	10.4	9.6	9.5	8.8	8.5	8.5	8.6	8.2	8.1	7.8	7.8	8.0	8.3	9.7	10.3	9.4	10.1	10.1	10.1
100	7.5	8.6	9.3	10.4	9.6	9.2	8.6	8.4	8.3	8.5	8.1	8.0	7.8	7.8	7.5	8.2	8.6	10.1	9.4	10.1	10.0	9.8
110	7.4	8.7	9.3	10.4	9.6	9.2	8.6	8.4	8.3	8.4	8.1	8.0	7.7	7.7	7.3	8.1	8.3	9.8	9.3	9.9	9.9	9.8
120	6.9	8.5	9.3	10.3	9.5	9.0	8.4	8.4	8.3	8.2	8.1	7.9	7.7	7.6	7.2	8.0	8.1	8.8	9.3	9.9	9.9	9.8
130	6.9	8.5	9.3	10.2	9.5	9.0	8.4	8.4	8.3	8.2	8.2	7.9	7.6	7.5	7.3	7.9	8.0	8.5	9.3	9.9	9.9	9.7
140	6.8	8.3	9.2	10.2	9.5	8.6	8.2	8.2	8.1	8.0	8.1	7.8	7.1	7.8	7.3	7.8	7.8	8.1	9.3	9.9	9.9	9.4
150	6.4	8.2	9.2	10.2	9.3	8.5	8.1	8.1	7.9	7.8	7.9	7.6	7.0	7.2	7.3	7.7	7.8	8.1	9.3	9.8	9.8	9.4

Secchi depth

(m)	12	14.5	13.5	17.5	11	14.5	15.5	16	13	13	15	14.5	19	22	16.4	17	16	15.5	12	9.5	12	15.5
-----	----	------	------	------	----	------	------	----	----	----	----	------	----	----	------	----	----	------	----	-----	----	------

**Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Additional site B (Kuratau Basin) for the period starting 8 January 2002 on**

2001-2002

Temperature	8/01/2002	22/01/2002	6/03/2002	4/04/2002	22/04/2002	5/05/2002	19/06/2002	1/07/2002	17/07/2002	31/07/2002	29/08/2002	18/09/2002
Date												
Depth (m)												
0	18.1	18.8	18.64	17.38	16.84	15.12	12.45	12.13	11.48	11.3	11	11.08
10	17.55	18.45	18.58	17.35	16.61	15.14	12.44	12.09	11.49	11.1	10.8	11.05
20	15.72	17.4	18.56	17.1	16.6	15.05	12.44	12.09	11.48	11.1	10.8	11.03
30	13.74	13.9	15.07	16.74	16.4	14.75	12.43	12.09	11.48	11.1	10.8	11.03
40	12.62	12.73	13.08	14.3	13.4	14.4	12.24	12.08	11.48	11.1	10.8	11.02
50	11.92	11.98	11.91	12.77	12.12	14.07	12.11	11.97	11.49	11.1	10.8	10.91
60	11.31	11.41	11.5	12.03	11.53	12.96	11.73	11.93	11.49	11.1	10.8	10.9
70	11.21	11.25	11.24	11.5	11.32	12.2	11.49	11.87	11.48	11.1	10.8	10.89
80	11.15	11.19	11.21	11.29	11.24	11.97	11.38	11.78	11.48	11.1	10.8	10.89
90	11.1	11.13	11.15	11.2	11.18	11.69	11.3	11.37	11.46	11.1	10.7	10.87
100	11.1	11.12	11.12	11.19	11.15	11.39	11.22	11.28	11.3	11	10.7	10.85
110										10.7	10.7	
Dissolved Oxygen (g m⁻³)												
Depth (m)												
0	8.7	8.8	9.3	9.3	9.3	10.9	10.4	10.3	10.4	9.9	10.4	10.4
10	8.6	9	9.1	9.2	9.3	9.5	10.3	10.3	10.8	9.7	10.3	10.5
20	8.8	9	9.1	9.2	9.2	9.4	10.2	10.2	10.6	9.6	10.3	10.5
30	8.8	8.9	8.6	9.1	9.2	9.3	10.2	10.2	10.6	9.6	10.2	10.5
40	8.7	8.7	8.7	8.9	8.5	9.1	10.1	10.1	10.5	9.6	10.2	10.4
50	8.7	8.4	8.5	8.6	8.2	9	10	10.1	10.5	9.6	10.1	10.3
60	8.7	8.3	8.4	8.4	8	8.6	9	9.8	10.4	9.6	10.1	10.2
70	8.7	8.3	8.3	8.3	7.9	8.1	8.7	9.7	10.4	9.5	10	10.1
80	8.7	8.2	8.1	8.1	7.8	7.9	8.4	9.5	10.3	9.5	10	10.1
90	8.2	8.1	7.9	7.7	7.7	7.8	8.2	9.1	10.3	9.5	10	10
100	8	7.6	7.5	7.7	7.5	7.7	7.8	8.7	9.8	9.6	9.9	9.9
110	8				6.2					9.2	9.8	
Secchi depth												
Depth (m)	13.5	12	14.5	19.5	19	13.2	15	16	12.5	10.5	8	11

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Additional site C (Western Bays) for the period starting 8 January 2002 on

2001-2002

Temperature	8/01/2002	22/01/2002	6/03/2002	4/04/2002	22/04/2002	5/05/2002	19/06/2002	1/07/2002	17/07/2002	31/07/2002	29/08/2002	18/09/2002	9/10/2002
Date													
Depth (m)													
0	18.72	18.82	18.68	17.47	16.88	15.6	12.58	12.22	11.52	11.6	11.4	11.24	12.10
10	17.41	18.46	18.47	17.24	11.63	15.64	12.56	12.15	11.5	11.2	10.9	11.23	11.30
20	16.95	18.21	18.32	17.16	16.58	15.64	12.56	12.14	11.49	11.2	10.9	11.16	11.30
30	14	13.77	15.9	17.12	16.5	15.61	12.56	12.14	11.49	11.2	10.8	11.06	11.20
40	13.14	12.01	12.98	13.17	13.02	12.26	12.56	12.13	11.49	11.2	10.8	11.02	11.20
50	11.97	11.5	12.13	12.11	11.87	11.57	12.56	12.13	11.49	11.2	10.8	11.02	11.20
60	11.44	11.26	11.59	11.57	11.47	11.37	11.9	11.92	11.49	11.2	10.8	11	11.10
70	11.26	11.17	11.36	11.38	11.32	11.29	11.36	11.55	11.49	11.2	10.8	10.99	11.10
80	11.18	11.16	11.25	11.32	11.26	11.24	11.28	11.5	11.49	11.2	10.8	10.95	11.10
90	11.15	11.14	11.18	11.21	11.23	11.21	11.23	11.47	11.49	11.2	10.8	10.94	11.00
100	11.12	11.11	11.18	11.19	11.19	11.19	11.22	11.45	11.49	11.2	10.8	10.92	11.00
110	11.11	11.1			11.16	11.15				11.2	10.8		10.90
120										11.2	10.8		10.90
Dissolved Oxygen (g m⁻³)													
Depth (m)													
0	8.6	8.9	9.3	9.4	9.3	10.6	10.3	10.4	10.5	9.7	10.3	10.5	10.4
10	8.4	8.9	9	9.1	9.2	9.5	10.2	10.4	10.8	9.5	10.2	10.7	10.4
20	8.9	8.9	9	9.1	9.2	9.5	10.2	10.4	10.8	9.5	10.2	10.7	10.4
30	8.6	8.9	8.8	9.1	9.1	9.4	10.1	10.3	10.7	9.4	10.1	10.6	10.4
40	8.6	8.5	8.6	8.6	8.5	8.9	10.1	10.3	10.5	9.4	10	10.5	10.3
50	8.5	8.2	8.5	8.5	8.1	8.6	10	10.2	10.5	9.4	10	10.4	10
60	8.6	8.1	8.5	8.2	7.9	8.3	9.7	10	10.5	9.4	10	10.4	10
70	8.6	8.1	8.2	8.2	7.8	8.2	9.1	9.6	10.5	9.4	9.9	10.3	9.9
80	8.7	8.1	8.1	8	7.7	8	8.4	8.8	10.5	9.3	9.9	10.2	9.9
90	8.6	8.1	8.1	7.9	7.7	7.9	8	8.7	10.4	9.3	9.9	10.1	9.8
100	8.7	8.1	8.1	7.9	7.6	7.8	7.7	8.6	10.2	9.3	10	10	9.6
110	8.5	7.9			7.6	7.7				9.3	10		9.7
120	8.5	7.7								9.1	9.9		9.6
Secchi depth													
Depth (m)	14.5	15.5	16	19	18.5	15.6	16	14	12.5	12	8	12	19

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
 For the period starting 11 July 2000

2000-2001

Temperature		2000-2001																		
Date	11-7-00	04-8-00	21-8-00	11-9-00	28-9-00	25-10-00	13-11-00	06-12-00	03-1-01	15-1-01	20-2-01	01-3-01	19-3-01	09-4-01	11-4-01	10-5-01	29-5-01	02-7-01	25-7-01	13-8-01
Depth (m)																				
0	11.87	11.32	11.19	11.80	12.47	14.04	13.27	15.73	18.16	18.98	20.47	20.87	19.01	16.99	16.99	15.78	13.62	12.11	11.26	11.15
10	11.87	11.32	11.15	11.46	11.52	13.03	13.09	15.06	17.37	18.51	19.37	20.71	19.05	16.87	16.99	15.78	13.74	12.04	11.26	11.12
20	11.86	11.32	11.14	11.33	11.36	11.99	12.98	14.15	15.46	14.79	18.08	18.98	19.06	16.78	16.97	15.78	13.78	12.00	11.26	11.12
30	11.86	11.33	11.14	11.30	11.33	11.83	12.80	13.31	13.61	13.63	16.06	15.95	16.46	15.82	16.84	15.73	13.79	11.99	11.26	11.11
40	11.86	11.33	11.14	11.27	11.31	11.60	12.36	12.49	12.73	12.81	13.39	13.36	13.05	13.13	13.87	13.19	13.80	11.98	11.26	11.11
50	11.86	11.33	11.14	11.22	11.30	11.49	12.10	12.16	12.21	12.27	12.67	12.58	12.42	12.35	12.68	12.42	13.80	11.98	11.26	11.11
60	11.64	11.33	11.15	11.18	11.27	11.42	11.69	11.78	11.76	11.87	12.01	12.01	11.84	11.89	11.90	11.92	11.95	11.26	11.10	
70	11.42	11.33	11.15	11.15	11.24	11.39	11.41	11.53	11.64	11.67	11.77	11.79	11.67	11.67	11.69	11.61	11.76	11.26	11.09	
80	11.31	11.33	11.15	11.14	11.20	11.38	11.29	11.40	11.47	11.55	11.56	11.63	11.55	11.54	11.54	11.52	11.54	11.51	11.26	11.08
90	11.22	11.33	11.15	11.13	11.17	11.33	11.26	11.36	11.43	11.46	11.50	11.55	11.49	11.46	11.48	11.47	11.46	11.45	11.26	11.08
100	11.21	11.32	11.15	11.13	11.14	11.33	11.21	11.32	11.38	11.39	11.43	11.50	11.43	11.41	11.43	11.42	11.42	11.41	11.26	11.08
110	11.19	11.32	11.15	11.13	11.06	11.29	11.19	11.28	11.36	11.40	11.46	11.41	11.37	11.39	11.40	11.38	11.39	11.38	11.26	11.08
120	11.19	11.31	11.15	11.13	11.04	11.27	11.19	11.27	11.33	11.34	11.39	11.44	11.39	11.33	11.35	11.38	11.35	11.36	11.26	11.08
130	11.18	11.26	11.15	11.12	11.02	11.23	11.17	11.26	11.30	11.32	11.37	11.43	11.37	11.32	11.34	11.36	11.33	11.35	11.26	11.07
140	11.16	11.18	11.14	11.12	11.01	11.18	11.15	11.25	11.30	11.31	11.35	11.40	11.35	11.31	11.32	11.34	11.31	11.34	11.26	11.07
150	11.15	11.18	11.14	11.12	11.01	11.15	11.15	11.25	11.32	11.31	11.33	11.41	11.34	11.31	11.32	11.34	11.31	11.33	11.26	11.07
Dissolved Oxygen (g m ⁻³)		2000-2001																		
Depth (m)																				
0	9.0	9.0	9.2	9.3	9.1	8.9	8.2	8.7	8.2	8.0	8.0	8.2	8.4	8.3	8.4	8.2	8.7	9.2	10.2	9.6
10	9.0	9.0	9.4	9.5	8.7	8.8	8.4	8.3	8.3	8.6	8.0	8.5	8.3	8.3	8.2	8.0	8.5	9.1	10.5	9.6
20	9.0	9.1	9.4	9.5	8.7	9.1	8.4	8.5	8.4	8.1	8.2	8.6	8.6	8.4	7.9	7.9	8.4	9.4	9.6	
30	9.0	9.1	9.6	9.5	8.7	8.9	8.4	8.5	8.5	8.2	8.0	8.3	8.0	8.0	8.0	7.8	8.4	9.8	9.2	9.6
40	9.0	9.1	9.6	9.5	9.1	8.7	8.2	8.2	8.4	7.9	8.1	8.1	7.6	7.8	7.6	7.7	8.3	9.8	9.1	9.6
50	9.0	9.1	9.6	9.5	9.1	8.5	8.2	8.2	8.2	8.1	7.9	7.8	7.6	7.5	7.4	7.5	8.3	9.6	8.9	9.6
60	9.0	9.1	9.7	9.5	8.7	8.4	8.0	7.9	8.0	7.5	7.7	7.4	6.8	7.2	7.2	7.5	7.2	9.4	8.9	9.5
70	8.9	9.1	9.7	9.5	8.7	8.3	7.9	7.8	7.9	7.4	7.6	7.2	6.8	7.1	7.4	7.3	7.0	9.5	9.0	9.4
80	7.8	9.0	9.7	9.5	8.7	8.2	7.6	7.6	7.8	7.5	7.4	7.0	6.5	6.9	7.3	7.3	7.0	7.7	8.9	9.4
90	7.4	8.9	9.7	9.5	8.7	8.2	7.6	7.6	7.7	7.5	7.4	6.9	6.5	6.9	7.1	7.1	7.0	7.8	8.9	9.4
100	7.2	8.7	9.7	9.5	8.7	8.0	7.5	7.6	7.6	7.3	7.2	6.8	6.6	6.8	7.0	7.0	6.9	7.5	8.6	9.3
110	7.1	8.3	9.7	9.5	8.7	8.0	7.5	7.5	7.6	7.2	7.1	6.7	6.5	6.8	7.0	7.0	6.7	7.4	8.7	9.3
120	6.9	7.9	9.7	9.5	8.2	8.1	7.4	7.4	7.5	7.1	7.0	6.5	6.5	6.7	6.8	6.9	6.6	8.5	9.3	
130	6.9	7.3	9.7	9.5	8.5	8.1	7.4	7.3	7.4	7.0	7.0	6.5	6.5	6.6	6.7	6.6	6.5	6.9	8.5	9.3
140	6.9	7.1	9.7	9.5	8.6	8.0	7.3	7.2	7.2	6.9	6.8	6.4	6.5	6.4	6.4	6.7	6.3	6.8	8.3	9.2
150	6.8	7.4	9.7	9.3	8.5	7.9	7.3	7.1	7.1	6.6	6.5	6.3	6.4	6.3	6.3	6.6	6.1	6.4	8.2	9.2
Secchi depth		2000-2001																		
Depth (m)																				
11	12	15	12	13	11	12	17	17	18	17	14.5	17	13.5	13.5	17	14.5	12	14.5	13.5	

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
 For the period starting 13 July 1999

1999-2000

Temperature

Date	13-7-99	6-8-99	3-9-99	29-9-99	18-10-99	19-12-99	18-1-00	12-4-00	4-5-00	25-5-00	20-6-00	11-7-00	4-8-00	21-8-00	11-9-00	28-9-00	25-10-00	13-11-00	6-12-00	
Depth (m)																				
0	12.0	11.8	11.8	11.5	12.8	16.56	18.63	17.41	15.82	14.22	12.28	11.87	11.32	11.19	11.80	12.47	14.04	13.27	15.73	
10	12.0	11.4	11.3	11.5	12.7	16.40	18.35	17.25	15.77	14.28	12.28	11.87	11.32	11.15	11.46	11.52	13.03	13.09	15.06	
20	12.0	11.4	11.2	11.5	12.4	15.96	17.22	17.21	15.76	14.31	12.28	11.86	11.32	11.14	11.33	11.36	11.99	12.98	14.15	
30	12.0	11.4	11.1	11.4	11.6	15.23	14.94	16.65	15.75	14.28	12.27	11.86	11.33	11.14	11.30	11.33	11.83	12.86	13.31	
40	12.0	11.3	11.1	11.2	11.4	12.16	13.29	12.55	13.64	14.22	12.26	11.86	11.33	11.14	11.27	11.31	11.60	12.36	12.49	
50	12.0	11.3	11.1	11.1	11.3	11.64	11.91	11.67	12.14	12.53	12.26	11.86	11.33	11.14	11.22	11.30	11.49	12.10	12.16	
60	12.0	11.3	11.0	11.1	11.1	11.35	11.45	11.39	11.56	11.56	12.21	11.85	11.33	11.15	11.18	11.27	11.42	11.69	11.78	
70	12.0	11.3	11.0	11.0	11.1	11.25	11.31	11.29	11.36	11.34	11.58	11.64	11.33	11.15	11.15	11.24	11.39	11.41	11.53	
80	11.4	11.3	11.0	11.0	11.0	11.18	11.21	11.23	11.24	11.23	11.32	11.42	11.33	11.15	11.14	11.20	11.38	11.29	11.40	
90	11.3	11.3	11.0	11.0	11.0	11.16	11.17	11.20	11.21	11.20	11.24	11.31	11.33	11.15	11.13	11.17	11.33	11.26	11.36	
100	11.2	11.2	11.0	11.0	11.0	11.14	11.14	11.17	11.17	11.17	11.15	11.17	11.22	11.32	11.15	11.13	11.14	11.33	11.21	11.32
110	11.2	11.2	11.0	11.0	11.0	11.12	11.12	11.15	11.14	11.12	11.16	11.21	11.32	11.15	11.13	11.06	11.29	11.19	11.28	
120	11.2	11.1	11.0	11.0	11.0	11.10	11.09	11.13	11.12	11.10	11.14	11.19	11.31	11.15	11.13	11.04	11.27	11.19	11.27	
130	11.1	11.1	11.0	11.0	11.0	11.08	11.08	11.11	11.10	11.09	11.12	11.18	11.26	11.15	11.12	11.02	11.23	11.17	11.26	
140	11.1	11.1	11.0	11.0	11.0	11.07	11.07	11.09	11.09	11.09	11.10	11.16	11.18	11.14	11.12	11.01	11.18	11.15	11.25	
150	11.1	11.0	11.0	10.9	11.0	11.10	11.06	11.09	11.09	11.07	11.10	11.15	11.18	11.14	11.12	11.01	11.15	11.15	11.25	

Dissolved Oxygen (g m⁻³)

Depth (m)	0	10.5	10.1	9.2	9.5	8.9	8.3	7.9	9.2	8.7	8.5	8.1	9.0	9.0	9.2	9.3	9.1	8.9	8.2	8.7
10	10.7	10.2	9.8	9.8	8.9	8.6	7.9	9.2	8.6	8.3	8.3	9.0	9.0	9.4	9.5	8.7	8.8	8.4	8.3	
20	10.7	9.9	9.8	9.9	8.9	8.7	8.1	9.2	8.8	8.5	8.7	9.0	9.1	9.4	9.5	8.7	9.1	8.4	8.5	
30	10.6	10.0	9.8	9.7	8.9	8.7	8.3	9.0	8.8	8.5	8.6	9.0	9.1	9.6	9.5	8.7	8.9	8.4	8.5	
40	10.6	9.7	9.5	9.6	8.8	8.7	8.1	8.3	8.2	8.6	8.6	9.0	9.1	9.6	9.5	9.1	8.7	8.2	8.2	
50	10.4	9.9	9.5	9.3	8.6	8.7	8.0	8.0	7.9	8.2	8.6	9.0	9.1	9.6	9.5	9.1	8.5	8.2	8.2	
60	10.4	9.8	9.4	9.2	8.6	8.6	8.0	8.0	7.9	7.7	8.7	9.0	9.1	9.7	9.5	8.7	8.4	8.0	7.9	
70	10.3	9.7	9.3	9.0	8.6	8.6	8.7	8.0	8.0	7.8	7.7	8.4	8.9	9.1	9.7	9.5	8.7	8.3	7.9	
80	10.3	9.0	9.2	9.0	8.5	8.5	7.9	7.9	7.7	7.6	7.6	7.8	9.0	9.7	9.5	8.7	8.2	7.6	7.6	
90	8.1	8.6	9.2	9.0	8.6	8.5	7.7	7.9	7.8	7.4	7.4	7.4	8.9	9.7	9.5	8.7	8.2	7.6	7.6	
100	7.9	7.3	9.2	8.9	8.6	8.5	8.3	7.7	7.6	7.4	7.3	7.2	8.7	9.7	9.5	8.7	8.0	7.5	7.6	
110	7.5	7.1	9.1	8.9	8.6	8.3	8.1	7.7	7.6	7.4	7.4	7.1	8.3	9.7	9.5	8.7	8.0	7.5	7.5	
120	7.4	6.8	9.1	8.9	8.3	8.4	8.1	7.7	7.4	7.5	7.3	6.9	7.9	9.7	9.5	8.2	8.1	7.4	7.4	
130	7.3	6.7	9.0	8.8	7.9	8.2	8.0	7.5	7.4	7.5	7.3	6.9	7.3	9.7	9.5	8.5	8.1	7.4	7.3	
140	7.1	6.7	8.9	8.7	7.5	8.1	8.0	7.5	7.2	7.4	7.2	6.9	7.1	9.7	9.5	8.6	8.0	7.3	7.2	
150	6.9	6.4	8.9	8.6	7.5	8.0	7.5	7.2	6.8	7.0	6.9	6.8	7.4	9.7	9.3	8.5	7.9	7.3	7.1	

Secchi depth

Depth (m)	16	14.5	10	10	14.9	18	19.1	15	14	14	14	11	12	15	12	13	11	12	17
-----------	----	------	----	----	------	----	------	----	----	----	----	----	----	----	----	----	----	----	----

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.

For the period starting 28 July 1998

Temperature

Date	28-7-98	22-8-98	29-9-98	1-11-98	26-11-98	22-12-98	12-2-99	3-3-99	14-4-99	30-4-99	19-5-99	1-6-99	17-6-99	13-7-99	6-8-99	3-9-99	29-9-99	18-10-99
Depth (m)																		
0	11.4	11.5	12.9	13.6	18.4	18.5	20.1	20.9	18.3	16.4	14.4	14.2	13.0	12.0	11.8	11.8	11.5	12.8
10	11.6	11.3	11.9	13.2	15.6	16.7	20.1	19.8	18.3	16.4	14.4	14.1	13.4	12.0	11.4	11.3	11.5	12.7
20	11.6	11.3	11.5	12.7	15.4	15.7	20.1	19.8	18.3	16.4	14.5	14.1	13.4	12.0	11.4	11.2	11.5	12.4
30	11.6	11.3	11.3	12.4	12.7	14.5	14.9	15.1	18.1	16.0	14.5	14.1	13.4	12.0	11.4	11.1	11.4	11.6
40	11.6	11.3	11.2	12.4	12.1	12.7	13.2	13.1	12.9	13.1	14.5	13.9	13.4	12.0	11.3	11.1	11.2	11.4
50	11.6	11.3	11.1	12.2	11.8	11.8	12.1	12.1	11.9	12.2	13.1	13.0	13.4	12.0	11.3	11.1	11.1	11.3
60	11.6	11.3	11.1	11.7	11.5	11.5	11.6	11.8	11.6	12.0	11.8	12.0	12.1	12.0	11.3	11.0	11.1	11.1
70	11.6	11.1	11.0	11.2	11.3	11.3	11.4	11.5	11.4	11.8	11.3	11.4	11.5	12.0	11.3	11.0	11.0	11.1
80	10.6	10.9	11.0	11.1	11.2	11.2	11.2	11.4	11.3	11.2	11.2	11.3	11.3	11.4	11.3	11.0	11.0	11.0
90	10.6	10.9	10.9	11.1	11.1	11.1	11.1	11.3	11.2	11.1	11.1	11.2	11.2	11.3	11.3	11.0	11.0	11.0
100	10.5	10.8	10.9	11.0	11.1	11.1	11.1	11.3	11.2	11.1	11.1	11.1	11.2	11.2	11.2	11.0	11.0	11.0
110	10.5	10.5	10.9	11.0	11.0	11.1	11.1	11.2	11.2	11.1	11.1	11.1	11.1	11.2	11.2	11.0	11.0	11.0
120	10.5	10.5	10.9	11.0	11.0	11.0	11.0	11.2	11.2	11.1	11.1	11.1	11.1	11.2	11.1	11.0	11.0	11.0
130	10.5	10.5	10.7	11.0	11.0	11.1	11.1	11.1	11.1	11.1	11.0	11.1	11.1	11.1	11.1	11.0	11.0	11.0
140	10.5	10.5	10.7	10.9	11.0	11.1	11.1	11.1	11.1	11.1	11.0	11.1	11.0	11.1	11.1	11.0	11.0	11.0
150	10.5	10.5	10.7	10.9	11.0	11.1	11.1	11.1	11.1	11.1	11.0	11.1	11.0	11.1	11.0	11.0	10.9	11.0

Dissolved Oxygen (g m⁻³)

Depth (m)	0	10.6	10.6	10.6	10.4	9.6	9.7	9.0	8.6	9.1	9.5	9.9	10.0	10.4	10.5	10.1	9.2	9.5	8.9
10	10.5	10.5	10.7	10.7	9.9	10.1	9.0	8.7	9.2	9.5	10.5	10.4	10.4	10.3	10.7	10.2	9.8	9.8	8.9
20	10.4	10.4	10.6	10.7	9.8	10.2	8.9	8.7	9.1	9.6	10.4	10.4	10.4	10.4	10.7	9.9	9.8	9.9	8.9
30	10.4	10.3	10.5	10.6	10.1	10.2	9.9	9.5	9.1	9.6	10.1	10.7	10.5	10.6	10.0	9.8	9.7	8.9	8.9
40	10.3	10.3	10.3	10.4	10.0	10.1	9.9	9.2	9.1	9.1	10.0	10.4	10.4	10.4	10.6	9.7	9.5	9.6	8.8
50	10.3	10.2	10.2	10.2	9.8	9.9	9.6	8.9	9.0	8.7	9.2	9.6	10.4	10.4	10.4	9.9	9.5	9.3	8.6
60	10.3	10.1	10.1	10.0	9.7	9.7	9.5	8.8	8.9	8.7	8.7	9.4	9.0	10.4	9.8	9.4	9.2	8.6	8.6
70	10.3	9.5	9.9	9.6	9.5	9.5	9.4	8.7	8.7	8.6	8.3	9.1	8.9	10.3	9.7	9.3	9.0	8.6	8.6
80	8.6	8.2	9.5	9.1	9.2	9.3	9.2	8.6	8.6	8.4	8.2	9.1	8.6	10.3	9.0	9.2	9.0	8.5	8.5
90	8.5	7.9	9.3	8.8	9.1	9.1	9.1	8.4	8.6	8.0	7.8	8.8	8.5	8.1	8.6	9.2	9.0	8.6	8.6
100	8.3	7.4	8.9	8.5	9.1	8.9	8.9	8.3	8.6	8.6	8.0	7.7	8.5	8.2	7.9	7.3	9.2	8.9	8.6
110	8.3	7.4	8.5	8.3	8.8	8.9	8.7	8.2	8.5	8.0	7.5	8.2	8.1	7.5	7.1	9.1	8.9	8.6	8.6
120	8.2	7.4	7.7	8.0	8.6	8.8	8.3	7.9	8.3	7.9	7.4	8.2	8.0	7.4	6.8	9.1	8.9	8.3	8.3
130	8.2	7.4	7.6	7.8	8.4	8.6	8.1	7.7	8.1	7.7	7.3	8.1	7.7	7.3	6.7	9.0	8.8	7.9	7.9
140	8.1	7.4	7.4	7.6	8.2	8.4	7.9	7.5	7.9	7.5	7.2	7.8	7.4	7.1	6.7	8.9	8.7	7.5	7.5
150	8.1	7.4	7.4	7.6	8.0	8.2	7.7	7.3	7.7	7.3	7.0	7.5	7.3	6.9	6.4	8.9	8.6	7.5	7.5

Secchi depth

Depth (m)	10.0	10.5	10.4	13.5	15.0	14.5	12.5	14.3	13.0	12.2	15.0	15.0	15.0	16.0	14.5	10.0	10.0	14.9
-----------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
 For the period starting 16 September 1997

1997-1998

Temperature

Date	16-9-97	11-10-97	28-10-97	02-12-97	21-1-98	04-3-98	24-3-98	26-3-98	07-4-98	29-5-98	28-7-98	22-8-98
Depth (m)												
1	10.8	11.8	12.2	14.5	17.7	20.0	19.3	18.6	17.7	14.2	11.4	11.49
10	10.5	11.4	12.0	13.7	17.6	19.9	18.6	18.6	17.7	14.3	11.6	11.32
20	10.5	11.1	11.5	13.6	16.5	19.7	18.5	18.5	17.7	14.0	11.6	11.27
30	10.5	10.8	11.5	13.1	14.3	16.4	18.0	18.1	17.5	13.1	11.6	11.27
40	10.5	10.6	11.4	12.5	12.0	13.3	13.0	12.6	13.7	12.0	11.6	11.27
50	10.5	10.5	11.1	11.5	11.2	12.0	11.9	11.7	11.5	11.2	11.6	11.26
60	10.5	10.5	11.1	11.0	11.0	11.5	11.1	11.1	11.0	10.9	11.6	11.26
70	10.5	10.5	10.8	10.8	10.8	11.0	10.7	10.8	10.8	10.8	11.6	11.12
80	10.5	10.5	10.7	10.7	10.7	10.8	10.6	10.7	10.6	10.6	10.6	10.90
90	10.5	10.5	10.6	10.6	10.6	10.7	10.5	10.6	10.6	10.6	10.6	10.86
100	10.5	10.5	10.5	10.5	10.6	10.7	10.5	10.6	10.6	10.6	10.5	10.82
110	10.5	10.5	10.4	10.5	10.6	10.6	10.5	10.5	10.5	10.6	10.5	10.5
120	10.5	10.5	10.5	10.5	10.5	10.6	10.5	10.5	10.5	10.5	10.5	10.5
130	10.5	10.5	10.5	10.5	10.5	10.6	10.5	10.5	10.5	10.5	10.5	10.5
140	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5
150	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5	10.5

Dissolved Oxygen (g m⁻³)

Depth (m)	1	10.37	10.68	9.89	9.27	9.17	9.43	9.10	9.14	9.92	10.60	10.64
1	10.55	10.37	10.68	9.89	9.27	9.17	9.43	9.10	9.14	9.92	10.60	10.64
10	10.52	10.51	10.22	9.86	9.38	9.19	9.53	9.07	9.10	9.88	10.46	10.50
20	10.50	10.46	10.24	9.86	9.46	9.22	9.61	8.95	9.07	9.87	10.40	10.36
30	10.29	10.46	10.00	9.74	9.81	9.30	9.78	8.97	9.09	9.68	10.35	10.27
40	10.31	10.39	9.96	9.66	9.85	9.32	9.73	9.47	9.32	9.40	10.32	10.26
50	10.27	10.36	9.89	9.47	9.53	9.16	9.55	9.45	9.34	9.26	10.30	10.20
60	10.16	10.31	9.77	9.44	9.37	9.17	9.30	9.47	9.30	9.18	10.28	10.10
70	10.08	10.24	9.76	9.19	9.30	9.11	9.21	9.38	9.24	9.20	10.25	9.54
80	10.06	10.15	9.85	9.04	9.13	9.04	9.14	9.30	9.13	9.12	8.58	8.15
90	10.03	10.09	9.33	9.00	9.10	8.93	9.03	9.24	9.05	9.08	8.52	7.90
100	9.99	10.06	9.23	8.96	9.01	8.89	8.39	9.16	8.97	8.94	8.34	7.36
110	9.96	10.02	9.03	8.87	8.89	8.83	8.38	8.98	8.94	8.78	8.26	7.36
120	9.91	10.00	8.96	8.87	8.84	8.75	8.38	8.87	8.88	8.69	8.21	7.36
130	9.86	9.92	8.76	8.84	8.68	8.63	8.38	8.38	8.79	8.41	8.21	7.36
140	9.82	9.87	8.76	8.71	8.45	8.30	8.38	8.38	8.58	8.41	8.14	7.36
150	9.56	9.69	8.76	8.65	8.38	8.22	8.38	8.38	8.40	8.41	8.14	7.36

Secchi depth data (m)

Depth (m)	12.0	13.7	12.5	14.5	14.7	11.5	13.5	13.5	13.5	15.5	10.0	10.5

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.

For the period starting 3 September 1996

Temperature

Date	3-9-96	17-9-96	27-9-96	17-10-96	24-10-96	6-11-96	28-11-96	11-12-96	23-12-96	8-1-97	29-1-97	26-3-97	2-4-97	15-4-97	20-5-97	29-5-97	7-7-97	29-7-97
Depth (m)																		
1	10.5	10.7	12.5	13.3	12.6	13.5	13.6	14.8	16.3	17.9	17.8	17.7	17.3	16.7	14.1	14.2	11.7	10.9
10	10.4	10.6	11.6	12.0	12.3	13.6	13.6	14.8	15.3	16.8	17.6	17.6	17.3	16.7	14.0	14.1	11.7	11.0
20	10.3	10.4	11.1	11.9	12.3	13.4	13.3	14.4	15.1	16.5	17.4	17.2	17.2	16.7	14.0	14.1	11.7	11.0
30	10.3	10.3	11.0	11.8	12.3	13.3	13.3	14.2	15.0	15.6	14.8	16.6	17.2	16.7	12.6	14.1	11.7	11.0
40	10.3	10.3	10.5	11.7	11.9	11.7	11.6	12.7	13.5	13.0	13.4	13.8	14.5	14.0	11.5	14.0	11.7	11.0
50	10.4	10.3	10.4	11.5	11.6	10.8	10.9	12.5	12.4	11.9	11.8	12.4	11.5	11.9	11.0	12.1	11.7	11.0
60	10.3	10.3	10.4	10.9	11.1	10.6	10.9	11.7	11.3	11.2	10.9	11.2	10.9	11.1	10.5	11.8	11.7	11.0
70	10.3	10.3	10.3	10.6	10.6	10.5	10.5	11.7	10.7	10.8	10.7	10.7	10.6	10.9	10.5	11.1	11.7	11.0
80	10.3	10.3	10.3	10.5	10.5	10.4	10.4	11.1	10.6	10.6	10.6	10.5	10.5	10.7	10.5	10.8	10.9	11.0
90	10.3	10.3	10.3	10.4	10.4	10.4	10.4	10.4	10.5	10.5	10.4	10.5	10.5	10.6	10.5	10.6	10.8	10.9
100	10.3	10.3	10.3	10.3	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.4	10.5	10.5	10.5	10.5	10.6	10.7
110	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.4	10.4	10.4	10.4	10.4	10.4	10.5	10.5	10.5	10.5	10.6
120	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.4	10.4	10.4	10.4	10.4	10.4	10.5	10.5	10.5	10.5	10.5
130	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.4	10.4	10.4	10.4	10.4	10.4	10.5	10.5	10.5	10.5	10.5
140	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.4	10.3	10.3	10.3	10.4	10.4	10.5	10.5	10.5	10.5	10.5
150	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.3	10.4	10.4	10.5	10.4	10.4	10.5	10.5

Dissolved Oxygen (g m⁻³)

Depth (m)																		
1	8.81	9.08	10.03	9.78	10.32	9.96	9.99	10.03	9.10	8.71	8.80	9.70	9.40	9.06	9.09	9.3	9.9	10.53
10	9.17	9.17	10.43	9.85	10.27	9.84	9.87	9.97	9.30	8.70	8.80	9.30	9.25	8.95	9.10	9.2	9.8	10.42
20	9.14	8.98	10.32	9.84	10.15	9.80	9.80	9.90	9.30	8.70	8.70	8.93	8.94	8.91	9.06	9.2	9.8	10.45
30	8.98	8.95	10.16	9.84	9.89	9.79	9.81	9.76	9.30	8.80	9.10	8.80	8.82	8.87	9.01	9.2	9.8	10.43
40	8.90	8.93	9.98	9.80	9.89	9.73	9.77	9.70	9.30	9.00	8.90	8.78	8.79	8.82	8.94	9.1	9.8	10.46
50	8.78	8.87	9.69	9.76	9.80	9.29	9.35	9.10	9.30	8.80	8.90	8.51	8.58	8.65	8.86	9.1	9.7	10.40
60	8.73	8.80	9.54	9.67	9.67	9.19	9.14	9.04	9.15	8.60	8.70	8.49	8.56	8.71	8.70	9.0	9.7	10.36
70	8.74	8.80	9.45	9.56	9.44	9.14	9.09	9.03	9.07	8.60	8.60	8.47	8.52	8.71	8.64	8.9	9.7	10.34
80	8.70	8.77	9.37	9.42	9.33	9.03	9.01	9.01	9.00	8.60	8.50	8.36	8.46	8.69	8.48	8.5	8.6	10.34
90	8.63	8.70	9.24	9.29	9.30	8.99	8.96	8.92	8.98	8.60	8.50	8.30	8.45	8.63	8.32	8.3	8.2	10.24
100	8.59	8.61	9.11	9.22	9.21	8.94	8.93	8.88	8.95	8.60	8.40	8.27	8.40	8.54	8.29	8.2	8.1	8.70
110	8.48	8.49	9.13	9.15	9.20	8.90	8.87	8.80	8.89	8.50	8.30	8.18	8.29	8.48	8.27	8.1	8.0	8.02
120	8.44	8.33	9.07	8.91	8.98	8.77	8.74	8.73	8.85	8.40	8.20	8.08	8.20	8.41	8.22	8.1	8.0	8.05
130	8.19	8.27	9.07	8.83	8.98	8.71	8.69	8.66	8.30	8.30	7.96	8.02	8.20	8.19	8.1	7.9	8.09	
140	8.39	8.35	9.05	8.89	8.89	8.62	8.65	8.60	8.33	8.20	8.20	7.40	7.60	7.87	7.97	7.8	7.4	7.79
150	8.81	8.84	8.98	8.49	8.94	8.48	8.43	8.47	8.25	8.10	8.10	7.40	7.50	7.71	7.88	7.7	7.2	7.13

Secchi depth data (m)

Secchi d	13.1	14.2	11.2	12.6	13.4	14.9	14.1	14.7	17.7	15.1	15.2	15.3	16.0	17.7	14.6	14.5	12.5	13.5
----------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.

For the period starting 12 September 1995

Temperature

Date	12-9-95	25-9-95	30-10-95	24-11-95	06-12-95	12-1-96	31-1-96	13-2-96	29-2-96	20-3-96	28-3-96	18-4-96	19-5-96	14-6-96	9-7-96
Depth (m)															
1	10.7		13.7		17.7	21.1	21.7	22.7	20.5	18.2	16.8	17.7	14.8	12.2	11.2
10	10.7		11.9		16.2	20.7	20.7	21.0	20.1	18.2	16.7	17.4	14.8	12.2	11.2
20	10.7		11.4		15.3	18.1	18.5	20.6	20.0	18.2	16.6	17.3	14.8	12.1	11.2
30	10.7		11.2		12.4	14.8	13.5	15.1	15.5	18.1	13.7	17.0	14.8	12.1	11.2
40	10.7		10.9		11.4	12.4	12.3	12.2	11.9	12.3	12.4	12.6	14.7	12.0	11.2
50	10.7		10.8		11.0	11.5	11.6	11.6	11.3	11.4	11.6	11.4	11.6	11.2	11.2
60	10.7		10.7		10.7	11.0	11.2	11.0	11.0	11.1	11.4	11.1	11.1	10.9	11.2
70	10.7		10.5		10.6	10.9	10.8	10.8	10.8	10.9	11.6	11.1	10.9	10.8	11.2
80	10.5		10.5		10.6	10.9	10.7	10.7	10.7	10.8	11.2	10.9	10.8	10.8	11.2
90	10.4		10.5		10.6	10.7	10.7	10.7	10.7	10.7	11.3	10.8	10.7	10.8	10.8
100	10.4		10.5		10.5	10.6	10.6	10.7	10.7	10.7	10.9	10.8	10.8	10.7	10.8
110	10.4		10.5		10.5	10.5	10.6	10.7	10.7	10.6	10.8	10.8	10.7	10.7	10.8
120	10.4		10.5		10.5	10.5	10.5	10.6	10.6	10.6	10.7	10.7	10.7	10.7	10.8
130	10.4		10.5		10.5	10.5	10.5	10.7	10.6	10.6	10.7	10.7	10.7	10.7	10.8
140	10.4		10.5		10.5	10.5	10.5	10.6	10.6	10.6	10.6	10.7	10.7	10.7	10.8
150	10.4		10.5		10.5	10.5	10.5	10.6	10.6	10.6	10.6	10.7	10.7	10.7	10.8
160	10.4		*		10.5	10.5	10.5	*	*	*	*	*	*	*	*

Dissolved oxygen (g m⁻³)

Depth (m)															
1	9.6		10.3		9.5	8.5	8.5	8.1	8.2	8.4	8.7	8.6	9.0	9.2	9.3
10	9.6		10.5		9.9	8.7	8.5	8.1	8.2	8.3	8.7	8.6	9.0	9.2	9.1
20	9.6		10.6		10.0	9.1	9.1	8.2	8.1	8.3	8.8	8.6	8.9	9.2	9.1
30	9.6		10.7		10.5	9.7	10.1	9.2	9.0	8.1	9.0	8.4	8.9	9.1	9.0
40	9.7		10.7		10.5	10.1	10.2	9.5	9.1	8.7	8.8	8.7	8.9	9.0	8.9
50	9.6		10.3		10.3	9.9	9.9	9.0	9.0	8.6	8.6	8.4	8.7	8.4	8.8
60	9.5		10.3		10.0	9.6	8.9	8.7	8.8	8.5	8.5	8.4	8.5	8.1	8.7
70	9.4		10.2		10.0	9.6	8.9	8.6	8.6	8.5	8.5	8.4	8.3	7.9	8.7
80	9.4		10.2		9.9	9.6	8.8	8.5	8.5	8.4	8.3	8.4	8.3	7.8	8.6
90	9.0		10.1		9.8	9.5	8.8	8.4	8.4	8.3	8.2	8.3	8.2	7.7	8.1
100	9.0		10.0		9.7	9.4	8.8	8.3	8.3	8.3	8.2	8.3	8.1	7.7	7.5
110	9.0		9.9		9.6	9.4	8.8	8.1	8.3	8.2	8.1	7.9	7.8	7.6	7.3
120	8.8		9.9		9.4	9.3	8.3	8.1	8.3	8.1	8.3	7.9	7.8	7.5	7.1
130	8.8		9.8		9.3	9.2	8.3	7.9	8.2	7.8	8.3	7.8	7.5	7.5	7.1
140	8.7		9.6		9.1	8.9	7.9	7.6	8.2	7.5	8.0	7.6	7.7	7.4	7.0
150	8.7		9.2		8.9	8.7	7.9	7.6	8.0	7.4	7.8	7.4	7.5	7.4	7.0

Secchi depth

Depth (m)	11.9	11.9	13.0	13.6	15.1	16.3	15.7	17.8	18.4	14.1	14.6	14.4	14.7	14.4	12.9
-----------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Lake Taupo Temperature, Dissolved Oxygen, and Secchi Depth Database.
Started 27 October 1994
1994-1995
Temperature

Date	27-10-94	21-11-94	01-12-94	13-12-94	27-12-94	13-1-95	25-1-95	09-2-95	26-2-95	08-3-95	24-3-95	12-4-95	19-4-95	04-5-95	21-5-95	08-6-95	14-7-95	30-7-95
	Depth (m)																	
1	11.7	12.8	15.7	17.5	17.8	18.6	19.9	20.6	20.9	20.9	18.5	19.4	18.4	17.0	15.0	13.4	11.3	10.8
10	11.5	12.6	14.2	16.4	17.3	18.4	19.9	20.0	19.9	19.8	18.4	18.6	18.2	16.9	15.0	13.5	11.3	10.8
20	11.5	12.6	13.2	15.5	16.9	18.0	17.8	19.6	19.9	19.7	18.4	18.4	18.2	16.8	15.0	13.4	11.3	10.8
30	11.3	12.6	13.0	13.2	13.3	15.9	15.6	15.0	15.0	15.1	18.4	15.7	16.5	14.6	15.0	13.4	11.3	10.8
40	10.9	12.6	12.1	12.5	12.2	13.1	13.3	12.9	13.0	12.8	12.7	13.0	12.5	12.2	12.7	13.3	11.3	10.8
50	10.9	12.4	11.4	11.7	11.6	12.0	11.8	11.9	11.9	11.8	12.0	11.8	11.6	11.3	11.7	12.8	11.2	10.8
60	10.8	11.8	10.7	11.1	*	11.4	11.5	11.4	11.1	11.2	11.3	11.3	11.1	11.2	11.3	11.7	11.2	10.8
70	10.7	10.9	10.6	10.8	*	*	11.2	11.0	10.9	10.9	11.0	10.9	10.9	10.9	10.9	11.0	11.2	10.8
80	10.6	10.7	10.5	10.7	*	*	11.0	10.8	10.8	10.8	10.8	10.8	10.8	10.8	10.8	11.0	10.9	10.8
90	10.5	10.6	10.5	10.6	*	*	10.7	10.7	10.7	10.7	10.7	10.7	10.7	10.7	10.8	10.8	10.8	10.8
100	10.5	10.5	10.5	10.5	*	*	10.7	10.6	10.6	10.7	10.7	10.7	10.7	10.7	10.7	10.8	10.7	10.8
110	10.5	10.5	10.4	10.4	*	*	10.6	10.6	10.6	10.6	10.6	10.7	10.7	10.7	10.7	10.8	10.7	10.8
120	10.4	10.4	10.4	10.4	*	*	10.6	10.5	10.6	10.6	10.6	10.7	10.7	10.7	10.7	10.8	10.7	10.8
130	10.4	10.4	10.4	10.3	*	*	10.5	10.5	10.5	10.6	10.6	10.6	10.7	10.7	10.7	10.8	10.7	10.8
140	10.4	10.3	10.4	10.3	*	*	10.5	10.5	10.6	10.6	10.6	10.6	10.7	10.6	10.7	10.8	10.7	10.8
150	10.3	10.3	10.3	10.3	*	*	10.5	10.5	10.6	10.6	10.6	10.6	10.6	10.6	10.7	10.8	10.7	10.8
160	10.3	10.3	10.3	10.3	*	*	10.5	10.5	10.6	10.6	10.6	10.6	10.6	10.7	*	10.7	*	*

Dissolved oxygen (g m⁻³)

	Depth (m)																	
1	10.5	9.6	9.8	9.2	9.0	8.0	8.9	8.4	8.5	8.5	8.7	*	9.2	9.3	9.0	9.0	9.6	9.6
10	10.6	9.4	10.3	9.4	10.6	10.4	10.2	8.5	8.4	8.0	*	*	9.3	9.1	8.8	9.1	9.6	9.5
20	10.8	9.4	10.3	9.4	11.0	10.5	11.5	8.5	8.4	8.0	*	*	9.2	9.0	8.8	9.1	9.4	9.4
30	10.7	9.4	10.2	9.7	12.5	11.2	11.4	9.8	9.6	9.7	*	*	9.3	9.2	8.7	9.0	9.4	9.3
40	10.5	9.3	10.1	9.6	12.5	11.9	12.0	9.7	9.4	9.7	*	*	9.7	9.3	8.6	9.0	9.3	9.3
50	10.4	9.3	9.9	9.5	12.6	11.9	12.0	9.4	9.4	9.5	*	*	9.5	9.2	8.5	8.8	9.2	9.3
60	10.4	9.4	9.9	9.5	*	10.3	11.9	9.4	9.3	9.4	*	*	9.5	9.2	8.5	8.3	9.2	9.2
70	10.4	*	9.8	9.5	*	*	11.7	9.3	9.3	9.3	*	*	9.5	9.2	8.4	8.3	9.2	9.2
80	10.4	*	9.8	9.5	*	*	11.6	9.3	8.9	9.1	*	*	9.0	9.2	8.3	8.3	8.5	9.1
90	10.4	*	9.7	9.5	*	*	11.4	9.2	8.8	9.0	*	*	8.7	9.0	8.1	7.9	8.3	9.0
100	10.2	*	9.6	9.4	*	*	11.3	9.0	8.6	8.8	*	*	8.6	8.6	8.0	7.6	7.8	8.9
110	10.3	*	9.7	9.3	*	*	11.1	9.0	8.3	8.7	*	*	8.3	8.2	8.0	7.5	7.4	8.8
120	10.2	*	9.4	9.2	*	*	10.9	8.7	8.2	8.4	*	*	8.2	7.9	7.8	7.1	7.2	8.6
130	9.8	*	9.2	9.0	*	*	10.6	8.5	7.9	8.3	*	*	8.0	7.7	7.6	7.0	7.2	8.4
140	9.8	*	8.9	9.0	*	*	10.5	8.3	7.6	8.1	*	*	8.0	7.5	7.4	7.0	7.1	8.4
150	9.9	*	8.6	8.7	*	*	10.4	8.3	7.3	7.9	*	*	7.5	7.3	7.0	7.0	7.1	8.3
160	*	*	8.5	8.5	*	*	10.0	8.2	7.5	7.7	*	*	6.6	7.2	*	6.8	*	*

Secchi depth

Depth (m)	11.7	11.4	12.5	12.9	15.6	17.8	15.7	17.0	16.5	17.1	14.7	15.7	16.1	15.1	14.3	15.0	12.5	15.7
-----------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

* = missing or invalid data

8.4 Appendix 4: Nutrient data

Includes accumulated 10-m tube data since 1994. Blank cells represent missing data.

For completeness, 10-m tube data collected from the Kuratau Basin (site B) and Western Bays (site C) from January 2002 to December 2004 are included as separate sheets following the mid-lake data from site A for those years.

In the spring/autumn profile data, two different analytical methods are used to measure particulate nitrogen:

1. a wet digestion method involving high temperature refluxing in digestion mixture [persulphate / sulphuric acid / Selenium catalyst] for 3 hours followed by colorimetric determination of the nitrogen as the ammoniacal form; and
2. a CHN combustion method which converts all nitrogen compounds to N₂ gas in a furnace at ~1000°C to be measured in a thermal conductivity detector.

Particulate nitrogen analysed by the wet digestion method may not include some refractory nitrogen components which may be detected by the CHN combustion furnace method. Consequently the PN value from the CHN combustion furnace method should always be greater than or equal to the PN value obtained by the wet digestion method. Occasionally they are reported as less than the wet digestion method value in which case the wet digestion value should be regarded as correct. The cause of this difference is unknown but may be associated with the presence of low molecular weight organic nitrogen compounds lost during the drying step before combustion. The PN values for the time series data are all from wet digestion method analyses and hence are directly comparable with the profile data.

Low level NH₄-N results are likely to be subject to interference from low molecular weight DON and hence may not be biologically available for phytoplankton growth.

From February 2002, DRP, NO₃-N, and NH₄-N were measured on a Lachat Flow Injection Analysis (FIA) system but using essentially the same chemistry as previously used on the Technical Auto-Analyzer system. The reported detection limits for these nutrients remains the same at 0.5 mg m⁻³ for DRP and NO₃-N, and 1 mg m⁻³ for NH₄-N, however, the greater precision of the FIA system provides confidence in reporting results to a lower level as an indication of likely absolute values near zero. Such values are provided as an indication only and the true value should be expressed

as less than the detection limit. TN and TP values are the sum of all N and P components, excluding Urea-N which is part of the DON component. All analytical values ‘on-the-day’ are used wherever possible or $<DL = DL/2$ for summation in TN and TP. See **Appendix 1** for discussion on detection limits.

The DON value for 5/08/2000 was corrected from 12 to 43.5 in March 2006. This was a transcription error from the original analytical result sheet.

Lake Taupo cumulative database of 10m tube sample data from October 1994 to September 2002.

Samples collected from central lake site.

Date Collected	Temp. °C	Secchi m	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH₄-N mg m⁻³	NO₃-N mg m⁻³	DON mg m⁻³	PN mg m⁻³	TN mg m⁻³	Chlorophyll a mg m⁻³	PC mg m⁻³	
27/10/1994	11.7	11.7	1.2	0.7	2.5	4.4	1.1	0.2	56	16.6	73.4	1.16		
24/11/1994	12.8	11.4	0.5	2.7	1.7	4.8	1.7	1.0	51	12.6	66.5	0.41		
1/12/1994	15.7	12.5	0.6	2.4	2.4	5.4	2.2	1.2	56	18.5	78.0	0.41		
13/12/1994	17.5	12.9	0.8	4.2	1.4	6.4	<0.2	0.9	51	9.3	60.8	0.24		
28/12/1994	17.8	15.6	0.5	1.7	1.9	4.1	1.1	1.3	51	16.7	69.6	0.41		
13/01/1995	18.6	17.8	0.1	2.2	1.6	3.8	<0.2	0.8	53	11.6	64.9	0.22		
24/01/1995	19.9	15.7	0.2	2.1	1.2	3.6	<0.2	0.8	57	13.3	71.0	0.25		
10/02/1995	20.6	17.0	0.3	2.2	1.2	3.6	<0.2	1.5	62	10.2	73.3	0.32		
27/02/1995	20.8	16.5	0.4	<0.5	2.5	2.8	1.9	1.5	71	16.5	90.8	0.35		
30/02/1995	20.8	17.1	0.4	1.7	1.7	3.7	0.2	0.7	55	11.5	67.5	0.28		
24/03/1995	18.5	14.7	-	-	1.9	1.9	-	-	-	13.0	-	0.37		
12/04/1995	19.4	15.7	0.2	1.4	1.7	3.2	0.3	0.7	51	17.3	69.6	0.57		
19/04/1995	18.4	16.1	2.8	1.5	1.4	5.7	4.0	0.9	71	14.1	90.0	0.92		
4/05/1995	17.0	15.1	1.4	1.1	3.0	5.5	1.4	2.3	76	24.7	104.4	0.96		
21/05/1995	15.0	14.3	1.2	0.9	2.2	4.3	0.4	2.1	50	29.2	81.8	0.98		
8/06/1995	13.4	15.0	0.7	0.4	1.8	2.9	0.2	0.6	54	15.4	70.2	1.05		
14/07/1995	11.3	12.5	0.3	2.5	1.7	4.5	0.3	2.1	53	15.0	70.8	1.32		
30/07/1995	10.8	15.7	0.7	0.7	1.9	3.3	<0.2	4.6	35	17.3	57.3			
13/08/1995	-	-	0.5	0.4	1.9	2.8	<0.2	4.6	39	14.2	57.4	0.99		
12/09/1995	10.7	11.9	0.5	2.2	2.2	4.9	1.0	40.9	177	19.1	237.6	1.37		
25/09/1995	15.5	11.9	<0.2	0.7	2.1	2.8	<0.2	0.1	84	17.6	101.6	0.64		
30/10/1995	13.0	13.0	<0.2	2.4	1.9	4.3	<0.2	<0.1	56	14.7	70.4	0.93		
24/04/1996	13.7	13.6	0.8	1.8	1.6	4.3	1.9	<0.1	59	12.6	73.3	0.29		
6/12/1996	17.7	16.1	2.2	0.4	1.4	3.9	1.7	<0.1	64	11.3	70.8	0.20		
12/01/1996	21.1	16.3	2.6	0.6	1.2	4.4	3.6	<0.1	64	10.1	77.8	0.24		
3/01/1996	21.7	15.7	1.3	1.6	1.3	4.2	4.2	<0.1	59	11.9	75.5	0.29		
13/02/1996	22.7	17.8	2.1	3.3	1.2	6.6	7.4	<0.1	81	10.4	98.9	0.15		
29/02/1996	20.5	18.4	1.9	2.2	1.2	5.3	4.2	<0.1	61	10.8	76.3	0.31		
2/03/1996	18.2	14.1	0.8	2.2	1.4	4.5	5.4	<0.1	76	14.2	95.3	0.56		
28/03/1996	16.8	14.6	1.3	1.8	1.4	4.5	4.7	<0.1	91	12.6	108.3	0.81		
18/04/1996	17.7	14.4	0.8	2.2	-	-	4.3	<0.1	61	-	-	0.41		
19/05/1996	14.8	14.7	0.8	3.0	-	-	6.8	<0.1	59	-	-	0.70		
14/06/1996	12.2	14.4	1.6	3.2	-	-	5.7	<0.1	71	-	-	0.70		
19/06/1996	12.2	14.4	1.0	1.2	-	-	4.0	<0.1	49	-	-	0.70		
9/07/1996	11.2	12.9	3.0	-	1.9	-	4.0	<0.1	47	11.3	-	0.80		
3/09/1996	10.5	13.1	0.7	2.0	3.0	5.7	2.5	0.2	52	17.0	71.7	1.03		
18/09/1996	10.7	14.2	1.3	1.2	2.4	4.9	2.1	0.2	42	14.0	58.3	0.75		
30/09/1996	12.5	11.2	0.9	1.6	1.8	4.3	3.3	0.2	58	16.0	72.5	0.28		
17/10/1996	13.3	12.6	0.6	2.1	2.6	5.3	2.9	2.5	64	18.0	88.4	0.59		
24/04/1997	15.2	13.4	0.7	2.3	2.2	5.2	2.4	0.4	64	15.0	81.8	0.47		
6/11/1997	13.5	14.9	0.8	2.6	2.2	5.6	3.2	1.0	64	17.0	88.2	0.45		
28/11/1997	13.6	14.1	0.4	1.9	2.4	4.7	4.6	0.4	49	20.0	72.0	0.90		
11/12/1997	14.8	14.7	1.3	1.7	1.3	4.3	6.2	0.8	98	17.0	122.0	0.33		
23/12/1997	16.3	17.7	1.3	1.1	-	-	5.2	0.3	46	-	-	0.23		
8/01/1998	17.9	15.1	0.7	1.7	1.9	4.3	2.0	0.6	50	15.0	67.6	0.33		
29/01/1998	17.8	15.2	0.7	1.8	1.6	4.1	1.9	0.4	54	17.0	73.3	0.21		
26/03/1998	17.7	15.3	0.6	1.7	2.1	4.4	2.4	1.8	57	19.0	80.2	0.46		
2/04/1998	17.3	16.0	0.9	1.3	1.6	3.8	1.7	0.3	51	16.0	69.0	0.69		
15/04/1998	16.7	17.7	0.7	2.5	1.5	4.7	3.2	0.8	57	12.0	73.0	0.40		
1/05/1998	15.6	16.0	0.6	-	-	-	1.7	0.1	-	-	-	0.58		
21/05/1998	14.2	14.6	1.0	8.8	1.7	11.5	4.5	0.3	92	15.0	111.8	1.05		
29/05/1998	14.3	14.5	1.1	1.1	-	-	3.3	0.1	51	-	-	0.89		
7/07/1998	11.6	12.5	0.6	0.9	-	-	4.7	2.1	53	-	-	0.90		
29/07/1998	10.9	13.5	0.5	1.6	-	-	5.1	1.5	21	39	-	1.13		
2/08/1998	14.0	14.4	1.1	1.1	1.7	4.2	7.0	1.5	47	13.1	68.9	1.08		
16/08/1998	10.6	12.0	0.5	1.1	-	-	1.3	0.7	35	-	-	2.16		
11/10/1998	11.6	13.7	2.4	2.8	1.7	6.9	4.8	0.9	63.3	16.2	85.2	1.14		
29/10/1998	12.1	12.5	0.7	1.9	1.9	4.5	1.3	1.3	32	19.0	59.6	1.49		
2/12/1998	14.5	14.5	0.2	2.3	-	-	3.2	1.7	-	-	-	0.83		
21/01/1998	17.7	14.7	1.4	1.1	1.2	3.7	2.8	1.5	46.0	10.0	60.3	0.48		
2/03/1998	20.0	11.5	1.5	1.7	2.6	5.8	6.4	4.0	75.0	19.8	106.2	0.58		
24/03/1998	19.3	13.5	1.0	1.4	1.8	3.2	2.1	1.1	48.0	13.2	64.4	1.25		
7/04/1998	17.7	13.5	0.9	1.4	1.8	4.1	1.9	2.5	52.0	13.7	70.1	1.04		
29/05/1998	14.2	15.5	1.0	1.9	1.9	4.8	5.0	3.5	51.0	16.4	75.9	1.36		
28/07/1998	11.4	10.0	1.2	1.0	3.1	5.3	2.1	1.4	45.0	26.0	74.5	1.19		
29/09/1998	12.9	10.5	1.5	1.0	-	-	2.2	0.5	41.0	20.3	64.0	0.70		
8/10/1998	12.9	10.4	1.5	<1	-	-	2.4	2.4	46.0	37.6	88.4	1.00		
1/11/1998	13.6	13.5	0.6	1.3	2.6	4.5	4.5	<0.5	36.0	15.2	53.6	0.90		
26/11/1998	18.4	15.0	1.3	2.6	2.1	6.0	9.6	1.6	42.0	16.4	69.6	0.61		
22/12/1998	18.5	14.5	1.1	0.4	2.5	4.0	2.7	1.1	36.0	17.7	61.5	0.25		
2/01/1999	20.1	12.5	0.8	2.8	1.7	5.3	4.0	1.6	39.0	11.4	56.0	0.60		
3/03/1999	20.9	14.3	0.6	2.9	2.0	5.5	1.6	1.1	40.0	16.8	59.5	0.82		
14/04/1999	18.3	13.0	0.6	<1	1.8	2.4	3.0	<0.5	41.0	19.0	61.6	1.20		
30/04/1999	16.4	12.2	1.1	1.5	1.7	4.3	2.1	<0.5	38.0	19.6	60.2	0.94		
19/05/1999	14.4	15.0	0.8	<1	1.5	5.1	1	<1	46.0	16.2	63.7	1.2		
8/06/1999	14.1	14.5	1.0	<1	3.9	4.9	1	<1	48.0	25.4	74.9	1.1		
18/06/1999	13.0	15.0	0.8	<1	2.0	3.1	3.6	1	<1	45.0	28.3	74.3	1.0	
20/07/1999	12.0	16.0	0.5	<1	3.1	5.3	4	8	45.0	18.4	75.4	1.7		
9/08/1999	11.5	14.5	1.3	1.7	2.3	5.3	4	7	45.0	14.7	88.8	1.7		
6/09/1999	11.1	10.0	<0.5	2.5	2.1	5.1	2	1	60	16.2	79.2	0.5		
29/09/1999	11.5	10.0	0.7	1	4	5.7	3	1	54	32.6	90.6	1.8		
18/10/1999	12.7	14.9	0.5	3	2.5	6	<1	<1	41	19.4	60.4	0.4		
20/12/1999	16.4	18.0	0.7	2.3	5	8	4	2	39	38	83	1.6		
1/01/2000	17.6	15.1	0.9	2	2	4	5	2	42	52	72.0	1.7		
2/02/2000	15.0	14.5	1.1	2	1.5	3.5	1	<1	40	11	52.0	0.5		
16/02/2000	19.0	18.0	0.5	2.5	1.5	4.5	1	0.5	43.5	19.5	66.0	0.5		
2/03/2000	20.5	17.0	0.9	1.1	2	4.0	2	2	46	22.5	73.5	1.65		
1/04														

Lake Taupo cumulative database of 10 m tube sample data from June 2000 on
 Samples collected from Mid Lake (Site A)

Date Collected	Temp. °C	Secchi (m)	DRP (mg m ⁻³)	DOP (mg m ⁻³)	PP (mg m ⁻³)	TP (mg m ⁻³)	NH ₄ -N (mg m ⁻³)	NO ₃ -N (mg m ⁻³)	DON (mg m ⁻³)	PN (mg m ⁻³)	TN (mg m ⁻³)	Chlorophyll a (mg m ⁻³)	PC (mg m ⁻³)
20/06/2000	12.3	14.0	<1	4	0	4.0	2	2	52	16	72.0	1.7	193.5
11/07/2000	11.9	11.0	<1	4	3	7.0	3	2	46	22.5	73.5	1.65	198
5/08/2000	11.3	12.0	2	2	3	7.0	1	3.5	43.5	19.5	36.0	2.5	153.5
22/08/2000	11.2	15.0	2	2	2	6.0	2	4	49	16.5	71.5	1.65	158.5
12/09/2000	11.5	12.0	2	5	3.5	10.5	2	<1	63	23.5	88.5	1	148
29/09/2000	11.5	13.0	2	4	2	8.0	1	1	54	21	77.0	1.15	236.5
26/10/2000	13.1	11.0	0.8	4.2	3	8.0	1.0	0.4	41.6	25	68.0	1.3	237
14/11/2000	13.1	12.0	<1	4	2	6.0	1	<1	41	14.5	56.5	0.9	171
7/12/2000	15.1	17.0	2	2	1.55	5.6	7	4	63	14.75	88.8	0.6	165.5
4/01/2001	18.0	14.5	<1	2	1.5	3.5	1	<1	40	11	52.0	0.5	127
16/01/2001	19.0	18.0	0.5	2.5	1.5	4.5	1	0.5	53.5	13	68.0	0.5	118.5
21/02/2001	20.5	17.0	0.9	1.1	1.5	3.5	<1	0.5	46.5	12.5	59.5	0.6	190.5
2/03/2001	20.7	14.5	<1	2	2	4.0	2	<1	53	18	73.0	0.9	193
20/03/2001	19.0	17.0	<1	3	1.4	4.4	<1	<1	46	14.25	60.3	0.9	154
9/04/2001	17.0	13.5	0.8	1.2	2.15	4.2	<1	3	62	19.45	84.5	1.05	199
8/05/2001	15.8	17.0	0.8	3.2	1.7	5.7	2	<1	61	23	86.0	1.1	248
30/05/2001	13.6	14.5	1.5	1.5	2	5.0	1	<1	57	12	70.0	1.4	203
2/07/2001	12.1	12.0	<1	3	2.3	5.3	1	1	50	18.3	70.3	1.5	155.5
25/07/2001	11.3	14.5	2	1	2.65	5.7	<1	6	45	19.75	70.8	2.2	188
13/08/2001	11.2	13.5	1	1	2.85	4.9	1	<1	41	21.9	63.9	2.1	225
3/09/2001	10.2	17.5	1	1	2.6	4.6	<1	<1	37	19	56.0	1.7	203
25/09/2001	11.6	11.0	1.1	0.9	2.8	4.8	1	<1	56	24.5	81.5	0.9	283
25/10/2001	13.0	14.5	0.8	1.2	2.4	4.4	<1	<1	46	19.4	65.4	1.1	246
12/11/2001	14.3	15.5	1.0	2	2.55	5.6	0.9	0.1	48	17.6	66.6	0.5	227.5
10/12/2001	15.5	16.0	1.0	2	2.55	5.6	0.9	0.1	48	17.6	66.6	0.5	227.5
20/12/2001	17.0	13.0	0.6	2.7	2.05	5.4	1.3	0.1	48	14.85	64.3	0.5	203.5
8/01/2002	18.3	13.0	0.3	2	2.2	4.5	0	<1	50	17.15	67.2	0.8	246.5
22/01/2002	19.3	15.0	0	7	2.25	9.3	0	<1	40	20.35	60.4	0.9	188
6/03/2002	18.7	14.5	1.2	0.8	2.05	4.1	0.0	0.4	74	17.7	92.1	1.7	226.5
4/04/2002	17.4	19.0	0.6	3	1.45	5.1	1.1	0.1	46	10.7	57.9	0.8	138
17/04/2002	17.4	22.0	0.0	3	1.65	4.7	0.5	0.5	47	13.1	61.1	0.9	157
5/05/2002	15.5	16.4	0.7	1			3.1	0.7	48			1	
19/06/2002	12.6	17.0	1.2	1.8	1.9	4.9	0.5	1.4	43.6	15.8	61.3	1.1	165.0
1/07/2002	12.1	16.0	1.2	1.8	1.8	4.8	0.9	1.7	37.3	14.3	54.2	1.5	214
17/07/2002	11.4	15.5	2.3	2.7	1.7	6.7	2.3	7.8	41.9	14.6	66.6	1.5	153.5
31/07/2002	11.2	12.0	2.3	2.7	2.5	7.5	0.9	5.9	177.2	16.7	200.7	2.2	193

Date Collected	Temp. °C	Secchi (m)	DRP (mg m ⁻³)	DOP (mg m ⁻³)	PP (mg m ⁻³)	TP (mg m ⁻³)	NH ₄ -N (mg m ⁻³)	NO ₃ -N (mg m ⁻³)	DON (mg m ⁻³)	PN (mg m ⁻³)	TN (mg m ⁻³)	Chlorophyll a (mg m ⁻³)	PC (mg m ⁻³)
29/08/2002	11.1	9.5	1.6	1.4	3.1	6.1	0.0	0	90	23	113.0	2.6	196
18/09/2002	11.4	12	1.3	1.7	2	5.0	0	0.3	47	13	60.3	0.9	196.5
9/10/2002	11.6	15.5	1.3	2.7	2.1	6.1	2.9	0	29	12	43.9	0.6	159.5
13/11/2002	12.6	18	0.9	1.1	2.4	4.4	1.7	1.3	41	14.0	58.0	0.7	158.5
28/11/2002	14.1	12.7	0.7	2.3	2.7	5.7	0.1	0.0	43.0	22.0	65.1	0.7	201.5
18/12/2002	15.0	13.5	0.6	1.8	2.5	4.9	0.2	0.1	47.0	14.0	61.3	0.4	123.0
30/01/2003	17.8	18	0.4	3.6	1.9	5.9	0.4	0.1	56.5	12.0	69.0	0.7	166.0
13/02/2003	19.3	19	0.5	2.5	1.6	4.6	0.0	0.4	43.6	8.0	52.0	0.5	146.0
17/03/2003	18.5	15	0.8	2.2	1.7	4.7	<1	0.4	45.6	13.0	59.0	1.0	212
3/04/2003	19.3	13.5	1.1	2.9	1.8	5.8	<1	0.5	78.5	17.7	96.7	1.1	234.5
28/04/2003	16.7	14	0.3	3.7	1.9	5.9	<1	0.3	73.7	15.6	89.6	1.5	208.5
15/05/2003	15.6	16.5	0.1	3.9	2.2	6.2	0.3	0.3	50.4	19.5	70.5	1.4	228.5
12/06/2003	13.5	11	1.3	2.7	2.2	6.2	0.3	0.4	40.3	13.7	54.7	1.3	111.0
14/07/2003	11.8	14.5	2.2	1.8	2.6	6.6	1.1	1.1	34.8	18.0	55.0	1.8	102.0
31/07/2003	11.4	14	2.4	1.6	2.4	6.4	1.3	3.7	46.0	16.7	67.7	2.0	89.5
14/08/2003	11.2	13.5	1.8	2.2	3.1	7.1	0.7	0.2	46.1	21.1	68.1	2.9	91.5
26/08/2003	11.2	13	3.0	1.0	4.0	8.0	1.0	0.2	42.8	21.7	65.7	2.9	135.5
8/09/2003	11.1	12.5	2.6	0.4	3.3	6.3	0.4	0.2	45.2	17.4	63.2	1.5	199.5
7/10/2003	11.4	13.0	2.6	1.6	2.8	7.0	0.3	0.2	54.5	17.8	72.8	1.2	157.5
21/10/2003	13.0	17.0	2.0	1.0	2.3	5.3	0.1	1.3	39.6	14.0	55.0	0.6	146.0
19/11/2003	13.9	16.0	1.7	1.3	2.8	5.8	0.3	0.1	45.6	20.0	66.0	0.8	148.0
4/12/2003	16.0	18.5	1.6	2.4	1.8	5.8	0.2	0.1	53.7	13.4	67.4	0.3	106.5
18/12/2003	17.7	17.5	1.1	3.9	3.1	8.1	0.0	0.0	49.0	20.6	69.6	0.4	151.5
13/01/2004	20.3	19.0	0.5	3.5	1.6	5.6	0.0	0.3	52.0	12.5	64.8	0.4	127.0
26/02/2004	17.2	17.0	1.4	1.7	1.6	4.7	0.0	0.1	40.9	15.5	56.5	0.7	139.0
8/03/2004	17.5	15.0	0.6	2.4	2.0	5.0	0.4	0.1	42.5	12.4	55.4	0.6	177.5
31/03/2004	16.4	16.0	0.8	5.2	1.9	7.9	0.2	0.2	78.6	11.5	90.5	1.2	159.5
14/04/2004	15.3	15.0	1.0	3.0	2.4	6.4	0.1	0.3	46.6	16.0	63.0	1.3	187.5
10/05/2004	14.7	18.0	0.6	4.4	1.8	6.8	0.1	0.2	64.7	16.8	81.8	1.2	215.0
10/06/2004	13.6	13.5	0.9	2.1	2.1	5.1	0.0	0.6	63.4	17.8	81.8	1.0	371.5
13/07/2004	11.6	12.0	1.8	3.2	2.4	7.4	0.3	4.5	37.2	19.4	61.4	1.6	193.3
26/07/2004	11.3	11.0	1.6	2.4	3.0	7.0	0.5	2.4	38.1	23.4	64.4	2.7	196.0
24/08/2004	10.9	12.5	0.8	3.2	2.7	6.7	0.0	0.5	58.5	18.6	77.6	2.3	181.5
7/09/2004	10.7	12.0	0.6	2.4	2.7	5.7	0.0	0.1	40.9	15.5	56.5	1.4	162.5
21/10/2004	11.6	15.0	1.0	3.0	2.0	6.0	0.0	0.0	33.0	13.0	46.0	0.7	185.0
2/11/2004	12.9	16.0	1.0	3.0	1.9	5.9	2.2	0.8	62.0	14.7	79.7	0.6	147.0
22/11/2004	15.1	16.0	0.7	2.3	2.1	5.1	0.1	0.2	49.7	16.4	66.4	0.4	195.0
15/12/2004	14.1	19.5	0.7	3.3	2.2	6.2	0.0	0.2	45.8	14.7	60.7	0.2	127.5

Date Collected	Temp. °C	Secchi (m)	DRP (mg m ⁻³)	DOP (mg m ⁻³)	PP (mg m ⁻³)	TP (mg m ⁻³)	NH ₄ -N (mg m ⁻³)	NO ₃ -N (mg m ⁻³)	DON (mg m ⁻³)	PN (mg m ⁻³)	TN (mg m ⁻³)	Chlorophyll a (mg m ⁻³)	PC (mg m ⁻³)
11/01/2005	16.0	20	0.4	2.6	1.4	4.4	0	0.1	42.9	12.5	55.5	0.2	137
25/01/2005	19.3	19.5	0.5	2.5	1.5	4.5	0.0	0.1	54.9	14.5	69.5	0.3	131.0
9/02/2005	20.7	18	2.2	0.8	1.4	4.4	0.5	0.0	38.5	12.7	51.7	0.2	136.0
22/02/2005	20.0	21.5	0.8	5.2	1.7	7.7	1.5	0.5	58.0	15.8	75.8	0.2	159.0
10/03/2005	19.3	18.5	0.2	2.8	1.4	4.4	1.8	0.2	34.0	14.5	50.5	0.4	158.0
21/03/2005	19.3	20	0.8	3.2	1.2	5.2	0.5	0.1	43.4	10.0	54.0	0.5	140.0
14/04/2005	17.9	17.2	0.9	2.1	1.6	4.6	0.8	0.2	54.0	14.0	69.0	0.7	177.0
18/05/2005	14.3	16	0.8	2.2	1.9	4.9	0.0	0.5	46.5	13.9	60.9	1.3	177.5
9/06/2005	13.0	14.1	0.6	3.4	2.2	6.2	0.1	1.6	41.3	17.4	60.4	1.3	140.5
20/06/2005	12.7	13.8	0.6	3.4	2.0	6.0	0.1	1.0	39.9	18.5	59.5	1.2	158.5
20/07/2005	11.5	13	3.9	6.1	2.5	12.5	0.8	0.8	97.4	19.1	118.1	2.1	169
3/08/2005	11.1	14	2.6	1.4	2.3	6.3	2.0	1.4	61.6	20.3	85.3	1.2	116
17/08/2005	11.2	13	3.1	1	3.2	7.3	0.3	2.1	49.6	26.4	78.4	1.7	172.5
31/08/2005	11.7	13	2	1	2.4	5.4	<1	1	69	22.2	92.2	1.3	330
14/09/2005	12.4	13	1	1	2.5	4.5	<1	<1	60	19.9	79.9	0.8	243
29/09/2005	11.9	14	1	1	2.4	4.4	<1	<1	67	18	85	0.8	253.5
12/10/2005	11.9	14	0.7	2.3	2.7	5.7	0.0	0.7	56.3	23.2	80.2	0.8	301
25/10/2005	13.4	15	0.8	4.2	1.8	6.8	0.6	0.7	54.7	16.8	72.8	0.6	193
10/11/2005	16.3	17.5	1.2	3.8	1.5	6.5	0.2	0.1	52.7	15.6	68.6	0.5	160
1/12/2005	15.1	19.3	0.6	2.4	1.4	4.4	0	0.3	39.7	16.1	56.1	0.4	141
10/01/2006	17.4	19	1	2	1.4	4.4	0.1	1	49.9	17.8	68.8	0.5	167
2/02/2006	20.2	15.5	1.1	8.9	1.5	11.5	0.0	0.0	54	18	72	1.1	193.5
1/03/2006	19.5	15.3	0.3	7.7	1.6	9.6	0.0	1.3	38.7	18.5	58.5	0.9	160.5
12/04/2006	16.7	15.8	0.6	2.4	1.6	4.6	0.0	0.0	43	20.4	63.4	1.0	230
27/04/2006	16.3	17	1.0	2	1.6	4.6	0.1	0.0	52.9	17.6	70.6	1.1	196.5
9/05/2006	15.7	17.5	0.7	2.3	1.6	4.6	0.7	0.1	46.2	17.2	64.2	0.9	233
30/05/2006	14.2	18.2	0.8	2.2	1.6	4.6	1.8	0.9	61.3	16.6	80.6	1.3	233
27/06/2006	11.9	15.2	0.8	3.2	1.9	5.9	0.8	1.3	61.9	23.2	87.2	2	243
11/07/2006	11.5	13.5	1.4	5.6	2.3	9.3	0.2	1.7	93.1	21	116	1.7	209
25/07/2006	11.1	12	1.0	0	2.1	3.1	0.9	7.4	48.7	17.6	74.6	2.8	192
4/09/2006	11.1	11	1.8	1.2	2.5	5.5	0.0	0.6	31.4	24.5	56.5	2.8	218
26/09/2006	11.9	17.5	1.0	0.8	2.3	4.1	0.0	0.1	39.9	18.6	58.6	0.8	347
18/10/2006	11.7	13	0.8	1.2	2.5	4.5	0.0	0.3	35.7	18.2	54.2	0.9	227.5
1/11/2006	12.4	14.5	0.3	2.7	2.4	5.4	0.0	0.0	41	19.4	60.4	0.8	203
5/12/2006	14.7	16	0.0	3	2	5	0.0	0.0	52	20.2	72.2	0.7	186
19/12/2006	15.6	15.5	0.2	1.8	1.8	3.8	1.0	0.1	48.9	15.4	65.4	0.7	150
9/01/2007	16.5	13.5	0.5	1.5	1.6	3.6	0.9	0.4	60.7	15	77	0.3	207
25/01/2007	18.5	14.5	0.6	0	1.6	2.2	1.5	0.5	59	18.6	79.6	0.3	212
8/02/2007	19.3	16	0.6	0	1.6	2.2	0.4	0.5	58.1	16.8	75.8	0.4	156
21/02/2007	19.6	18.2	0.4	0	1.8	2.2	0.8	0.5	68.3	24.4	94	0.3	182
21/03/2007	18.6	16.5	1.1	0	2.1	3.2	1.8	1.3	47.2	22.1	72.4	0.8	175
3/04/2007	18.0	19	0.9	6.1	1.8	8.8	0.6	0.3	66.9	23.8	91.6	0.7	
19/04/2007	16.5	16	0.9	3.1	2.7	6.7	2.4	1.0	69.6	29.2	102.2	0.6	193

Date Collected	Temp. °C	Secchi (m)	DRP (mg m ⁻³)	DOP (mg m ⁻³)	PP (mg m ⁻³)	TP (mg m ⁻³)	NH ₄ -N (mg m ⁻³)	NO ₃ -N (mg m ⁻³)	DON (mg m ⁻³)	PN (mg m ⁻³)	TN (mg m ⁻³)	Chlorophyll a (mg m ⁻³)	PC (mg m ⁻³)
8/05/2007	19.3	16	1.1	3.9	1.2	6.2	0.3	0.4	63.3	17.8	81.8	1.2	169
22/05/2007	15.2	18.5	0.7	2.3	1.3	4.3	2.0	0.5	53.5	15.4	71.4	0.8	201
14/06/2007	13.6	18	0.6	2.4	1.8	4.8	4.0	0.8	65.2	21.8	91.8	1	159
27/06/2007	12.4	18.5	0.8	0.2	3.6	4.6	2.1	1.4	45.5	25.8	74.8	1.2	162
18/07/2007	11.4	14.5	1.1	1.9	2.9	5.9	1.3	1.0	44.7	37.8	84.8	1.7	
8/08/2007	11.1	14	1.1	1.9	2.8	5.8	2.0	2.2	46.8	28.2	79.2	1.3	229
23/08/2007	11.0	13	0.8	2.2	2.5	5.5	0.4	0.4	39.2	30.3	70.3	2.2	202
11/09/2007	11.0	11	1	4	3.3	8.3	0	1	67	34.7	102.7	1.4	324
9/10/2007	12.1	15	1	1	2.6	4.6	1.4	1.5	59.1	23.8	85.8	0.8	184
30/10/2007	12.8	16	1.1	0.9	2.4	4.4	1.2	0.6	64.2	30.5	96.5	0.7	253
15/11/2007	13.5	14	1.8	2.2	2.1	6.1	1.8	0.3	53.9	24.8	80.8	0.5	262
4/12/2007	16.6	15	0.9	2.1	2	5	0.9	0.6	40.5	20.6	62.6	0.3	196
20/12/2007	17.4	17.5	1.1	2.9	1.1	5.1	0.2	0.4	44.4	17	62	0.6	112
17/01/2008	21.1	22.5	1	4	1.5	6.5	0.9	0.4	62.7	24.5	88.5	0.3	230
31/01/2008	19.8	21.5	0.5	1.5	1.3	3.3	1.5	0.3	75.2	17.6	94.6	0.3	190
14/02/2008	19.9	25	0.3	1.7	1.6	3.6	1.4	0.7	75.9	19.8	97.8	0.4	138
27/02/2008	19.3	22	0.1	1.9	1.6	3.6	0.7	0.2	70.1	20	91	0.4	143
13/03/2008	18.8	22	1	1	1.2	3.2	1.2	0.6	56.2	19.6	77.6	0.5	147
26/03/2008	19.3	19	1	0	0.9	1.9	0.4	0.5	63.1	17.1	81.1	0.5	160
17/04/2008	17.8	20.5	1.2	0.8	1.3	3.3	1.1	1	51.9	14.2	68.2	0.8	189
7/05/2008	15.7	16	0.7	2.3	1.5	4.5	1.3	0.3	60.4	21.1	83.1	0.6	189
22/05/2008	14.7	17	0.2	1.8	1.5	3.5	0.4	0.4	71.2	23.6	95.6	0.7	191
5/06/2008	13.6	15	1.3	0.7	1.6	3.6	1	2.1	29.9	17.5	50.5	1	177
19/06/2008	12.9	16.5	0.5	1.5	1.6	3.6	2	0.7	34.3	29.2	66.2	1.2	259
1/07/2008	12.0	14	0.9	2.1	2.15	5.15	0.6	0.7	50.7	34.6	86.6	1.7	242
15/07/2008	11.4	13	1.3	1.7	2.7	5.7	0.0	0.9	38.1	26.5	65.5	1.9	193
7/08/2008	11.1	12.5	1.8	1.2	3.4	6.4	0.0	0.7	25.3	28.8	54.8	3.0	119
20/08/2008	10.7	12.5	1.3	1.7	2.1	5.1	0.7	0.6	24.7	25	51	1.5	179
4/09/2008	11.0	13	0.6	3.4	2	6	1.0	0.0	50	21.5	72.5	1.1	217
16/09/2008	11.3	14.5	1.4	2.6	2.1	6.1	2.2	0.5	28.3	24.3	55.3	0.7	202
14/10/2008	12.6	12.2	0.5	2.5	2.6	5.6	0.5	0.0	45.5	27.1	73.1	0.6	203
4/11/2008	13.4	12	1.0	4	2.5	7.5	3.2	0.5	35.3	28.5	67.5	0.9	140
26/11/2008	15.7	10	1.1	1.9	2.4	5.4	0.4	0.0	47.6	27.6	75.6	1	217
22/12/2008	18.8	12	0.3	1.7	2.3	4.3	1.8	0.0	53.2	35.2	90.2	0.6	245
13/01/2009	19.7	13	1.4	1.6	2.1	5.1	0.3	1.4	61.3	29.4	92.4	0.5	266
28/01/2009	20.9	18	0.4	4.6	1.8	6.8	0.0	3.8	52.2	27.6	83.6	0.3	204
11/02/2009	21.4	22	0.1	4.9	1.6	6.6	4.1	0.5	49.4	25.6	79.6	0.4	185.5
25/02/2009	20.5	20	0.5	2.5	1.6	4.6	2.7	0.4	37.9	21.3	62.3	0.5	186.5
26/03/2009	18.0	18.5	1.1	1.9	2.7	5.7	0.0	1.3	56.7	25.1	83.1	0.6	285
15/04/2009	16.6	18	1.5	2.5	3.4	7.4	1.1	0.7	60.8	22.7	85.3	0.8	240
7/05/2009	15.0	16	1.4	4.6	2.3	8.3	1.3	1.1	56.6	21.7	80.7	1.3	223
27/05/2009	13.0	15	1.2	4.8	1.5	7.5	0.0	0.6	58.4	16.7	75.7	1.2	190
18/06/2009	11.6	16	1.9	0.1	1.7	3.7	0.7	1.7	45.6	23.5	71.5	1.5	201
6/07/2009	10.9	15	2.8	1.2	2.4	6.4	0.1	8.1	46.8	23.4	78.4	1.6	190
13/08/2009	10.43	12	1.9	2.1	2.7	6.7	0.6	0.5	46.9	31.4	79.4	1.9	230
7/09/2009	10.56	15	4.2	0	2.9	7.1	0.1	0.6	54.3	32.3	87.3	1.5	301

Lake Taupo cumulative database of 10 m tube sample data
Samples collected from Kuratau Basin (Site B)

Date Collected	Temp. °C	Secchi m	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	PN mg m ⁻³	TN mg m ⁻³	Chlorophyll a mg m ⁻³	PC mg m ⁻³
8/01/2002	18.1	13.5	0.4	2	2.2	4.6	0.4	1.3	48	16.7	66.4	0.9	233
22/01/2002	18.8	12	0.9	2	2.6	5.5	0.9	0.3	41	19.9	62.1	0.9	221
6/03/2002	18.6	14.5	0.3	2	2.3	4.6	1.4	0.5	73	18.3	93.2	0.9	207
4/04/2002	17.4	19.5	0.6	2	1.5	4.1	0.4	0.1	40	11.2	51.7	0.9	162
17/04/2002	16.8	19	0.0	3	1.6	4.6	0.5	0.1	45	12.3	57.9	0.9	143
5/05/2002	15.1	13.2	0.3	1.1			1.6	0.4	40			0.9	
19/06/2002	12.5	15	1.0	1	2.2	4.2	0.4	0.8	48.2	17.4	66.8	1.5	182
1/07/2002	12.1	16	1.5	1.5	1.8	4.8	0.8	1.7	41.5	14.2	58.2	1.6	146
17/07/2002	11.5	12.5	1.8	2.2	2	6	0.8	5.1	51.1	16.1	73.1	1.5	156.5
31/07/2002	11.3	10.5	2.0	3	2.5	7.5	1.5	2.2	81.5	18.5	103.7	2.6	194.5
29/08/2002	11.0	8	1.2	4.8	3.3	9.3	0	0.2	184.0	22.9	207.1	2.3	221
18/09/2002	11.1	11	1.9	2.1	2.1	6.1	0.4	0.6	43.4	14	58.4	1.1	149
9/10/2002	11.7	16	1.4	1.6	1.7	4.7	4.4	0.2	19.6	11.7	35.9	0.5	149
13/11/2002	12.0	14	1	3	2.5	6.5	0.3	0	35	15.2	50.5	1.8	478
28/11/2002	13.8	12.7	0.9	2.9	2	5.8	0	0	40	16.7	56.7	0.7	203.5
18/12/2002	15.2	14	0.6	1.4	2.1	4.1	0	0.1	36	11.2	47.3	0.4	143
30/01/2003	16.8	18	0.5	2.5	1.7	4.7	<1	0.8	43	12.1	55.9	0.6	148.5
13/02/2003	18.8	11	0.7	1.3	1.6	3.6	0.4	0.2	45	9.3	54.9	0.7	131
17/03/2003	18.7	14	0.5	3.5	2	6	<1	0.7	49	16.3	66.0	1.0	208
3/04/2003	19.0	12.8	0.6	3.4	2.1	6.1	<1	0.1	50	19.6	69.7	1.1	239.5
28/04/2003	16.7	13.5	0.6	3.4	1.6	5.6	<1	0.2	57	13.1	70.3	1.4	218.5
15/05/2003	15.7	15.5	0.4	3.6	1.8	5.8	<1	0.2	63	13.5	76.7	1.7	229.5
12/06/2003	12.5	12	1.7	1.3	2.2	5.2	0.1	2.8	39.1	13.9	55.9	1.3	
14/07/2003	11.8	12	1.7	2.3	2.2	6.2	0.9	1.9	39.4	15.9	58.1	1.7	96.5
31/07/2003	11.3	13	2.1	1.9	2.7	6.7	1.2	2.0	43.8	18.0	65.0	2.1	108.5
14/08/2003	11.4	13	1.8	2.2	3.3	7.3	0.3	0.3	33	22.3	55.9	2.5	112.0
26/08/2003	11.3	11.5	3.1	0.9	4.0	8	0.4	0.1	37	22.4	59.9	3.1	148.0
8/09/2003	11.1	11	2.5	1.5	3.3	7.3	0.4	0.1	36	23.5	60.0	1.4	196.5
7/10/2003	11.7	9.5	2.3	1.7	3.0	7.0	0.0	0.1	49.9	20.5	70.5	1.2	185.5
21/10/2003	13.2	15.0	2.2	0.8	2.7	5.7	0.3	0.2	38.5	14.9	53.9	0.8	155.5
19/11/2003	13.8	17.0	1.6	2.4	2.4	6.4	0.0	0.1	51.0	14.6	65.7	0.6	139.5
4/12/2003	15.6	17.0	1.8	2.2	1.8	5.8	0.2	0.1	44.7	13.5	58.5	0.4	126.5
18/12/2003	17.0	15.0	0.5	3.5	1.9	5.9	0.0	0.2	56.0	12.4	68.6	0.5	145.5
13/01/2004	20.3	16.0	0.4	4.6	1.8	6.8	0.0	0.2	54.0	13.7	67.9	0.5	125.0

Date Collected	Temp. °C	Secchi (m)	DRP (mg m ⁻³)	DOP (mg m ⁻³)	PP (mg m ⁻³)	TP (mg m ⁻³)	NH ₄ -N (mg m ⁻³)	NO ₃ -N (mg m ⁻³)	DON (mg m ⁻³)	PN (mg m ⁻³)	TN (mg m ⁻³)	Chlorophyll a (mg m ⁻³)	PC (mg m ⁻³)
26/02/2004	16.8	13.5	1.1	1.9	1.8	4.8	0.6	0.1	42.3	15.8	58.8	0.8	157.0
8/03/2004	17.6	5.0	0.8	2.2	3.1	6.1	1.0	0.3	41.7	17.5	60.5	0.9	172.0
31/03/2004	15.9	11.0	0.8	3.2	1.8	5.8	0.7	0.2	45.1	9.9	55.9	1.4	124.5
14/04/2004	15.0	14.0	0.9	4.1	2.2	7.2	0.6	0.3	52.1	14.9	67.9	1.3	171.5
10/05/2004	14.7	15.5	0.8	2.2	1.7	4.7	0.0	0.2	59.8	15.9	75.9	1.3	179.0
10/06/2004	12.9	12.0	1.4	2.6	2.1	6.1	0.0	0.2	108.8	18.6	127.6	1.2	183.0
13/07/2004	11.4	11.0	2.1	2.9	2.5	7.5	0.0	8.4	40.6	19.3	68.3	1.4	154.0
26/07/2004	11.2	10.0	1.3	2.7	3.2	7.2	0.2	5.8	38.0	25.0	69.0	2.7	204.0
24/08/2004	10.9	10.0	0.7	3.3	3.1	7.1	0.0	0.0	47.0	20.9	67.9	2.5	158.0
7/09/2004	10.8	11.0	0.7	2.3	2.6	5.6	0.0	0.2	44.8	17.1	62.1	1.5	172.5
21/10/2004	11.7	11.0	1.2	1.8	2.1	5.1	0.2	0.0	30.8	16.1	47.1	0.8	172.5
2/11/2004	13.1	15.0	1.0	2.0	1.7	4.7	0.2	0.1	42.7	11.0	54.0	0.5	152.0
22/11/2004	14.9	15.0	0.6	3.4	1.6	5.6	0.6	0.0	33.4	9.5	43.5	0.5	141.5
15/12/2004	13.2	17.2	0.6	3.4	1.6	5.6	0.4	0.1	39.5	12.6	52.6	0.2	120.0

Lake Taupo cumulative database of 10 m tube sample data
Samples collected from Western Bays (site C)

Date Collected	Temp. °C	Secchi m	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	PN mg m ⁻³	TN mg m ⁻³	Chlorophyll a mg m ⁻³	PC mg m ⁻³
8/01/2002	18.72	14.5	0.9	4	2.3	7.2	0.9	0.6	88	16.1	105.6	0.8	213
22/01/2002	18.82	15.5	0.7	2	2.2	4.9	0.7	0.0	37	16.8	54.5	0.8	221
6/03/2002	18.68	16	0.2	2	2	4.2	0	0.1	45	16	61.1	0.7	177
4/04/2002	17.47	19	0.6	2	1.4	4	0.0	0.0	38	8.8	46.8	0.9	152
17/04/2002	16.88	18.5	0	3	1.6	4.6	0.7	0.2	44	11.8	56.7	0.9	167
5/05/2002	15.6	15.6	0.4	1		2	0.2	0.2	45			1.1	
19/06/2002	12.58	16	0.9	2.1	2	5	0.3	1.2	38.8	15.9	56.2	0.9	161
1/07/2002	12.22	14	1.3	1.7	1.9	4.9	0.3	0.4	45	15	60.7	1.4	148
17/07/2002	11.52	12.5	1.9	2.1	2	6	0.9	4.9	46.1	16.3	68.2	1.5	160
31/07/2002	11.6	12	2.3	2.7	2.3	7.3	1.7	4.0	113.3	16.7	135.7	2.3	150
29/08/2002	11.4	8	1	3	3.2	7.2	0	0	177	22.3	199.3	2.4	217
18/09/2002	11.24	12	2.8	2.2	2	7	1.7	0.4	45.3	11.7	59.1	0.9	152
9/10/2002	12.10	19	1.5	1.5	1.7	4.7	0.3	0.2	28	10.2	38.7	0.4	116
13/11/2002	12.60	16	1.1	2.9	2	6	0.1	0	51	12.2	63.3	0.6	141
28/11/2002	13.90	15.5	0.9	2.1	2	5	0.4	0.4	40	14.4	55.2	0.8	125.5
18/12/2002	15.10	13.5	0.8	2.2	1.9	4.9	0	0.3	45	10.2	55.5	0.5	136.5
30/01/2003	17.60	18.5	0.5	2.5	1.5	4.5	<1	0.1	46	8.6	54.7	0.4	141.5
13/02/2003	19.50	19	0.6	1.4	1.6	3.6	0	0.1	42	8.4	50.5	0.5	104
17/03/2003	18.70	15	0.5	2.5	1.7	4.7	<1	0.4	46	14.6	61.0	1.1	215
3/04/2003	18.80	14.5	0.5	2.5	1.6	4.6	<1	0.4	49	16.5	65.9	1.2	204
28/04/2003	17.00	14.5	0.4	2.6	1.4	4.4	<1	0.4	54	12.2	66.6	1.5	191
15/05/2003	15.60	17	0.1	3.9	2.2	6.2	<1	0.1	56	18	74.1	1.3	197
12/06/2003	13.70	11	1.3	1.7	2	5	0.1	0.9	40	13.8	54.8	1.3	
14/07/2003	11.80	14	1.9	2.1	2	6	1	4.7	39.3	14.9	59.9	1.5	85.0
31/07/2003	11.40	12	3.1	5.9	2.8	11	0.1	4.0	55	20.3	79.4	2.3	101.5
14/08/2003	11.50	14.5	2.4	2.6	2.9	7.9	1.1	3.8	46.1	19.5	70.5	2.8	92.5
26/08/2003	11.30	13	2.8	2.2	3.8	8.8	0.5	0.2	39	25.0	64.7	3.2	174.5
8/09/2003	11.30	12	2.6	0.4	3	6	0.1	0.1	40	19.5	59.7	1.3	233.0
7/10/2003	11.7	12.5	2.7	1.3	2.8	6.8	0.0	0.3	44.7	18.4	63.4	1.5	157.5

Date Collected	Temp. °C	Secchi (m)	DRP (mg m ⁻³)	DOP (mg m ⁻³)	PP (mg m ⁻³)	TP (mg m ⁻³)	NH ₄ -N (mg m ⁻³)	NO ₃ -N (mg m ⁻³)	DON (mg m ⁻³)	PN (mg m ⁻³)	TN (mg m ⁻³)	Chlorophyll a (mg m ⁻³)	PC (mg m ⁻³)
21/10/2003	13.0	12.0	1.5	1.5	3.1	6.1	0.3	0.0	44.7	17.4	62.4	1.1	195.0
19/11/2003	14.3	17.2	1.5	1.5	2.3	5.3	0.8	0.0	38.2	14.4	53.4	0.7	123.0
4/12/2003	15.5	17.0	1.7	3.3	1.7	6.7	0.0	0.2	46.8	11.2	58.2	0.5	129.0
18/12/2003	17.0	19.0	0.5	4.5	1.5	6.5	0.0	0.0	47.0	9.9	56.9	0.4	124.5
13/01/2004	20.2	17.5	0.7	4.3	1.6	6.6	0.0	0.1	53.0	11.9	65.0	0.4	118.5
26/02/2004	16.9	14.0	0.9	2.1	2.2	5.2	0.8	0.4	40.8	17.2	59.2	0.7	156.0
8/03/2004	18.4	13.0	0.8	2.2	2.0	5.0	0.7	0.1	34.2	11.1	46.1	0.6	124.0
31/03/2004	16.4	12.5	0.6	3.4	2.0	6.0	0.7	0.3	51.0	12.3	64.3	1.2	175.5
14/04/2004	15.4	16.5	0.9	3.1	2.3	6.3	0.6	0.3	50.1	14.2	65.2	1.2	159.0
10/05/2004	14.9	16.0	0.8	3.2	1.6	5.6	0.0	0.2	48.8	15.4	64.4	1.1	153.0
10/06/2004	13.1	14.0	0.8	2.2	2.0	5.0	0.0	0.2	41.8	16.6	58.6	1.0	151.0
13/07/2004	11.6	12.5	1.3	2.7	2.5	6.5	0.0	5.9	39.1	19.9	64.9	1.6	156.5
26/07/2004	11.5	11.0	1.5	2.5	2.9	6.9	0.3	2.7	46.0	22.2	71.2	2.4	180.5
24/08/2004	10.9	10.0	1.0	3.0	2.9	6.9	0.0	0.4	37.6	18.5	56.5	2.5	161.0
7/09/2004	11.1	12.0	1.2	3.8	2.6	7.6	0.0	0.0	54.0	16.8	70.8	1.5	202.0
21/10/2004	11.7	12.0	1.1	1.9	1.9	4.9	0.2	0.0	35.8	14.8	50.8	0.6	167.5
2/11/2004	12.4	17.0	1.0	3.0	1.7	5.7	0.3	1.2	45.5	16.3	63.3	0.4	173.0
22/11/2004	14.8	16.0	0.5	3.5	1.7	5.7	0.0	0.2	37.8	10.8	48.8	0.5	149.0
15/12/2004	14.2	20.8	0.9	4.1	1.4	6.4	0.0	0.0	42.0	12.2	54.2	0.2	131.0

Lake Taupo biannual nutrient database
2008-2009

Started 27 October 1994

Collection date 14 October 2008

Code	Depth m	Secchi depth = 12.2 m																		
		pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³
SZ1	1	7.66	119	12.59	10.29	1.1	<0.5	0.7	1.0	2.0	4.2	7.2	4.1	0.0	70.9	26.1	101.1	816	235.0	24.6
SZ2	10	7.70	121	12.09	10.29	0.7	<0.5	0.8	0.6	2.4	3.9	6.9	0.1	0.0	39.9	18.7	58.7	690	169.5	23.5
SZ3	20	7.70	121	11.93	10.50	0.8	<0.5	0.8	0.7	2.3	7.8	10.8	0.0	0.0	59.0	32.7	91.7	638	250.0	33.1
SZ4	30	7.70	120	11.85	10.46	1.0	0.6	0.7	0.7	2.3	5.6	8.6	0.0	0.0	65.0	24.2	89.2	632	195.5	31.8
SZ5	40	7.70	120	11.75	10.34	0.7	<0.5	0.9	0.3	1.7	4.6	6.6	0.0	0.0	52.0	16.2	68.2	597	162.5	15.5
SZ6	50	7.69	120	11.59	10.05	0.5	<0.5	0.9	0.4	2.6	4.5	7.5	0.5	0.0	48.5	15.6	64.6	602	139.5	29.2
SZ7	60	7.56	120	10.90	9.89	0.8	0.5	0.8	1.0	2.0	5.0	8.0	0.7	1.6	69.7	16.7	88.7	603	94.0	18.2
SZ8	70	7.52	121	10.76	9.86	0.6	<0.5	0.6	1.2	1.8	3.6	6.6	0.0	2.6	45.4	20.4	68.4	593	77.2	16.8
SZ9	80	7.45	122	10.71	9.81	0.7	<0.5	0.4	1.3	2.7	3.1	7.1	0.0	4.7	36.3	9.5	50.5	589	61.8	25.9
SZ10	90	7.49	121	10.69	9.85	0.7	<0.5	0.3	1.8	0.2	2.3	4.3	0.0	5.7	29.3	9.7	44.7	561	57.5	9.1
SZ11	100	7.23	121	10.68	10.03	0.6	<0.5	0.2	1.5	0.5	2.5	4.5	2.2	6.6	33.2	9.2	51.2	605	71.8	23.1
SZ12	110	7.32	121	10.66	10.13	<0.5	<0.5	0.3	1.5	1.5	2.2	5.2	3.5	7.4	33.1	8.0	52.0	617	46.8	10.6
SZ13	120	7.36	122	10.64	10.09	0.7	<0.5	0.2	1.2	2.8	2.5	6.5	1.6	9.5	34.9	9.9	55.9	613	57.6	28.5
SZ14	130	7.45	121	10.60	9.83	0.8	<0.5	0.2	2.6	0.4	2.1	5.1	1.6	11.7	34.7	7.5	55.5	652	56.6	27.2
SZ15	140	7.43	120	10.59	9.76	<0.5	<0.5	<0.1	2.9	3.1	2.5	8.5	1.4	17.1	37.5	8.7	64.7	686	46.6	24.1
SZ16	150	7.40	121	10.59	9.85	<0.5	<0.5	0.2	2.7	2.3	3.5	8.5	2.3	17.3	39.4	11.0	70.0	656	68.9	23.5

Collection date 15 April 2009

Code	Depth m	Secchi depth = 18.0 m																			
		pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
EU1	1	7.89	123	16.60	9.33	<0.5	<0.5	0.7	1.1	0.9	1.7	3.7	4.3	1.4	74.3	17	16.7	96.7	834	187.0	19.2
EU2	10	7.84	122	16.59	10.11	<0.5	<0.5	0.8	1.3	1.7	2.0	5.0	0.1	0.0	26.9	<1	13.1	40.1	669	116.0	16.2
EU3	20	7.83	121	16.59	10.76	<0.5	<0.5	0.9	1.2	2.8	2.0	6.0	0.3	0.0	29.7	1	17.2	47.2	691	152.0	18.4
EU4	30	7.84	123	16.58	10.83	<0.5	<0.5	0.9	0.9	3.1	1.8	5.8	0.8	0.0	38.2	2	15.8	54.8	650	143.0	19.1
EU5	40	7.8	121	12.53	10.39	<0.5	<0.5	1.0	1.4	6.6	1.5	9.5	0.7	0.1	37.3	1	13.0	51.1	627	81.9	13.2
EU6	50	7.79	121	11.56	9.58	<0.5	<0.5	0.7	2.2	3.8	1.2	7.2	0.0	2.0	20.0	<1	9.3	31.3	574	79.5	12.1
EU7	60	7.58	122	11.12	9.06	<0.5	<0.5	0.5	3.9	3.1	1.2	8.2	0.0	8.5	24.5	2	7.4	40.4	581	68.6	11.6
EU8	70	7.49	123	10.98	8.84	<0.5	<0.5	0.3	5.5	4.5	1.1	11.1	0.7	18.7	14.6	2	8.7	42.7	553	59.6	15.2
EU9	80	7.03	124	10.92	8.21	<0.5	<0.5	0.2	6.6	6.4	1.2	14.2	0.0	24.5	26.5	<1	9.3	60.3	635	51.7	11.8
EU10	90	7.03	124	10.88	8.24	12	12	0.1	7.2	2.8	1.1	11.1	0.0	27.0	16.0	1	6.7	49.7	514	46.6	9.4
EU11	100	7.16	123	10.86	8.07	<0.5	<0.5	0.1	6.3	5.7	0.9	12.9	0.0	24.7	32.3	1	5.1	62.1	554	35.9	8.8
EU12	110	7.21	124	10.84	8.12	<0.5	<0.5	0.1	7.0	4	1.0	12.0	0.2	26.3	12.5	<1	6.9	45.9	562	42.7	10.1
EU13	120	7.2	123	10.82	8.02	<0.5	<0.5	0.1	7.1	4.9	1.0	13.0	0.2	26.8	25.0	4	6.8	58.8	549	53.7	10.1
EU14	130	7.61	123	10.79	8.15	<0.5	<0.5	<0.1	7.6	8.4	1.0	17.0	0.0	27.6	<1	2	7.2	34.8	562	45.4	11.8
EU15	140	7.23	122	10.78	8.01	<0.5	<0.5	<0.1	8.1	4.9	1.1	14.1	0.0	29.0	8.0	<1	7.3	44.3	661	50.3	9.8
EU16	150	7.22	122	10.78	7.55	<0.5	<0.5	<0.1	9.0	2	1.3	12.3	1.3	30.6	21.1	1	7.1	60.1	544	42.8	12.7

 NH₄, NO₃, DON, Urea all as N

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient database
2007-2008

Started 27 October 1994

Collection date 30 October 2007

Code	Depth m	Secchi depth = 12.8 m																			
		pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
ZA1	1	7.80	119	12.84	10.18	0.7	<0.5	0.6	1.3	0.7	2.1	4.1	1.5	0.7	79.8	16	20.1	102.1	617	170.0	19.2
ZA2	10	7.83	120	11.83	10.27	<0.5	<0.5	1.0	0.9	1.1	2.5	4.5	0.0	0.0	42.0	<5	18.5	60.5	553	204.0	19.8
ZA3	20	7.79	115	11.76	10.25	0.5	<0.5	1.1	1.1	0.9	2.6	4.6	0.2	0.0	42.8	<5	19.0	62.0	405	169.0	19.4
ZA4	30	7.76	119	11.70	10.07	0.7	<0.5	1.2	0.8	1.2	2.5	4.5	0.0	0.0	49.0	<5	19.1	68.1	417	173.5	19.0
ZA5	40	7.72	120	11.64	10.02	0.7	<0.5	1.1	1.0	1.0	2.6	4.6	0.0	0.0	36.0	<5	16.8	52.8	417	131.5	17.4
ZA6	50	7.61	121	11.51	9.85	0.8	<0.5	1.4	0.9	1.1	3.3	5.3	0.0	0.0	39.0	<5	18.3	57.3	434	140.0	18.1
ZA7	60	7.54	120	11.43	9.52	0.9	<0.5	1.4	1.2	0.8	2.7	4.7	0.2	0.0	32.8	<5	19.5	52.5	414	127.5	17.1
ZA8	70	7.46	123	11.32	9.77	0.8	<0.5	1.5	1.5	0.5	2.7	4.7	0.1	0.3	46.6	<5	19.1	66.1	443	130.0	19.0
ZA9	80	7.42	122	11.23	9.58	0.8	<0.5	1.1	1.9	1.1	2.1	5.1	0.4	2.6	41.0	5	15.8	59.8	422	95.8	14.4
ZA10	90	7.42	121	11.16	9.42	0.7	<0.5	0.9	2.1	0.9	2.1	5.1	0.3	4.8	42.9	<5	13.3	61.3	410	92.0	13.0
ZA11	100	7.38	122	11.07	9.49	<0.5	<0.5	0.7	2.8	0.2	1.8	4.8	0.0	8.5	36.5	<5	11.2	56.2	400	64.0	11.0
ZA12	110	7.40	122	11.04	9.16	0.7	<0.5	0.7	2.9	0.1	1.8	4.8	0.0	9.2	56.8	<5	11.6	77.6	386	68.3	11.1
ZA13	120	7.38	122	11.02	9.27	0.7	<0.5	0.6	2.8	1.2	2.1	6.1	0.0	10.0	46.0	<5	12.7	68.7	359	105.3	12.5
ZA14	130	7.44	120	11.00	9.01	0.6	<0.5	0.6	2.6	1.4	1.9	5.9	0.0	10.4	35.6	<5	10.9	56.9	348	61.8	10.5
ZA15	140	7.44	121	10.98	9.11	0.6	<0.5	0.6	3.0	0.0	1.7	4.7	0.0	10.8	39.2	<5	10.3	60.3	351	64.1	11.2
ZA16	150	7.42	121	10.96	8.91	<0.5	<0.5	0.6	3.5	1.5	1.8	6.8	0.0	13.3	38.7	<5	10.8	62.8	305	63.1	10.6

Collection date 17 April 2008

Code	Depth m	Secchi depth = 17.8 m																			
		pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
KA1	1	7.79	122	17.88	9.49	<0.5	<0.5	0.4	0.8	0.2	0.7	1.7	2.8	0.4	64.8	14	13.3	81.3	656	138.5	8.4
KA2	10	7.87	121	17.87	8.97	<0.5	<0.5	0.8	0.5	0.5	0.7	1.7	1.1	0.3	48.6	<5	12.0	62.0	576	112.5	8.3
KA3	20	7.83	124	17.85	8.46	<0.5	<0.5	0.8	0.9	0.1	0.8	1.8	0.4	0.3	38.3	<5	13.7	52.7	528	142.0	9.4
KA4	30	7.71	122	15.58	8.52	<0.5	<0.5	0.5	1.0	0.0	0.9	1.9	3.1	0.1	27.8	<5	10.9	41.9	526	110.0	9.1
KA5	40	7.58	121	12.38	8.72	<0.5	<0.5	0.6	1.7	1.3	0.8	3.8	1.8	0.8	36.4	<5	14.6	53.6	459	107.0	6.7
KA6	50	7.38	121	11.72	8.48	<0.5	<0.5	0.5	1.9	2.1	0.6	4.6	0.2	3.4	29.4	<5	10.2	43.2	417	75.1	6.1
KA7	60	7.36	122	11.48	8.20	<0.5	<0.5	0.4	3.5	0.5	0.8	4.8	0.6	5.3	32.1	<5	9.6	47.6	353	84.9	6.7
KA8	70	7.31	122	11.34	7.84	<0.5	<0.5	0.3	3.5	1.5	0.7	5.7	0.9	10.8	42.3	<5	10.7	64.7	481	85.4	6.8
KA9	80	7.25	122	11.27	7.71	<0.5	<0.5	0.2	4.2	0.8	1.2	6.2	0.4	14.7	82.9	<5	9.5	107.5	347	97.5	4.9
KA10	90	7.19	122	11.20	7.57	<0.5	<0.5	0.1	5.1	0.0	0.7	5.8	0.3	19.8	43.9	<5	10.2	74.2	370	107.0	5.4
KA11	100	7.18	122	11.17	7.45	<0.5	<0.5	0.1	4.6	0.6	5.2	6.6	0.6	21.2	30.2	<5	8.6	60.6	412	59.8	4.0
KA12	110	7.12	123	11.14	7.29	<0.5	<0.5	0.1	5.0	1.0	0.6	6.6	0.8	28.2	26.0	<5	4.5	59.5	346	44.6	3.3
KA13	120	7.07	123	11.15	7.29	0.6	<0.5	<0.1	7.4	0.0	0.8	8.2	0.1	30.2	29.7	<5	7.9	67.9	373	85.8	5.8
KA14	130	7.28	123	11.12	7.18	<0.5	<0.5	<0.1	5.6	1.4	0.8	7.8	1.1	29.5	26.4	<5	9.0	66.0	395	89.1	4.4
KA15	140	7.12	123	11.11	7.13	<0.5	<0.5	<0.1	8.4	1.6	1.5	11.5	1.1	36.8	27.1	<5	8.5	73.5	393	72.6	4.1
KA16	150	7.11	123	11.11	6.72	<0.5	<0.5	8.3	0.7	1.5	10.5	0.4	36.4	27.2	<5	7.2	71.2	379	98.8	4.1	

 NH₄, NO₃, DON, Urea all as N

* = PN by wet digestion method, ** = PN by combustion furnace method.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient database
2006-2007

Started 27 October 1994

Collection date 1 November 2006

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	Secchi depth = 14.5 m		DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
							VSS g m ⁻³	Chlor_a mg m ⁻³													
HW1	1	7.79	118	12.43	10.2	0.5	<0.5	0.5	1.2	0.0	1.7	2.9	0.1	1.0	75.9	13.6	90.6	413	168.0	15.4	
HW2	10	7.77	119	12.27	10.1	0.8	<0.5	0.6	1.0	0.0	1.9	2.9	0.0	0.1	61.9	13.8	75.8	419	187.0	13.8	
HW3	20	7.77	120	12.25	10.1	0.7	<0.5	0.7	0.9	1.1	2.3	4.3	0.0	0.1	32.9	17.8	50.8	373	209.5	17.4	
HW4	30	7.81	119	12.20	10.1	0.8	<0.5	0.6	1.0	0.0	2.7	3.7	0.3	0.0	38.7	22.3	61.3	456	215.5	18.1	
HW5	40	7.78	119	12.10	10.1	0.9	<0.5	0.6	1.1	0.9	2.2	4.2	0.0	0.1	30.9	17.9	48.9	368	227.5	19.8	
HW6	50	7.74	119	11.96	10.0	0.6	<0.5	0.7	1.2	0.0	1.9	3.1	0.0	0.2	29.8	14.0	44.0	468	169.0	13.9	
HW7	60	7.67	120	11.34	9.7	0.7	<0.5	1.1	1.5	0.0	1.8	3.3	0.6	0.1	31.3	13.9	45.9	411	123.5	13.5	
HW8	70	7.64	119	11.17	9.5	<0.5	<0.5	1.3	1.2	1.8	2.0	5.0	0.5	0.1	29.4	14.5	44.5	378	98.0	12.3	
HW9	80	7.57	119	11.06	9.4	0.7	<0.5	1.3	1.3	0.7	2.2	4.2	2.5	1.8	27.7	14.1	46.1	330	91.5	11.2	
HW10	90	7.56	119	10.99	9.3	<0.5	<0.5	1.3	1.2	0.8	2.2	4.2	2.7	2.3	52.0	14.4	71.4	352	122.5	15.3	
HW11	100	7.56	119	10.94	9.3	0.5	<0.5	1.1	1.4	0.0	2.3	3.7	2.9	3.1	43.0	13.4	62.4	378	105.5	13.2	
HW12	110	7.50	121	10.91	9.2	<0.5	<0.5	0.9	1.8	0.0	2.3	4.1	3.7	4.6	73.7	14.3	96.3	382	106.5	12.8	
HW13	120	7.50	119	10.88	9.1	<0.5	<0.5	0.7	1.8	2.2	2.2	6.2	3.7	5.8	52.5	11.5	73.5	421	87.5	11.5	
HW14	130	7.57	120	10.85	9.0	<0.5	<0.5	0.9	1.8	2.2	2.2	6.2	3.3	4.4	38.3	12.0	58.0	354	84.5	11.6	
HW15	140	7.50	119	10.84	8.9	0.6	<0.5	0.8	1.4	0.6	2.3	4.3	3.0	4.5	43.5	13.4	64.4	428	110.5	12.9	
HW16	150	7.49	120	10.84	8.7	<0.5	<0.5	0.7	2.0	3.0	2.4	7.4	4.7	7.6	52.7	12.8	77.8	368	98.0	10.7	

Collection date 3 April 2007

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	Secchi depth = 19.0 m		DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
							VSS g m ⁻³	Chlor_a mg m ⁻³													
HW17	1	7.94	119	18.04	9.4	<0.5	<0.5	0.7	1.6	2.4	1.4	5.4	4.7	0.9	62.4	14.9	82.9	567	122.0	18.4	
HW18	10	8.09	119	18.03	9.5	<0.5	<0.5	0.8	1.1	3.9	1.8	6.8	0.0	0.1	59.9	14.9	74.9	522	317.5	19.2	
HW19	20	8.09	119	17.94	9.4	<0.5	<0.5	0.8	1.2	2.8	1.6	5.6	0.0	0.2	65.8	14.8	80.8	498	177.5	16.8	
HW20	30	7.95	119	16.72	9.3	<0.5	<0.5	1.2	1.0	4.0	2.0	7.0	0.0	0.1	63.9	17.5	81.5	481	133.0	19.6	
HW21	40	7.73	119	13.50	8.9	<0.5	<0.5	1.2	1.8	2.2	1.6	5.6	0.0	0.3	55.7	12.3	68.3	444	76.4	12.1	
HW22	50	7.62	120	12.33	8.9	<0.5	<0.5	0.8	1.5	4.5	1.3	7.3	0.1	0.8	53.2	9.0	63.1	419	68.1	10.1	
HW23	60	7.54	119	11.65	8.8	<0.5	<0.5	0.7	1.2	3.8	1.5	6.5	0.1	3.4	51.5	7.7	62.7	393	49.9	6.3	
HW24	70	7.48	120	11.28	8.8	<0.5	<0.5	0.9	2.0	2.0	1.3	5.3	0.0	9.7	70.2	6.4	86.3	434	68.3	8.6	
HW25	80	7.43	115	11.22	8.5	<0.5	<0.5	0.6	2.0	3.0	1.2	6.2	0.0	14.6	52.4	6.4	73.4	436	58.0	8.3	
HW26	90	7.39	121	11.11	8.5	<0.5	<0.5	0.3	1.7	3.3	1.0	6.0	0.1	16.3	54.7	7.1	78.2	460	62.7	8.4	
HW27	100	7.35	121	11.10	8.2	<0.5	<0.5	0.3	2.5	1.5	1.1	5.1	0.0	19.4	50.5	7.0	76.9	469	48.9	6.7	
HW28	110	7.31	121	11.04	8.2	<0.5	<0.5	0.2	2.7	2.3	0.9	5.9	1.5	20.9	47.1	5.9	75.4	437	40.4	7.5	
HW29	120	7.32	120	11.04	8.0	<0.5	<0.5	0.2	3.0	2.0	0.9	5.9	0.0	23.8	57.7	4.9	86.4	452	48.5	7.8	
HW30	130	7.73	121	11.01	8.1	<0.5	<0.5	0.2	2.7	3.3	0.9	6.9	0.0	24.8	51.2	3.8	79.8	389	42.7	6.7	
HW31	140	7.30	118	11.00	7.7	<0.5	<0.5	0.2	3.7	2.3	1.3	7.3	0.0	24.6	47.4	3.8	75.8	413	43.2	6.4	
HW32	150	7.25	121	10.99	7.4	<0.5	<0.5	0.2	4.5	3.5	1.6	9.6	0.0	30.5	50.5	6.1	87.1	439	51.7	9.5	

 NH₄, NO₃, DON, Urea all as N

* = PN by wet digestion method, ** = PN by combustion furnace method.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient database
2005-2006

Started 27 October 1994

Collection date 25 October 2005

Code	Depth	pH	EC @25oC	Temp	DO	SS	Secchi depth = 15.0 m													DOC	PC	PN**
							mS cm ⁻¹	°C	g m ⁻³	g m ⁻³	VSS	Chlor_a	DRP	DOP	PP	TP	NH ₄ -N	NO ₃ -N	DON	UREA	PN*	TN
	m						g m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	mg m ⁻³	
QD1	1	7.81	119	13.40	10.1	<0.5	<0.5	0.4	1.0	3.0	1.3	5.3	0.6	0.3	51.1	4	8.5	60.5	613	132.5	11.0	
QD2	10	7.88	119	12.88	10.0	0.5	<0.5	0.5	0.7	0.7	2.3	1.9	4.9	0.1	0.0	52.9	3	12.8	65.8	623	169.0	13.5
QD3	20	7.74	119	12.17	10.1	0.6	<0.5	0.7	0.6	0.6	2.4	2.7	5.7	0.4	0.2	43.4	2	17.0	61.0	625	216.5	20.0
QD4	30	7.77	118	11.65	9.9	0.7	<0.5	0.6	0.6	0.6	5.4	2.6	8.6	0.7	0.0	57.3	2	17.3	75.3	566	212.0	16.0
QD5	40	7.68	119	11.49	9.8	<0.5	<0.5	0.9	0.6	3.4	3.1	7.1	0.0	0.2	49.8	2	22.2	72.2	581	229.5	20.5	
QD6	50	7.59	119	11.29	9.5	<0.5	<0.5	1.4	0.8	1.2	2.2	4.2	1.4	0.1	35.5	2	15.9	52.9	599	172.5	14.0	
QD7	60	7.46	120	11.18	9.2	0.7	<0.5	0.7	0.7	1.7	2.3	1.6	5.6	1.7	9.6	41.7	2	9.8	62.8	503	103.5	6.5
QD8	70	7.37	120	11.07	9.0	0.5	<0.5	0.8	1.9	2.1	1.5	5.5	1.6	12.8	56.6	2	9.2	80.2	482	101.5	6.0	
QD9	80	7.35	120	11.01	8.8	0.6	<0.5	0.6	2.5	1.5	1.4	5.4	0.6	15.3	30.1	13	9.0	55.0	521	86.5	6.0	
QD10	90	7.36	121	10.97	8.8	0.7	<0.5	0.4	2.8	1.2	1.4	5.4	0.3	17.1	47.6	2	7.3	72.3	478	62.5	4.0	
QD11	100	7.29	121	10.97	8.6	<0.5	<0.5	0.5	2.8	1.2	1.4	5.4	0.4	17.4	39.2	2	7.8	64.8	476	77.5	4.5	
QD12	110	7.34	120	10.94	8.5	<0.5	<0.5	0.5	3.0	1.0	1.3	5.3	1.5	18.7	48.8	2	7.4	76.4	462	92.5	3.0	
QD13	120	7.29	121	10.94	8.5	<0.5	<0.5	0.5	2.8	2.2	1.2	6.2	0.8	20.4	42.8	2	6.2	70.2	549	5.0		
QD14	130	7.32	120	10.93	8.4	<0.5	<0.5	0.5	2.7	1.3	1.3	5.3	0.1	20.3	35.6	3	5.9	61.9	504	69.5	6.0	
QD15	140	7.34	121	10.93	8.4	<0.5	<0.5	0.6	3.0	2.0	1.4	6.4	1.4	20.9	34.7	1	7.8	64.8	352	77.5	6.5	
QD16	150	7.26	120	10.92	8.2	<0.5	<0.5	0.5	3.8	1.2	1.5	6.5	0.9	23.5	29.6	3	7.1	61.1	533	66.0	6.0	

Collection date 12 April 2006

Code	Depth	pH	EC @25oC	Temp	DO	Secchi depth = 15.8 m													DOC	PC	PN**
						SS	VSS	Chlor_a	DRP	DOP	PP	TP	NH ₄ -N	NO ₃ -N	DON	UREA	PN*	TN			
	m					g m ⁻³	g m ⁻³	mg m ⁻³													
ZD1	1	7.9	119	16.72	9.6	<0.5	<0.5	1.2	1.1	0.9	1.9	3.9	0.0	0.2	50.8	2	19.2	70.2	213.5	19.0	
ZD2	10	7.9	118	16.72	9.2	<0.5	<0.5	1.3	0.8	1.2	1.6	3.6	0.0	0.0	38.0	2	16.6	54.6	196.0	13.5	
ZD3	20	7.9	116	16.72	9.0	0.5	<0.5	1.1	0.7	0.3	1.3	2.3	0.0	0.0	42.0	<1	15.65	57.7	235.0	15.5	
ZD4	30	7.88	120	16.71	9.4	<0.5	<0.5	1.2	0.6	1.4	1.6	3.6	0.1	0.0	50.9	<1	15.45	66.5	172.0	13.5	
ZD5	40	7.9	116	16.64	9.2	0.8	0.7	1.3	0.5	1.5	1.55	3.6	0.0	0.0	41.0	2	15.45	56.5	224.5	13.0	
ZD6	50	7.6	119	12.11	8.7	<0.5	<0.5	1.0	0.7	2.3	1.2	4.2	0.0	0.1	33.9	8	11.4	45.4	133.0	8.5	
ZD7	60	7.43	121	11.52	8.5	<0.5	<0.5	1.0	0.7	2.3	1.05	4.1	0.0	0.5	44.5	2	9.15	54.2	171.5	8.0	
ZD8	70	7.49	121	11.31	8.3	<0.5	<0.5	0.9	0.7	2.3	1.15	4.2	0.0	0.7	37.3	6	9.55	47.6	130.5	9.0	
ZD9	80	7.9	120	11.18	8.3	<0.5	<0.5	1.1	0.5	2.5	1.4	4.4	0.3	0.0	50.7	5	16.1	67.1	182.0	12.5	
ZD10	90	7.31	122	11.11	8.1	<0.5	<0.5	0.2	3.0	1	0.45	4.5	0.0	23.0	28.0	2	4.1	55.1	62.5	6.0	
ZD11	100	7.31	122	11.08	8.1	<0.5	<0.5	0.3	3.2	0.8	0.5	4.5	0.1	22.8	24.1	<1	4.95	52.0	68.5	6.5	
ZD12	110	7.91	119	11.05	8.0	0.7	0.5	1.1	3.2	1.8	1.5	6.5	0.1	22.2	25.7	3	16.5	64.5	196.0	15.0	
ZD13	120	7.42	122	11.03	7.9	<0.5	<0.5	0.3	3.1	1.9	0.5	5.5	0.0	21.6	27.4	<1	5.2	54.2	86.5	7.0	
ZD14	130	7.5	121	11.02	7.7	<0.5	<0.5	0.3	3.0	2	0.55	5.6	0.0	19.9	32.1	2	5.45	57.5	69.5	6.5	
ZD15	140	7.3	119	11.02	7.3	<0.5	<0.5	0.2	3.4	1.6	0.55	5.6	0.0	23.1	31.9	2	6.5	61.5	87.0	7.5	
ZD16	150	7.24	122	11.02	7.2	<0.5	<0.5	0.3	2.9	1.1	0.55	4.6	0.2	21.0	28.8	5	5.85	55.9	77.5	7.0	

* = PN by wet digestion method, ** = PN by combustion furnace method.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient database
2004-2005

Started 27 October 1994

Collection date 21 October 2004

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	Secchi depth = 15.0 m													DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
							VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³				
VZ1	1	7.88	122	11.75	10.4	0.6	0.5	0.6	1.3	2.7	1.6	5.6	0.1	0.4	39.5	19	9.7	49.7	500	110.0	8	
VZ2	10	7.82	120	11.61	10.2	0.8	0.6	0.8	1.1	2.9	2.0	6.0	0.2	0.1	35.7	24	12.8	48.8	447	157.0	8.5	
VZ3	20	7.87	120	11.59	10.1	0.9	0.7	0.8	1.0	3.0	1.9	5.9	0.0	0.0	33.0	16	11.3	44.3	440	153.0	8.5	
VZ4	30	7.91	123	11.59	10.2	1.5	1.0	0.7	1.0	2.0	1.9	4.9	0.0	0.0	34.0	15	11.3	45.3	490	157.5	8	
VZ5	40	7.82	117	11.58	10.1	1.1	0.6	0.7	1.4	3.6	2.0	7.0	0.2	0.1	33.7	7	11.2	45.2	445	155.0	10	
VZ6	50	7.83	120	11.58	9.9	1.1	0.7	0.9	1.0	4.0	2.1	7.1	0.0	0.1	33.9	9	13.2	47.2	494	197.5	15	
VZ7	60	7.79	119	11.15	9.9	1.1	0.7	1.0	1.6	2.4	2.3	6.3	0.5	0.4	34.1	11	26.0	61.0	585	167.0	16	
VZ8	70	7.66	118	10.79	9.7	0.7	0.5	1.0	1.9	1.1	1.9	4.9	2.4	0.8	40.8	21	11.5	55.5	468	114.0	11.5	
VZ9	80	7.63	118	10.74	9.6	0.6	<0.5	0.9	2.0	1.0	1.7	4.7	2.8	1.3	47.9	16	8.9	60.9	440	103.0	9.5	
VZ10	90	7.61	119	10.72	9.5	0.6	<0.5	0.7	2.0	2.0	1.6	5.6	3.9	2.2	28.9	9	9.1	44.1	633	100.5	10	
VZ11	100	7.53	118	10.70	9.4	0.7	0.5	0.7	2.3	1.7	1.5	5.5	5.1	3.6	34.3	7	9.0	52.0	570	93.0	10	
VZ12	110	7.56	119	10.68	9.4	0.5	<0.5	0.7	2.0	5.0	1.6	8.6	5.3	2.8	28.9	9	9.2	46.2	514	101.5	9	
VZ13	120	7.49	119	10.66	9.3	0.5	<0.5	0.7	2.1	1.9	1.5	5.5	5.3	3.9	35.8	6	8.5	53.5	391	91.5	11	
VZ14	130	7.48	118	10.65	9.3	<0.5	<0.5	0.6	2.5	1.5	1.6	5.6	5.8	5.3	34.9	5	8.6	54.6	366	73.5	8.5	
VZ15	140	7.58	118	10.61	9.2	<0.5	<0.5	0.6	2.9	1.1	1.6	5.6	5.9	7.3	33.8	13	9.1	56.1	491	93.5	10.5	
VZ16	150	7.58	119	10.56	9.1	<0.5	<0.5	0.6	2.4	1.6	1.5	5.5	4.5	3.3	35.2	21	8.7	51.7	464	78.0	9	

Collection date 14 April 2005

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	Secchi depth = 17.2 m													DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
							VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³				
GC1	1	7.85	119	17.92	9.1	0.4	0.4	0.7	0.8	1.2	1.9	3.9	1.2	0.2	64.6	7	15.1	81.1	690	176.0	19.0	
GC2	10	7.86	118	17.96	9.0	0.3	0.4	0.9	0.8	2.2	1.9	4.9	0.0	0.0	46	3	14.1	60.1	580	199.5	19.0	
GC3	20	7.9	119	17.95	9.0	0.3	0.3	0.9	0.8	2.2	2.0	5.0	0.0	0.1	55.9	1	14.5	70.5	580	179.0	17.0	
GC4	30	7.82	118	15.13	8.4	0.3	0.3	0.9	0.8	2.2	1.8	4.8	0.0	0.3	49.7	2	12.8	62.8	570	176.5	17.0	
GC5	40	7.58	121	12.92	8.7	0.2	0.2	0.8	2.3	0.7	1.2	4.2	0.3	0.6	31.1	2	8.9	40.9	510	109.5	14.0	
GC6	50	7.51	120	12.00	8.3	0.1	0.1	0.6	3.1	0.9	1.0	5.0	0.0	6.4	39.6	3	6.8	52.8	480	84.0	9.0	
GC7	60	7.47	121	11.33	8.2	0.1	0.1	0.5	3.6	1.4	1.1	6.1	0.0	8.3	40.7	2	8.2	57.2	510	78.5	7.5	
GC8	70	7.48	120	10.99	8.2	0.1	0.1	0.3	4.2	0.8	0.9	5.9	0.0	15.7	38.3	2	6.5	60.5	490	96.0	7.0	
GC9	80	7.39	121	10.88	8.2	0.2	0.2	0.3	3.8	0.2	0.8	4.8	0.1	15.7	36.2	1	4.3	56.3	480	72.5	7.5	
GC10	90	7.21	121	10.82	8.3	0.0	0.1	0.1	5.6	1.4	0.9	7.9	0.2	23.8	38	2	5.6	67.6	480	64.0	7.0	
GC11	100	7.31	121	10.78	8.0	0.0	0.1	0.1	5.7	1.3	0.8	7.8	0.2	23.6	53.2	2	5.0	82.0	460	78.5	7.0	
GC12	110	7.32	121	10.76	7.8	0.1	0.1	0.1	5.7	1.3	0.8	7.8	0.0	25.9	47.1	2	5.6	78.6	470	43.5	6.0	
GC13	120	7.33	121	10.76	7.7	0.1	0.1	<0.1	6.4	1.6	0.8	8.8	0.3	26.8	37.9	1	4.9	69.9	450	56.0	6.5	
GC14	130	7.33	121	10.74	7.7	0.1	0.1	<0.1	6.1	0	0.8	6.8	0.3	26.7	57	1	4.4	88.4	470	43.5	5.5	
GC15	140	7.34	121	10.74	7.6	0.1	0.1	<0.1	6.6	0.4	0.9	7.9	0.2	28.8	39	2	5.8	73.8	490	54.5	6.0	
GC16	150	7.36	121	10.72	7.5	0.3	0.1	0.1	7.8	0.2	1.1	9.1	0.0	32.1	51.9	1	6.9	90.9	490	46.0	7.5	

 NH₄, NO₃, DON, Urea all as N

* = PN by wet digestion method, ** = PN by combustion furnace method.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient database
2003-2004
Started 27 October 1994
Collection date 19 November 2003

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
EU1	1	7.84	119	13.96	9.9	<0.5	<0.5	0.8	1.7	2.3	2.3	6.3	8.0	0.8	42.2	1	14.8	65.8	476	90.5	10.5
EU2	10	7.84	120	13.79	9.9	<0.5	<0.5	0.9	1.6	1.4	2.5	5.5	0.3	0.3	52.4	1	14.4	67.4	461	147.5	15.0
EU3	20	7.83	120	13.78	9.8	<0.5	<0.5	0.7	1.8	1.2	3.4	6.4	0.4	0.1	46.5	1	19.4	66.4	466	151.0	20.5
EU4	30	7.84	120	13.70	9.5	<0.5	<0.5	0.9	1.8	2.2	3.8	7.8	0.4	0.3	42.3	1	26.3	69.3	450	133.0	18.5
EU5	40	7.69	120	12.30	9.3	<0.5	<0.5	1.5	2.6	1.4	3.3	7.3	0.7	0.2	35.1	1	20.6	56.6	437	133.0	17.0
EU6	50	7.63	121	11.35	9.0	<0.5	<0.5	1.2	2.8	1.2	1.9	5.9	0.4	0.5	37.1	1	11.9	49.9	470	92.5	11.0
EU7	60	7.58	121	11.28	8.9	<0.5	<0.5	0.7	3.3	0.7	1.5	5.5	1.0	3.2	27.8	2	9.6	41.6	503	69.5	8.0
EU8	70	7.59	121	11.23	8.7	<0.5	<0.5	0.6	3.5	0.5	1.1	5.1	3.4	4.8	25.8	1	6.2	40.2	465	47.0	<6
EU9	80	7.6	121	11.19	8.6	<0.5	<0.5	0.5	3.6	0.4	1.1	5.1	0.6	5.9	29.5	2	5.1	41.1	430	65.0	<6
EU10	90	7.57	121	11.16	8.6	<0.5	<0.5	0.5	3.9	0.1	1.2	5.2	1.0	7.0	27	3	6.4	41.4	391	39.5	<6
EU11	100	7.59	121	11.15	8.6	<0.5	0.7	0.4	4.1	0.9	1.2	6.2	0.8	7.8	33.4	2	4.0	46.0	405	46.5	<6
EU12	110	7.6	121	11.12	8.4	<0.5	<0.5	0.4	4.1	0.9	1.1	6.1	1.1	11.8	29.1	3	3.4	45.4	428	45.5	<6
EU13	120	7.57	120	11.11	8.4	<0.5	<0.5	0.4	4.6	0.4	1.2	6.2	0.7	13.6	32.7	2	3.0	50.0	439	37.0	<6
EU14	130	7.53	121	11.09	8.3	<0.5	<0.5	0.3	5.1	0.4	1.2	6.7	0.8	16.1	32.7	3	3.7	53.3	408	33.0	<6
EU15	140	7.57	121	11.09	8.2	<0.5	<0.5	0.3	5.3	0.7	1.2	7.2	0.4	18.1	32.5	3	5.1	56.1	440	54.5	<6
EU16	150	7.54	120	11.09	8.0	0.5	<0.5	0.5	5.6	1.4	1.5	8.5	2.4	20.7	32.9	4	6.4	62.4	481	44.0	<6

Collection date 31 March 2004

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
MB1	1	7.86	118	16.49	9.2	<0.5	<0.5	0.7	0.9	4.1	1.4	6.4	1	0	69	-	9.7	79.7	622	91.0	-
MB2	10	7.83	118	16.29	9.1	<0.5	<0.5	1.2	0.5	3.5	2.0	6.0	0	0	47	-	12.4	59.4	548	141.5	17.0
MB3	20	7.83	118	16.23	9.0	<0.5	<0.5	1.1	0.6	3.4	2.1	6.1	1	0.2	47.8	-	14.8	63.8	561	140.5	17.0
MB4	30	7.83	118	16.19	9.0	<0.5	<0.5	1.1	0.8	3.2	1.9	5.9	1	0.2	50.8	-	13.5	65.5	749	131.5	15.5
MB5	40	7.66	118	16.15	8.9	<0.5	<0.5	0.9	1.5	1.5	1.9	4.9	1	2.8	71.2	-	11.6	86.6	560	114.5	14.0
MB6	50	7.46	120	12.51	8.2	<0.5	<0.5	0.5	3.3	2.7	1.5	7.5	1	12.1	58.9	-	7.2	79.2	467	109.0	7.5
MB7	60	7.41	121	11.59	8.0	<0.5	<0.5	0.3	4.7	2.3	1.0	8.0	1	18.0	41	-	4.2	64.2	394	54.5	7.0
MB8	70	7.36	121	11.40	8.0	<0.5	<0.5	0.2	4.5	1.5	0.8	6.8	1	19.1	36.9	-	3.7	60.7	404	45.0	<4
MB9	80	7.42	121	11.34	8.0	<0.5	<0.5	0.2	5.0	1.0	0.8	6.8	1	20.2	31.8	-	5.3	58.3	464	41.0	<4
MB10	90	7.36	121	11.30	7.9	<0.5	<0.5	0.1	5.2	1.8	0.7	7.7	3	22.1	35.9	-	3.9	64.9	453	52.0	<4
MB11	100	7.31	122	11.27	7.8	<0.5	<0.5	0.1	5.6	2.4	0.8	8.8	2	23.9	38.1	-	3.0	67.0	477	36.5	<4
MB12	110	7.29	122	11.26	7.7	<0.5	<0.5	<0.1	5.8	2.2	1.0	9.0	1	25.0	30	-	6.2	62.2	392	36.5	5.5
MB13	120	7.31	121	11.24	7.6	<0.5	<0.5	0.1	5.9	3.1	0.8	9.8	1	25.0	59	-	3.6	88.6	373	53.5	<4
MB14	130	7.3	121	11.22	7.5	<0.5	<0.5	<0.1	6.3	2.7	0.9	9.9	0	27.0	35	-	3.3	65.3	393	61.0	<4
MB15	140	7.3	121	11.21	7.4	<0.5	<0.5	<0.1	6.6	3.4	0.8	10.8	0	27.8	46.2	-	3.3	77.3	356	35.0	<4
MB16	150	7.31	120	11.21	7.1	<0.5	<0.5	0.1	7.2	2.8	1.0	11.0	0	30.1	48.9	-	4.0	83.0	394	34.0	<4

 NH₄, NO₃, DON, Urea all as N

* = PN by wet digestion method, ** = PN by combustion furnace method.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient
Collection date 13 November 2002

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	Secchi depth = 18.0 m														Started 27 October 1994
							VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
NZ1	1	7.87	122	12.58	10.2	0.6	<0.5	0.6	1.3	1.7	2.2	5.2	0.8	0.6	65.6	2	15.3	82.3	620	160.0	12.5
NZ2	10	7.86	120	12.58	10.3	0.5	<0.5	0.7	1.2	1.8	2.1	5.1	0.7	0.0	49.3	1	13.7	63.7	573	180.5	13.5
NZ3	20	7.93	120	12.49	10.2	1.0	<0.5	0.7	1.1	1.9	2.2	5.2	0.5	0.1	61.4	1	15.8	77.8	536	157.5	12.0
NZ4	30	7.85	121	12.38	10.2	<0.5	<0.5	0.8	0.9	3.1	2.6	6.6	0.7	0.5	74.8	2	17.7	93.7	657	242.0	14.0
NZ5	40	7.81	119	12.16	10.1	<0.5	<0.5	0.7	1.2	1.8	1.9	4.9	0.6	0.7	58.7	1	12.9	72.9	506	164.5	8.0
NZ6	50	7.83	120	12.00	10.1	<0.5	<0.5	0.7	1.6	1.4	1.7	4.7	1.6	0.0	55.4	1	11.5	68.5	505	170.0	9.5
NZ7	60	7.78	119	11.81	10.0	<0.5	<0.5	0.6	1.5	1.5	1.5	4.5	1.2	0.0	64.8	2	9.5	75.5	531	108.5	6.5
NZ8	70	7.72	120	11.51	9.9	<0.5	<0.5	0.6	2.8	1.2	1.3	5.3	3.4	2.2	42.4	7	7.1	55.1	514	53.5	5.0
NZ9	80	7.67	120	11.32	9.7	<0.5	<0.5	0.4	2.7	1.3	1.1	5.1	3.3	0.9	38.8	2	5.9	48.9	578	61.0	4.5
NZ10	90	7.77	121	11.13	9.6	<0.5	<0.5	0.4	2.8	1.2	1.0	5.0	3.7	0.4	44.9	4	6.6	55.6	487	41.0	<2
NZ11	100	7.53	122	11.08	9.4	<0.5	<0.5	0.2	3.0	2.0	0.8	5.8	4.2	3.7	65.1	5	6.1	79.1	525	31.0	<2
NZ12	110	7.64	121	11.05	9.4	<0.5	<0.5	0.1	3.3	1.7	0.7	5.7	3.4	5.4	57.2	4	4.4	70.4	472	38.0	<2
NZ13	120	7.55	122	11.01	9.3	<0.5	<0.5	0.2	3.6	0.4	1.0	5.0	3.0	7.0	51.0	6	5.9	66.9	473	64.5	4.0
NZ14	130	7.32	123	10.99	9.2	<0.5	<0.5	0.1	3.6	0.4	1.0	5.0	2.9	7.5	45.6	5	6.7	62.7	555	70.5	3.5
NZ15	140	7.47	121	10.97	9.1	0.5	<0.5	0.1	3.7	1.3	0.9	5.9	2.5	10.5	60.0	16	6.7	79.7	460	54.5	3.0
NZ16	150	7.46	121	10.96	9.0	<0.5	<0.5	0.2	4.3	1.7	1.0	7.0	0.5	12.9	58.6	4	6.4	78.4	461	52.5	3.0

Collection date 3 April 2003

Code	Depth m	pH	EC @25oC mS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	Secchi depth = 13.5 m														Started 27 October 1994
							VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
UJ1	1	8.01	119	19.20	8.8	3.0	0.5	0.7	0.8	3.2	1.8	5.8	5	0.4	75.6	5	18.8	99.8	546	219.0	19.5
UJ2	10	8.07	146	18.71	8.8	0.7	1.0	1.4	0.9	4.1	2.5	7.5	<1	0.6	45.4	1	24.0	70.0	511	304.5	29.0
UJ3	20	8.15	120	18.60	8.6	1.0	0.7	1.3	0.6	3.4	2.3	6.3	<1	0.6	40.4	1	23.7	64.7	520	270.0	31.5
UJ4	30	7.93	119	16.93	8.3	<0.5	<0.5	1.5	0.8	3.2	1.8	5.8	<1	0.3	39.7	1	20.4	60.4	503	181.0	39.0
UJ5	40	7.66	118	13.31	8.0	<0.5	<0.5	1.3	1.7	3.3	1.7	6.7	<1	0.8	39.2	1	12.2	52.2	443	115.0	54.0
UJ6	50	7.61	122	12.39	7.9	<0.5	1.0	0.7	2.9	2.1	1.3	6.3	<1	4.8	35.2	3	8.6	48.6	410	92.5	5.5
UJ7	60	7.57	138	11.80	7.7	<0.5	<0.5	0.5	3.9	2.1	1.1	7.1	<1	10.7	32.3	1	5.9	48.9	366	86.5	4.5
UJ8	70	7.42	121	11.50	7.6	<0.5	<0.5	0.2	4.4	1.6	0.9	6.9	<1	16.3	27.7	1	6.1	50.1	404	109.5	4.0
UJ9	80	7.39	121	11.32	7.5	<0.5	<0.5	0.1	4.5	1.5	1.0	7.0	<1	19.3	41.7	1	6.2	67.2	365	37.0	4.0
UJ10	90	7.32	121	11.20	7.3	<0.5	<0.5	0.1	4.7	1.3	0.8	6.8	<1	21.9	24.1	2	4.5	50.5	360	40.0	<4
UJ11	100	7.29	121	11.19	7.3	<0.5	<0.5	<0.1	5.3	2.7	0.9	8.9	<1	23.9	27.1	2	4.6	55.6	387	92.5	<4
UJ12	110	7.26	120	11.12	7.2	<0.5	<0.5	<0.1	5.5	0.5	0.7	6.7	<1	25.2	30.8	1	2.9	58.9	366	28.5	<4
UJ13	120	7.33	122	11.11	7.0	<0.5	<0.5	<0.1	6.6	0.4	0.7	7.7	<1	28.8	36.2	5	2.5	67.5	409	40.0	<4
UJ14	130	7.27	123	11.09	6.9	<0.5	<0.5	<0.1	7.7	0.3	0.9	8.9	<1	30.9	29.1	3	3.2	63.2	382	15.5	<4
UJ15	140	7.28	122	11.10	6.8	<0.5	<0.5	<0.1	7.6	0.4	0.8	8.8	<1	30.4	47.6	4	4.3	82.3	384	47.5	<4
UJ16	150	7.29	122	11.09	6.5	<0.5	<0.5	<0.1	9.0	5.0	1.6	15.6	<1	36.4	30.6	2	6.5	73.5	371	38.5	<4

* = PN by wet digestion method, ** = PN by combustion furnace method.

Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

FIA instrument results are given as a better indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N below nominal detection limit.

Lake Taupo biannual nutrient database
Collection date 12 November 2001

Code	Depth	pH	EC @25oC µS cm ⁻¹	Temp °C	2001-2002										Started 27 October 1994						
					DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH4-N mg m ⁻³	NO3-N mg m ⁻³	DON mg m ⁻³	UREA	PN*	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
XH1	1	7.85	122	14.23	9.5	0.5	<0.5	0.6	0.9	1.1	1.55	3.6	<1	<0.5	29	2	6	35	500	146.5	12.0
XH2	10	7.86	122	14.16	9.8	0.5	<0.5	0.7	1.1	0.9	4.3	6.3	<1	<0.5	32	2	16.5	49	520	212.0	31.3
XH3	20	7.82	119	13.37	9.4	<0.5	<0.5	1.0	1.1	<0.5	3.5	4.6	<1	<0.5	28	1	20	48	510	340.5	26.8
XH4	30	7.6	116	12.85	9.4	0.6	0.7	1.3	1.6	<0.5	3.1	4.7	<1	1.0	29	1	14.5	45	480	264.5	24.7
XH5	40	7.44	122	11.87	8.9	<0.5	<0.5	1.3	2.2	<0.5	2.8	5.0	1	2.5	25.5	2	11.5	41	470	200.5	21.7
XH6	50	7.46	121	11.57	9.0	<0.5	<0.5	0.9	2.6	<0.5	1.75	4.4	<1	7.2	26.8	2	6	40	470	136.5	12.6
XH7	60	7.41	121	11.24	8.7	1.3	1.2	0.7	2.6	<0.5	1.4	4.0	<1	8.0	24	2	<2	32	440	104.5	9.1
XH8	70	7.4	122	11.13	8.8	<0.5	<0.5	0.5	2.9	<0.5	1.15	4.1	<1	12.3	21.7	2	<2	34	450	142.0	7.2
XH9	80	7.38	122	11.03	8.6	<0.5	<0.5	0.4	3.2	<0.5	1.15	4.4	<1	13.6	29.4	4	<2	43	440	103.0	8.1
XH10	90	7.4	119	11.01	8.8	<0.5	<0.5	0.4	3.2	<0.5	1.05	4.3	<1	15.1	21.9	2	<2	37	420	79.0	6.2
XH11	100	7.35	120	10.99	8.6	<0.5	<0.5	0.3	3.8	<0.5	1.05	4.9	<1	17.8	25.2	2	4	47	460	98.0	6.6
XH12	110	7.36	122	10.97	8.6	<0.5	<0.5	0.3	4.0	<0.5	1.1	5.1	<1	19.5	24.5	2	<2	44	490	116.5	5.8
XH13	120	7.35	126	10.95	8.4	<0.5	<0.5	0.3	4.5	<0.5	1.3	5.8	<1	22.0	22	2	<2	44	490	93.5	5.6
XH14	130	7.38	127	10.94	8.4	<0.5	<0.5	0.3	4.4	<0.5	1.1	5.5	<1	21.1	21.9	2	<2	43	420	113.5	5.5
XH15	140	7.34	126	10.94	8.2	<0.5	<0.5	0.3	5.2	<0.5	1.3	6.5	<1	24.7	25.3	2	<2	50	440	93.5	7.3
XH16	150	7.38	127	10.94	8.1	1.3	0.6	0.3	5.3	<0.5	1.3	6.6	<1	25.2	26.8	3	<2	52	480	83.5	7.7

Collection date 4 April 2002

Code	Depth	pH	EC @25oC µS cm ⁻¹	Temp °C	Secchi depth = 19.0 m										Secchi depth = 15.5 m						
					DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH4-N mg m ⁻³	NO3-N mg m ⁻³	DON mg m ⁻³	UREA	PN*	TN mg m ⁻³	DOC g m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
EJ1	1	7.91	119	17.45	8.8	<0.5	<0.5	0.72	0.5	0.5	1	2.0	1.1	0.3	44.6	7.85	53.9	0.5	187.0	10.0	
EJ2	10	7.94	118	17.38	8.9	<0.5	<0.5	0.96	0.6	1.4	1.4	3.4	0.2	0.1	44.7	9.4	54.4	0.6	164.5	10.5	
EJ3	20	7.88	119	17.18	8.8	<0.5	<0.5	1.02	0.5	1.5	1.35	3.4	0.3	0.0	38.7	9.45	48.5	0.8	154.5	11.0	
EJ4	30	7.85	119	16.83	8.7	<0.5	<0.5	0.95	0.7	2.3	1.45	4.5	0.4	0.1	40.5	8.4	49.4	0.5	136.5	10.5	
EJ5	40	7.65	121	12.9	8.3	<0.5	<0.5	0.89	1.4	0.6	1.2	3.2	0.4	0.8	32.8	7.95	42.0	0.4	100.0	8.0	
EJ6	50	7.66	120	12.09	8.2	<0.5	<0.5	0.85	2.1	0.9	1.3	4.3	0.4	3.5	35.1	7.8	46.8	0.4	114.0	9.0	
EJ7	60	7.60	123	11.51	8.1	<0.5	<0.5	0.50	3.9	2.1	1	7.0	0.9	12.3	30.8	5.7	49.7	0.4	75.0	6.0	
EJ8	70	7.42	123	11.3	8.0	<0.5	<0.5	0.26	4.5	0.5	0.95	6.0	0.0	20.9	30.1	5.65	56.7	0.5	49.5	4.0	
EJ9	80	7.46	121	11.24	7.9	<0.5	<0.5	0.24	4.6	0.4	1.1	6.1	0.2	24.8	29	7.55	61.6	0.3	50.0	5.0	
EJ10	90	7.38	121	11.19	7.8	<0.5	<0.5	0.19	5.3	<0.5	0.75	6.1	0.3	28.1	23.6	4.45	56.5	0.4	48.0	4.0	
EJ11	100	7.33	121	11.17	7.8	<0.5	<0.5	0.11	5.4	0.6	0.8	6.8	0.1	28.6	30.3	5.05	64.1	0.3	76.0	5.5	
EJ12	110	7.37	122	11.14	7.7	<0.5	<0.5	0.10	6.0	<0.5	0.8	6.8	0.5	31.7	23.8	6.15	62.2	0.6	67.5	7.5	
EJ13	120	7.36	122	11.14	7.7	<0.5	<0.5	0.10	6.3	<0.5	0.6	6.9	0.2	32.2	24.6	3.25	60.3	0.3	46.5	4.0	
EJ14	130	7.32	122	11.13	7.6	<0.5	<0.5	0.09	6.5	<0.5	0.45	7.0	0.1	32.2	26.7	0.8	59.8	0.5	48.0	5.5	
EJ15	140	7.34	122	11.13	7.1	<0.5	<0.5	0.07	7.0	<0.5	0.7	7.7	1.1	34.0	29.9	4.9	69.9	0.4	44.0	4.0	
EJ16	150	7.44	122	11.13	7.0	<0.5	<0.5	0.09	8.7	<0.5	0.9	9.6	0.8	36.3	24.9	4.45	66.5	0.4	75.5	4.0	

 NH₄, NO₃, DON, Urea all as N * = PN by wet digestion method, ** = PN by combustion furnace method.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

New Analytical instrument (Flow Injection Analysis) from January 2002, gives greatly improved resolution at low levels.

 FIA instrument results are given for Autumn as an indication of likely absolute low levels of DRP, NO₃-N, and NH₄-N.

Lake Taupo biannual nutrient database
2000-2001
Started 27 October 1994
Collection date 26 October 2000

Code	Depth m	pH	EC @25°C μS cm⁻¹	Temp °C	DO g m⁻³	SS g m⁻³	VSS g m⁻³	Chlor_a mg m⁻³	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH4-N mg m⁻³	NO3-N mg m⁻³	DON mg m⁻³	UREA mg m⁻³	PN*	TN mg m⁻³	DOC g m⁻³	PC mg m⁻³	PN** mg m⁻³
FX1	1	7.87	120	12.5	9.1	0.5	<0.5	0.4	<1	3	2	5.0	1	<1	25	4	9	35	0.5	104.5	4.0
FX2	10	7.85	120	11.5	8.7	0.8	0.5	1.1	1	4	3	8.0	<1	<1	33	2	23	56	0.5	196.0	12.0
FX3	20	7.79	120	11.4	8.7	<0.5	<0.5	1.3	<1	2	4	6.0	<1	<1	41	2	29	70	0.5	237.0	19.0
FX4	30	7.74	120	11.3	8.7	1.1	0.5	1.3	<1	2	3	5.0	<1	<1	36	1	24	60	0.5	183.0	11.0
FX5	40	7.69	119	11.3	9.1	0.9	0.5	1.5	<1	2	3	5.0	1	<1	38	2	18	57	0.5	90.5	7.0
FX6	50	7.63	120	11.3	9.1	0.8	<0.5	1.4	1	2	2	5.0	2	<1	64	2	14	80	0.4	79.5	6.0
FX7	60	7.54	120	11.3	8.7	0.9	<0.5	1.2	1	1	2	4.0	<1	<1	45	2	14	59	0.4	58.0	5.0
FX8	70	7.52	120	11.2	8.7	<0.5	<0.5	1.2	1	1	2	4.0	4	1	38	4	14	57	0.5	61.5	5.0
FX9	80	7.52	120	11.2	8.7	0.9	<0.5	1.1	2	2	2.5	6.5	5	2	44	2	13	64	0.5	44.5	<4
FX10	90	7.59	120	11.2	8.7	0.9	<0.5	1.1	2	2	2	6.0	6	3	37	2	14	60	0.5	58.5	5.5
FX11	100	7.47	120	11.1	8.7	<0.5	<0.5	1.4	1	1	3	5.0	3	4	39	4	16	62	0.4	48.5	6.0
FX12	110	7.41	121	11.1	8.7	0.9	<0.5	1.2	2	2	3	7.0	3	4	38	3	15	60	0.4	29.5	<4
FX13	120	7.40	121	11.0	8.2	0.5	<0.5	0.8	2	2	2	6.0	6	7	38	5	8	59	0.4	104.0	5.5
FX14	130	7.42	121	11.0	8.5	0.6	<0.5	0.2	2	2	2	6.0	6	7	41	4	11	65	0.4	71.0	6.5
FX15	140	7.36	121	11.0	8.6	0.8	<0.5	0.6	4	1	3	8.0	5	11	40	3	11	67	0.4	65.5	5.0
FX16	150	7.32	121	11.0	8.5	0.6	<0.5	1.4	4	2	4	10.0	8	13	47	9	18	86	0.4	110.5	8.0

Collection date 8 April 2001

Code	Depth m	pH	EC @25°C μS cm⁻¹	Temp °C	DO g m⁻³	SS g m⁻³	VSS g m⁻³	Chlor_a mg m⁻³	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH4-N mg m⁻³	NO3-N mg m⁻³	DON mg m⁻³	UREA mg m⁻³	PN*	TN mg m⁻³	DOC g m⁻³	PC mg m⁻³	PN** mg m⁻³
NZ1	1	7.94	120	17.0	8.3	<0.5	<0.5	1.0	<1	2	2	4.0	2	1	40	7	20.0	63.0	0.6	201.0	15.5
NZ2	10	7.97	120	16.9	8.3	<0.5	<0.5	1.4	<1	1	2	3.0	<1	<1	29	1	19.0	48.0	0.6	189.0	13.0
NZ3	20	7.99	120	16.8	8.4	<0.5	<0.5	1.5	<1	1	2	3.0	<1	<1	36	1	19.0	55.0	0.6	208.5	14.5
NZ4	30	7.96	124	15.8	8.0	<0.5	<0.5	1.2	<1	2	2	4.0	1	<1	42	1	16.0	59.0	0.6	156.0	10.5
NZ5	40	7.76	120	13.1	7.8	<0.5	<0.5	1.2	<1	1	1.5	2.5	1	1	22	2	12.0	36.0	0.5	145.0	8.5
NZ6	50	7.69	119	12.4	7.5	<0.5	<0.5	1.0	2	0	1	3.0	1	2	22	2	10.0	35.0	0.5	100.0	5.5
NZ7	60	7.60	120	11.8	7.2	<0.5	<0.5	0.8	1	1	1	3.0	<1	9	16	2	7.0	32.0	0.5	82.0	<2
NZ8	70	7.57	120	11.7	7.1	<0.5	<0.5	0.4	3	0	<1	3.0	<1	19	25	2	5.5	49.5	0.4	80.5	<2
NZ9	80	7.44	121	11.5	6.9	<0.5	<0.5	0.3	3	0	<1	3.0	2	24	15	3	5.0	46.0	0.6	70.0	<2
NZ10	90	7.39	121	11.5	6.9	<0.5	<0.5	0.2	3	1	<1	4.0	2	26	14	4	4.0	46.0	0.5	57.5	<2
NZ11	100	7.38	122	11.4	6.8	<0.5	<0.5	0.2	4	0	<1	4.0	2	29	16	1	4.0	51.0	0.5	47.5	<2
NZ12	110	7.39	122	11.4	6.8	<0.5	<0.5	0.1	4	1	<1	4.0	2	31	18	4	3.5	54.5	0.5	42.5	<2
NZ13	120	7.41	121	11.3	6.7	<0.5	<0.5	0.1	5	0	<1	5.0	1	33	16	4	5.0	55.0	0.4	40.0	<2
NZ14	130	7.42	122	11.3	6.6	<0.5	<0.5	0.1	5	0	<1	5.0	1	33	20	4	5.0	59.0	0.5	42.5	<2
NZ15	140	7.34	123	11.3	6.4	<0.5	<0.5	0.1	6	1	<1	7.0	2	38	12	5	4.5	56.5	0.5	55.0	<2
NZ16	146	7.30	123	11.3	6.3	<0.5	<0.5	0.1	7	2	1	10.0	2	43	22	5	6.5	73.5	0.5	70.5	<2

 NH₄, NO₃, DON, Urea all as N

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

* = PN by wet digestion method, ** = PN by combustion furnace method.

Lake Taupo biannual nutrient database
Collection date 18 October 1999

Secchi depth = 14.9 m

1999-2000

Started 27 October 1994

Code	Depth m	pH	EC @25oC µS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a ⁺⁺ mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
PX1	1	7.71	119	12.8	8.9	0.5	<0.5	0.14	0.5	3	3.7	7.2	<1	<1	41	16	19.4	60.4	441	105.7	8.8
PX2	10	7.74	117	12.7	8.9	<0.5	<0.5	0.39	0.5	4	3.2	7.7	<1	<1	36	4	19.9	55.9	411	160.8	12.9
PX3	20	7.73	122	12.4	8.9	0.6	<0.5	0.80	1	2	5.5	8.5	<1	<1	34	1	37.8	71.8	437	254.7	37.3
PX4	30	7.76	120	11.6	8.9	<0.5	1.9	1.06	1	2	3.9	6.9	<1	<1	36	<1	26.7	62.7	413	198.3	24.2
PX5	40	7.57	117	11.4	8.8	<0.5	<0.5	3.14	2	2	2.4	6.4	5	<1	44	22	14.6	63.6	392	117.2	9.7
PX6	50	7.48	119	11.3	8.6	<0.5	<0.5	2.90	2.5	2	1.7	6.2	8	2	33	5	9.1	52.1	417	87.0	6.6
PX7	60	7.49	118	11.1	8.6	0.5	<0.5	1.45	3	1	1.5	5.5	7	9	36	5	12.6	64.6	449	95.0	11.1
PX8	70	7.41	117	11.1	8.6	<0.5	<0.5	0.65	3.5	1	1.5	6.0	4	15	27	9	5.6	51.6	421	49.9	4.9
PX9	80	7.39	117	11.0	8.5	<0.5	<0.5	0.75	3.5	2	1.4	6.9	4	17	31	7	5.7	57.7	398	42.7	5.7
PX10	90	7.36	118	11.0	8.6	<0.5	<0.5	0.54	4	2	1.3	7.3	3	17	29	2	5.8	54.8	393	51.2	5.7
PX11	100	7.36	118	11.0	8.6	<0.5	<0.5	0.63	4	1	1.6	6.6	4	18	30	2	7.3	59.3	492	56.1	5.8
PX12	110	7.35	118	11.0	8.6	0.5	<0.5	0.65	4	2	1.8	7.8	5	18	46	10	20.1	89.1	547	129.5	21.4
PX13	120	7.33	119	11.0	8.3	0.8	0.7	0.71	4	2	1.7	7.7	6	19	47	20	45.3	117.3	530	222.3	44.3
PX14	130	7.33	119	11.0	7.9	0.6	0.5	0.59	4	2	1.7	7.7	5	19	40	12	15.3	79.3	461	112.9	19.7
PX15	140	7.32	123	11.0	7.5	0.6	<0.5	0.90	4	1	2.3	7.3	4	19	53	12	16.5	92.5	514	84.5	9.7
PX16	150	7.29	119	11.0	7.5	1.6	<0.5	0.67	4.5	2	2.1	8.6	3	19	34	7	9.6	65.6	783	63.9	6.8

Collection date 12 April 2000

Secchi depth = 15 m

Code	Depth m	pH	EC @25oC µS cm ⁻¹	Temp °C	DO g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ -N mg m ⁻³	NO ₃ -N mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³
YX1	1	7.86	118	17.4	9.2	0.6		1.3	<1	4	2	6.0	6	2	72	8	16	96.0	542	255.0	31.0
YX2	10	7.88	118	17.3	9.2	1.1		1.3	<1	3	2	5.0	3	1	57	1	21	82.0	472	198.5	16.5
YX3	20	7.88	118	17.2	9.2	1.0		1.4	<1	3	2	5.0	1	<1	59	3	15.5	75.5	599	166.5	12.0
YX4	30	7.79	118	16.7	9.0	1.1		1.3	<1	3	2	5.0	1	<1	59	2	17	77.0	608	154.0	17.5
YX5	40	7.29	119	12.6	8.3	0.6		1.1	2	2	1	5.0	2	2	57	6	9.5	70.5	396	72.0	6.0
YX6	50	7.17	120	11.7	8.0	1.0		0.8	3	2	1	6.0	2	7	42	7	8.5	59.5	403	94.5	7.5
YX7	60	7.18	119	11.4	8.0	0.5		1.0	4	1	<1	5.0	1	16	44	1	4	65.0	402	48.5	<4
YX8	70	7.1	120	11.3	8.0	0.6		<0.1	6	1	<1	7.0	6	29	35	1	6.5	76.5	418	41.0	4.0
YX9	80	7.14	120	11.2	7.9	1.0		<0.1	6	1	<1	7.0	2	32	46	1	12	92.0	451	105.5	8.0
YX10	90	7.11	120	11.2	7.9	0.7		<0.1	7	<1	<1	7.0	1	35	34	2	11	81.0	428	67.5	5.0
YX11	100	7.12	125	11.2	7.7	0.7		<0.1	7	2	<1	9.0	2	37	41	1	8.5	88.5	417	68.5	<4
YX12	110	7.12	120	11.2	7.7	0.9		<0.1	7	2	<1	9.0	2	37	50	3	11	100.0	439	65.0	5.5
YX13	120	7.06	120	11.1	7.7	0.6		<0.1	8	1	<1	9.0	3	39	47	1	6.5	95.5	431	40.5	0.0
YX14	130	7.12	120	11.1	7.5	1.2		<0.1	8	1	<1	9.0	2	40	47	3	9	98.0	453	57.0	5.0
YX15	140	7.08	120	11.1	7.5	1.2		<0.1	9	<1	<1	9.0	2	42	45	2	8	97.0	415	50.5	<4
YX16	146	7.04	120	11.1	7.2	1.7		0.1	10	3	1	14.0	4	43	42	2	10	99.0	429	92.0	4.0

NH₄, NO₃, DON, Urea all as N

* = PN by wet digestion method, ** = PN by combustion furnace method.

Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

++ = from calibrated chlorophyll fluorescence profiler (filters damaged)

Lake Taupo biannual nutrient database
Collection date 1 November 1998

Secchi depth =

13.5 m

1998-1999
Started 27 October 1994

Code	Depth m	pH	EC @25oC $\mu\text{S cm}^{-1}$	Temp °C	DO g m^{-3}	SS g m^{-3}	VSS g m^{-3}	Chlor_a mg m^{-3}	DRP mg m^{-3}	DOP mg m^{-3}	PP mg m^{-3}	TP mg m^{-3}	NH4-N mg m^{-3}	NO3-N mg m^{-3}	DON mg m^{-3}	PN* mg m^{-3}	TN mg m^{-3}	DOC g m^{-3}	PC mg m^{-3}	PN** mg m^{-3}
DM1	1	7.91	118	13.6	10.4	0.8	<0.5	0.8	0.7	1.5	2.0	4.2	3.4	<0.5	35	10.8	49.2	133.5	12.0	
DM2	10	7.87	117	13.2	10.7	0.8	<0.5	1.0	0.6	1.3	2.6	4.5	2.4	<0.5	36	15.2	53.6	180.5	15.0	
DM3	20	7.82	118	12.7	10.7	0.5	<0.5	1.4	0.6	1.4	2.9	4.9	1.9	1.1	37	18.0	58.0	215.0	23.3	
DM4	30	7.80	118	12.4	10.6	<0.5	<0.5	1.1	0.5	1.3	2.3	4.1	1.9	<0.5	34	14.1	50.0	128.0	13.5	
DM5	40	7.75	118	12.4	10.4	<0.5	<0.5	0.6	0.6	1.2	1.7	3.5	2.5	<0.5	34	9.2	45.7	118.0	10.4	
DM6	50	7.70	118	12.2	10.2	<0.5	<0.5	0.6	0.6	1.2	1.7	3.5	2.6	0.6	31	8.1	42.3	114.5	7.9	
DM7	60	7.46	119	11.7	10.0	<0.5	<0.5	0.4	2.1	1.0	1.4	4.5	1.6	9.5	32	6.0	49.1	73.0	6.0	
DM8	70	7.30	120	11.2	9.6	<0.5	<0.5	0.3	3.3	0.9	1.0	5.2	2.7	16.0	32	3.8	54.5	56.0	2.7	
DM9	80	7.15	121	11.1	9.1	<0.5	<0.5	0.2	3.9	0.8	0.9	5.6	1.5	20.5	29	5.0	56.0	64.5	2.7	
DM10	90	7.07	122	11.1	8.8	<0.5	<0.5	0.2	4.9	0.5	0.9	6.3	2.6	24.8	32	5.0	64.4	45.0	2.9	
DM11	100	7.16	121	11.0	8.5	<0.5	<0.5	0.2	5.0	0.5	0.9	6.4	3.3	26.2	34	3.6	67.1	42.5	2.0	
DM12	110	7.16	122	11.0	8.3	<0.5	<0.5	0.1	6.2	0.4	0.8	7.4	2.0	29.2	30	4.0	65.2	54.0	2.9	
DM13	120	7.11	122	11.0	8.0	<0.5	<0.5	0.1	6.4	0.3	0.8	7.5	2.2	30.6	29	3.3	65.1	63.0	1.8	
DM14	130	7.08	122	11.0	7.8	<0.5	<0.5	0.1	7.0	0.2	0.8	8.0	2.2	31.4	28	3.1	64.7	48.5	2.0	
DM15	140	7.07	123	10.9	7.6	<0.5	<0.5	0.1	7.9	0.0	0.9	8.8	2.0	33.8	32	5.0	72.8	54.0	2.0	
DM16	150	7.10	123	10.9	7.6	2.5	<0.5	0.2	8.2	0.4	3.7	12.3	2.7	35.4	34	12.8	84.9	140.5	10.5	

Collection date 14 April
1999

Secchi depth =

13 m

Code	Depth m	pH	EC @25oC $\mu\text{S cm}^{-1}$	Temp °C	DO g m^{-3}	SS g m^{-3}	VSS g m^{-3}	Chlor_a mg m^{-3}	DRP mg m^{-3}	DOP mg m^{-3}	PP mg m^{-3}	TP mg m^{-3}	NH4-N mg m^{-3}	NO3-N mg m^{-3}	DON mg m^{-3}	PN* mg m^{-3}	TN mg m^{-3}	DOC g m^{-3}	PC mg m^{-3}	PN** mg m^{-3}
II1	1	119	18.3	8.9	<0.5	<0.5	1.2	0.6	1.8	2.4	3	<0.5	43	19.0	65.0	0.6	221.4	19.5		
II2	10	118	18.3	8.8	<0.5	<0.5	1.2	0.5	1.8	2.3	1	<0.5	40	19.3	60.3	0.5	216.3	17.6		
II3	20	118	18.3	8.8	<0.5	<0.5	1.2	0.5	1.7	2.2	1	2	41	19.0	63.0	0.5	132.3	8.9		
II4	30	118	18.1	8.7	<0.5	<0.5	1.2	1.1	1.4	2.5	1	3	34	14.0	52.0	0.6	136.8	9.7		
II5	40	118	12.9	8.4	<0.5	<0.5	0.7	2.3	0.9	3.2	1	6	31	8.9	46.9	0.7	91.2	6.5		
II6	50	119	11.9	8.1	<0.5	<0.5	0.4	3.1	0.7	3.8	1	14	28	7.9	50.9	0.5	63.1	4.8		
II7	60	121	11.6	8.0	<0.5	<0.5	0.3	4.3	0.7	5.0	1	19	33	7.3	60.3	0.6	42.3	5.0		
II8	70	121	11.4	8.0	<0.5	<0.5	0.2	5.5	0.8	6.3	1	23	27	8.6	59.6	0.4	48.4	7.0		
II9	80	122	11.3	7.8	<0.5	<0.5	0.1	5.9	0.8	6.7	2	28	29	8.3	67.3	0.5	51.5	6.1		
II10	90	123	11.2	7.6	<0.5	<0.5	0.1	6.1	0.6	6.7	1	30	31	6.4	68.4	0.5	62.1	4.2		
II11	100	122	11.2	7.4	<0.5	<0.5	0.1	6.1	0.5	6.6	2	27	28	6.1	63.1	0.6	33.1	1.5		
II12	110	120	11.2	7.2	<0.5	<0.5	0.1	6.6	0.5	7.1	2	28	27	6.1	63.1	0.5	35.7	2.9		
II13	120	122	11.2	7.1	<0.5	<0.5	0.1	6.4	0.5	6.9	2	24	26	5.2	57.2	0.6	34.1	2.2		
II14	130	122	11.1	6.8	<0.5	<0.5	<0.1	7.5	0.5	8.0	2	28	31	6.3	67.3	0.6	46.9	5.5		
II15	140	122	11.1	6.3	<0.5	<0.5	0.1	8.8	0.9	9.7	2	33	31	6.4	72.4	0.5	63.4	3.0		
II16	150	116	11.1	5.9	<0.5	<0.5	<0.1	8.6	0.9	9.5	4	28	60	7.7	99.7	0.9	51.1	1.1		

 NH₄, NO₃, DON, Urea all as N

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

* = PN by wet digestion method, ** = PN by combustion furnace method.

Lake Taupo biannual nutrient database
Collection Date 30 October 1997

ID	Depth m	pH	EC @25°C µS cm⁻¹	Temp C	DO g m⁻³	SS g m⁻³	Secchi depth = 12.5 m			1997-1998			Started 27 October 1994			PN* mg m⁻³	TN mg m⁻³	DOC g m⁻³	PC mg m⁻³	PN** mg m⁻³	SO₄ g m⁻³
							VSS g m⁻³	Chlor_a mg m⁻³	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH₄ mg m⁻³	NO₃ mg m⁻³	DON mg m⁻³	UREA mg m⁻³					
TT1	1	7.70	116.9	12.2	10.7	0.61	0.30	1.28	1.0	1.3	1.5	3.8	2.1	2.9	36	1.1	14.3	55.3	0.71	168.3	17.2
TT2	10	7.71	117.8	12.0	10.2	0.54	0.29	1.49	0.7	1.9	1.9	4.5	1.3	7.3	32	1.1	18.7	59.7	0.82	160.7	18.8
TT3	20	7.65	118.1	11.5	10.2	0.59	0.32	1.58	0.8	1.6	1.7	4.0	1.6	0.7	36	1.1	14.0	52.0	0.60	133.0	16.5
TT4	30	7.64	118.2	11.5	10.0	0.52	0.25	1.19	0.4	1.5	1.9	3.8	1.5	1.3	31	0.9	15.8	49.8	0.60	146.9	16.0
TT5	40	7.62	117.1	11.4	10.0	0.55	0.28	1.31	0.6	1.5	1.6	3.7	1.7	0.3	33	1.0	14.1	49.1	0.62	126.3	13.4
TT6	50	7.63	116.9	11.1	9.9	0.37	0.20	1.10	0.4	1.5	1.4	3.2	2.2	0.3	32	0.8	12.3	46.3	0.51	112.1	12.1
TT7	60	7.54	117.7	11.1	9.8	0.21	0.10	0.93	1.4	0.7	1.5	3.5	3.3	0.7	34	1.6	14.3	52.3	0.74	80.6	9.0
TT8	70	7.45	117.8	10.8	9.8	0.41	0.12	0.79	1.1	1.1	1.1	3.2	8.2	1.3	31	1.5	7.9	47.9	0.65	58.4	4.8
TT9	80	7.36	118.3	10.7	9.9	0.31	0.04	0.54	1.5	1.1	0.8	3.3	6.1	2.3	31	0.6	6.0	45.0	0.57	57.6	9.0
TT10	90	7.48	117.8	10.6	9.3	0.44	0.27	0.74	1.1	1.2	1.2	3.5	7.9	4.8	33	0.7	12.4	58.4	0.52	69.3	12.2
TT11	100	7.29	118.5	10.5	9.2	0.25	0.11	0.40	2.0	1.2	0.8	4.1	8.4	5.0	30	1.1	5.7	48.7	0.63	64.5	8.3
TT12	110	6.97	119.3	10.4	9.0	0.21	0.06	0.29	2.3	1.0	1.1	4.3	10.8	5.6	29	2.5	6.7	51.7	0.59	53.0	5.5
TT13	120	7.00	119.1	10.5	9.0	0.29	0.26	0.27	2.0	1.2	1.0	4.1	9.9	6.7	31	6.1	5.8	53.8	0.58	37.5	5.3
TT14	130	6.80	119.8	10.5	8.8	0.28	0.26	0.28	2.2	1.2	1.3	4.7	10.6	7.1	32	1.5	8.2	58.2	0.56	49.0	6.4
TT15	140	7.23	117.9	10.4	8.8	0.25	0.20	0.26	2.7	1.4	1.1	5.2	10.8	9.5	37	2.0	10.9	67.9	0.63	66.0	8.5
TT16	150	7.29	118.9	10.4	8.8	0.50	0.27	0.32	2.5	1.1	1.0	4.5	11.6	9.6	37	3.0	7.6	65.6	0.54	69.0	9.2

Collection Date:- 7 April 1998

ID	Depth m	pH	EC @25°C µS cm⁻¹	Temp C	DO g m⁻³	SS g m⁻³	Secchi depth = 13.5 m			1997-1998			Started 27 October 1994			PN* mg m⁻³	TN mg m⁻³	DOC g m⁻³	PC mg m⁻³	PN** mg m⁻³	SO₄ g m⁻³
							VSS g m⁻³	Chlor_a mg m⁻³	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH₄ mg m⁻³	NO₃ mg m⁻³	DON mg m⁻³	UREA mg m⁻³					
YE1	1	8.00	118	17.7	9.1	0.40	0.10	0.67	0.8	1.4	1.3	3.5	2.9	4.6	53	3.7	9.9	70.4	0.83	156.5	14.4
YE2	10	7.99	119	17.7	9.1	0.49	0.12	1.04	0.9	1.4	1.8	4.1	1.9	2.5	52	4.6	13.7	70.1	0.78	179.5	16.0
YE3	20	8.00	119	17.7	9.1	0.32	0.32	1.07	0.7	1.5	1.7	3.9	2.4	1.5	48	3.7	12.6	64.5	0.71	162.5	15.2
YE4	30	7.99	120	17.5	9.1	0.30	0.20	1.06	0.7	1.7	1.6	4.0	2.0	1.2	48	3.7	12.7	63.9	0.78	138.5	14.5
YE5	40	7.60	120	13.7	9.3	0.13	0.13	1.18	1.2	1.0	1.2	3.4	2.0	3.1	39	4.2	8.2	52.3	0.69	112.5	8.2
YE6	50	7.50	120	11.5	9.3	0.34	0.00	0.75	2.4	0.9	0.9	4.2	2.5	4.5	52	3.2	6.5	65.5	0.65	88.0	6.7
YE7	60	7.38	120	11.0	9.3	0.11	0.00	0.49	3.0	0.7	0.8	4.5	1.5	11.7	32	3.2	5.3	50.5	0.72	74.5	5.8
YE8	70	7.32	121	10.8	9.2	0.20	0.00	0.33	3.1	0.9	0.6	4.6	1.0	17.7	38	3.7	4.0	60.7	0.78	57.5	4.1
YE9	80	7.23	120	10.6	9.1	0.24	0.24	0.24	3.5	0.6	0.8	4.9	1.4	23.1	43	6.9	5.7	73.2	0.69	49.5	4.5
YE10	90	7.27	121	10.6	9.1	0.31	0.21	0.17	4.4	0.6	0.7	5.7	1.3	24.1	41	6.5	5.6	72.0	0.68	47.5	4.9
YE11	100	7.29	121	10.6	9.0	0.32	0.11	0.16	4.5	0.7	0.8	6.0	1.0	24.5	39	3.7	6.8	71.3	0.57	58.0	7.4
YE12	110	7.29	121	10.5	8.9	0.35	0.35	0.12	4.8	0.7	0.5	6.0	1.3	25.1	40	5.5	6.5	72.9	0.63	52.5	2.6
YE13	120	7.35	121	10.5	8.9	0.24	0.08	0.37	3.4	0.6	1.2	5.2	1.0	18.9	35	4.6	4.1	59.0	0.75	63.5	3.8
YE14	130	7.24	122	10.5	8.8	0.32	0.16	0.11	5.7	0.6	0.7	7.0	1.0	27.0	39	6.0	3.5	70.5	0.63	52.0	3.9
YE15	140	7.21	122	10.5	8.6	0.45	0.05	0.15	6.4	0.6	1.0	8.0	4.2	29.1	65	10.6	6.7	105.0	0.74	60.5	5.9
YE16	150	7.49	121	10.5	8.4	0.80	0.15	0.62	3.3	1.1	1.6	6.0	2.5	13.0	62	9.7	14.2	91.7	0.70	135.5	13.6

 NH₄, NO₃, DON, Urea all as N

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

* = PN by wet digestion method, ** = PN by combustion furnace method.

Lake Taupo biannual nutrient database
Collection Date 24 October 1996

ID	Depth m	pH	EC @25°C μS cm⁻¹	Temp C	DO g m⁻³	Secchi depth = 12.6 m						1996-1997				Started 27 October 1994						
						SS g m⁻³	VSS g m⁻³	Chlor_a mg m⁻³	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH₄ mg m⁻³	NO₃ mg m⁻³	DON mg m⁻³	UREA mg m⁻³	PN* mg m⁻³	TN mg m⁻³	DOC g m⁻³	PC mg m⁻³	PN** mg m⁻³	SO₄ g m⁻³
IG1	1			12.4	10.3	0.45	0.34	0.27	0.6	2.1	1.7	4.4	3.0	0.5	59.3	1.4	13.9	76.7	0.86	171	14.5	7.82
IG2	10			12.3	10.3	0.72	0.42	0.47	0.7	2.3	2.2	5.2	2.4	0.4	64.5	1.0	14.5	81.8	0.88	201	16.8	7.90
IG3	20			12.3	10.2	0.67	0.40	0.45	0.8	2.8	2.9	6.5	2.6	0.4	75.8	0.6	18.7	97.5	0.91	232	19.8	7.87
IG4	30			12.3	9.9	0.85	0.49	0.64	0.6	2.3	3.1	6.0	3.3	0.5	73.6	0.4	20.6	98.0	0.95	198	15.7	7.86
IG5	40			11.9	9.9	0.71	0.46	0.56	0.5	1.8	2.5	4.8	2.6	1.2	64.8	0.3	14.6	83.2	0.80	183	12.8	7.84
IG6	50			11.6	9.8	0.62	0.34	0.45	1.1	3.1	2.1	6.3	2.9	0.6	71.2	0.9	13.2	87.9	0.92	157	14.9	7.95
IG7	60			11.1	9.7	0.77	0.32	0.70	0.9	1.8	2.3	5.0	4.4	13.2	175.4	3.5	14.3	207.3	1.29	151	14.1	10.67
IG8	70			10.6	9.4	0.65	0.28	0.54	0.8	1.5	1.9	4.2	2.9	0.8	59.3	1.5	9.2	72.2	0.78	116	10.2	7.85
IG9	80			10.5	9.3	0.51	0.27	0.55	0.9	2.5	1.8	5.2	3.0	3.0	76.1	1.3	9.8	91.9	0.95	103	10.8	7.80
IG10	90			10.4	9.3	0.49	0.23	0.50	0.6	1.8	1.8	4.2	2.1	1.0	52.3	1.4	10.9	66.3	0.73	95	11.0	7.69
IG11	100			10.4	9.2	0.50	0.21	0.51	0.5	1.5	1.8	3.8	1.8	3.6	53.9	4.5	9.6	68.9	1.04	106	12.8	7.85
IG12	110			10.4	9.2	0.43	0.23	0.49	0.4	1.3	2.0	3.7	2.5	5.2	54.0	6.0	9.3	71.0	0.80	94	11.5	7.85
IG13	120			10.4	9.0	0.47	0.21	0.47	0.8	1.4	1.8	4.0	3.7	9.6	61.9	6.9	8.0	83.2	0.78	78	9.7	7.97
IG14	130			10.3	8.9	0.44	0.18	0.38	1.1	1.5	2.3	4.9	4.5	9.7	52.4	4.6	12.0	78.6	1.00	83	8.7	7.99
IG15	140			10.3	8.9	0.49	0.22	0.51	1.5	1.6	2.5	5.6	4.3	12.9	57.8	5.0	10.4	85.4	0.99	80	8.9	8.14
IG16	150			10.3	8.9	1.13	0.26	0.57	1.2	2.3	3.5	7.0	5.1	13.6	65.9	4.8	14.5	99.1	0.91	121	13.4	8.15

Collection Date:- 2 April 1997

ID	Depth m	pH	EC @25°C μS cm⁻¹	Temp C	DO g m⁻³	Secchi depth = 16.0 m						1996-1997				Started 27 October 1994						
						SS g m⁻³	VSS g m⁻³	Chlor_a mg m⁻³	DRP mg m⁻³	DOP mg m⁻³	PP mg m⁻³	TP mg m⁻³	NH₄ mg m⁻³	NO₃ mg m⁻³	DON mg m⁻³	UREA mg m⁻³	PN* mg m⁻³	TN mg m⁻³	DOC g m⁻³	PC mg m⁻³	PN** mg m⁻³	SO₄ g m⁻³
NA1	1	8.02	118.4	17.3	9.4	0.30	0.30	0.63	0.9	2.2	1.5	4.6	4.0	0.6	67.4	4.9	18.1	90.1	0.82	186.5	17.3	7.80
NA2	10	8.01	118.3	17.3	9.2	0.20	0.10	0.69	0.9	1.3	1.6	3.8	1.7	0.3	51.0	3.3	14.4	67.4	0.77	190.0	17.1	7.86
NA3	20	8.03	118.2	17.2	8.9	0.40	0.30	0.63	0.6	1.2	1.6	3.4	1.8	0.3	51.8	2.2	17.6	71.5	0.75	192.0	19.1	7.85
NA4	30	7.98	118.4	17.2	8.8	0.40	0.40	0.52	0.7	1.0	1.5	3.2	2.5	0.6	47.5	2.7	15.2	65.8	0.56	207.5	20.3	7.90
NA5	40	7.52	118.5	14.2	8.8	0.20	0.20	0.72	0.8	1.8	1.4	4.0	2.7	0.3	53.2	4.1	13.3	69.5	0.69	158.0	15.2	7.91
NA6	50	7.32	119.3	11.3	8.6	0.00	0.00	0.39	1.5	1.4	1.0	3.9	11.2	3.1	54.7	4.5	9.7	78.7	0.62	116.5	10.6	7.88
NA7	60	7.18	120.2	10.9	8.6	0.20	0.20	0.16	1.7	1.3	0.8	3.8	3.7	10.1	48.9	2.1	10.5	73.2	0.86	100.0	13.8	7.88
NA8	70	7.13	119.6	10.6	8.5	0.10	0.10	0.12	1.9	1.7	0.8	4.4	4.3	11.8	58.3	2.2	8.0	82.4	0.83	75.0	8.7	7.87
NA9	80	7.12	120.1	10.5	8.5	0.10	0.10	0.05	3.3	1.4	0.7	5.4	6.9	26.9	82.4	16.9	6.7	122.9	0.98	77.5	9.9	7.90
NA10	90	7.12	120.4	10.5	8.5	0.00	0.00	0.25	3.6	2.2	0.7	6.5	28.9	22.9	108.3	7.4	8.1	168.2	0.63	110.5	8.8	8.00
NA11	100	7.10	120.4	10.5	8.4	0.20	0.20	0.04	4.4	1.2	0.8	6.4	10.7	22.5	72.0	5.2	7.1	112.3	0.85	71.0	8.3	7.97
NA12	110	7.07	120.6	10.4	8.3	0.20	0.20	0.02	3.7	2.0	0.8	6.5	2.9	21.9	52.5	3.8	6.4	83.7	1.01	77.0	9.6	7.93
NA13	120	7.07	120.5	10.4	8.2	0.30	0.20	0.02	3.3	2.4	0.8	6.5	6.4	22.8	56.4	4.2	13.0	98.6	0.70	113.5	15.4	7.88
NA14	130	7.08	120.4	10.4	8.0	0.20	0.20	0.01	4.3	1.6	0.8	6.7	6.2	27.9	56.7	6.2	8.2	99.0	0.81	118.5	11.0	7.97
NA15	140	7.10	121.1	10.4	7.6	0.40	0.40	0.04	4.5	1.7	1.2	7.4	3.9	28.9	58.5	7.9	24.7	116.0	0.80	212.5	28.8	7.91
NA16	150	7.10	122.1	10.4	7.5	1.20	0.40	0.07	5.0	1.0	2.7	8.7	8.6	29.0	61.5	11.8	20.2	119.3	2.07	234.5	22.1	7.97

 NH₄, NO₃, DON, Urea all as N

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

* = analysed by wet digestion method, ** = analysed by CHN combustion furnace method.

Lake Taupo biannual nutrient database
Collection Date:- 30 October 1995

ID	Depth	pH	EC @25°C	Temp	DO	BOD ₅	Secchi depth = 13.0 m													1995-1996				
							μS cm ⁻¹	C	g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ mg m ⁻³	NO ₃ mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC g m ⁻³	PC mg m ⁻³
ZH1	1	7.40		115.1	13.7	10.3	0.80	0.60	0.38	0.45	<0.2	2.4	1.27	3.67	<0.2	<0.1	55.7	3	6.89	62.69	0.75	123	10.3	
ZH2	10	7.59		116.1	11.9	10.5	0.40	0.95	0.53	0.96	<0.2	0.8	1.94	2.74	<0.2	<0.1	48.0	3	14.69	62.69	0.61	217	18.0	
ZH3	20	7.39		117.8	11.4	10.6	-0.05	1.09	0.59	1.18	0.3	1.5	2.41	4.21	0.2	<0.1	51.5	4	19.47	71.17	0.58	285	22.3	
ZH4	30	7.58		116.6	11.2	10.7	-0.15	1.15	0.58	1.26	0.2	0.7	2.21	3.11	<0.2	<0.1	44.6	2	17.83	62.43	0.45	242	19.4	
ZH5	40	7.48		116.2	10.9	10.7	0.00	0.91	0.57	1.22	<0.2	1.1	1.88	2.98	<0.2	<0.1	41.9	2	13.00	54.90	0.44	183	15.8	
ZH6	50	7.36		117.0	10.8	10.3	0.25	0.69	0.42	1.10	<0.2	0.8	1.71	2.51	<0.2	<0.1	41.7	3	8.55	50.25	0.43	116	10.3	
ZH7	60	7.28		117.2	10.7	10.3	0.70	0.49	0.28	1.03	<0.2	0.8	1.55	2.35	<0.2	0.1	41.1	3	7.75	48.95	0.40	110	10.3	
ZH8	70	7.25		117.8	10.5	10.2	0.50	0.64	0.43	1.03	<0.2	0.6	1.50	2.10	<0.2	0.2	40.4	2	7.27	47.87	0.38	108	9.9	
ZH9	80	7.25		117.5	10.5	10.2	0.40	0.72	0.43	1.19	<0.2	0.8	1.58	2.38	<0.2	0.7	41.4	2	7.19	49.39	0.48	115	12.1	
ZH10	90	7.30		118.0	10.5	10.1	0.00	0.72	0.40	1.27	0.3	0.6	1.59	2.49	<0.2	1.5	38.5	3	7.30	47.30	0.47	101	12.1	
ZH11	100	7.25		117.5	10.5	10.0	0.15	0.71	0.39	1.30	<0.2	0.2	1.77	1.97	<0.2	2.4	36.4	3	10.67	49.47	0.49	107	12.5	
ZH12	110	7.25		117.5	10.5	9.9	0.35	0.71	0.38	1.32	<0.2	0.9	1.69	2.59	0.5	4.6	44.3	3	10.26	59.66	0.52	93	13.1	
ZH13	120	7.23		117.3	10.5	9.9	0.30	0.70	0.41	1.35	<0.2	1.3	1.55	2.85	0.5	5.6	51.3	9	7.99	65.39	0.51	99	12.9	
ZH14	130	7.25		117.3	10.5	9.8	0.20	0.69	0.47	1.32	<0.2	0.4	1.89	2.29	1.3	6.6	49.7	7	13.42	71.02	0.55	112	18.5	
ZH15	140	7.25		117.3	10.5	9.6	0.40	0.97	0.47	1.60	<0.2	0.2	2.54	2.74	5.7	11.7	60.6	9	11.77	89.77	0.57	113	15.8	
ZH16	150	7.25		117.5	10.5	9.2	0.40	1.77	0.91	1.77	0.7	0.4	3.05	4.15	8.3	13.2	90.9	15	48.30	160.70	0.69	357	55.1	

Collection Date:- 28 March 1996

ID	Depth	pH	EC @25°C	Temp	DO	BOD ₅	Secchi depth = 14.6 m													1995-1996				
							μS cm ⁻¹	C	g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor_a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ mg m ⁻³	NO ₃ mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC g m ⁻³	PC mg m ⁻³
DR1	1	8.02		117.4	16.8	8.7	0.15	0.31	0.18	0.48	1.3	1.8	0.93	4.03	<0.2	4.7	91.0	1.4	12.69	108.39	0.35	118	9.7	
DR2	10	8.02		117.4	16.7	8.7	0.20	0.44	0.25	0.81	1.3	1.5	1.43	4.23	<0.2	7.4	111.0	6.2	12.60	131.00	0.42	149	12.3	
DR3	20	7.95		117.6	16.6	8.8	0.25	0.34	0.23	0.76	1.0	1.8	1.30	4.10	0.6	<0.1	60.0	2.0	11.70	72.30	0.35	126	11.7	
DR4	30	7.59		119.0	13.7	9.0	0.25	0.39	0.15	1.13	1.5	1.7	1.51	4.71	0.5	0.2	64.0	2.0	11.72	76.42	0.26	101	12.8	
DR5	40	7.43		118.9	12.4	8.8	0.25	0.35	0.16	0.97	1.3	1.4	1.41	4.11	1.1	<0.1	51.0	2.2	11.77	63.87	0.22	68	8.6	
DR6	50	7.34		119.5	11.6	8.6	0.10	0.32	0.14	0.71	1.8	1.5	1.17	4.47	0.8	5.0	68.0	3.5	8.76	82.56	0.18	60	6.4	
DR7	60	7.32		119.4	11.4	8.5	0.25	0.27	0.10	0.48	2.2	1.0	1.06	4.26	1.8	5.9	59.0	1.8	8.32	75.02	0.17	46	5.7	
FR8	70	7.29		120.4	11.6	8.5	0.25	0.23	0.13	0.28	2.3	1.5	0.80	4.60	<0.2	14.1	87.0	3.4	6.65	107.75	0.26	48	6.4	
DR9	80	7.20		120.8	11.2	8.3	0.20	0.30	0.14	0.17	2.9	1.3	0.83	5.03	1.5	10.0	68.0	1.4	5.15	84.65	0.23	45	5.5	
DR10	90	7.20		121.2	11.3	8.2	0.20	0.39	0.14	0.12	2.7	2.1	0.89	5.69	2.5	11.5	55.0	1.4	5.34	74.34	0.17	51	6.7	
DR11	100	7.24		121.3	10.9	8.2	0.05	0.45	0.19	0.10	2.8	1.8	0.93	5.53	2.2	11.4	72.0	8.1	9.25	94.85	0.22	46	6.9	
DR12	110	7.32		122.1	10.8	8.1	0.25	0.25	0.15	0.08	2.7	1.8	0.88	5.38	1.0	11.5	68.0	1.6	5.86	86.36	0.23	52	8.1	
DR13	120	7.39		120.2	10.7	8.3	0.15	0.24	0.11	0.09	2.8	1.2	0.74	4.74	2.2	11.2	75.0	3.8	3.91	92.31	0.26	34	5.3	
DR14	130	7.47		120.3	10.7	8.3	0.25	0.31	0.15	0.08	3.1	1.5	0.70	5.30	1.5	12.4	70.0	2.5	3.43	87.33	0.27	45	3.8	
DR15	140	7.43		121.1	10.7	8.0	0.15	0.33	0.15	0.08	4.6	1.4	0.96	6.96	2.9	16.0	88.0	5.7	4.28	111.18	0.26	51	7.4	
DR16	150	7.52		120.1	10.6	7.8	0.75	0.75	0.63	0.07	4.7	1.5	2.13	8.33	3.2	15.9	140.0	32.4	69.74	228.84	0.52	349	70.7	

 NH₄, NO₃, DON, UREA all as N

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

* = analysed by wet digest method, ** = analysed by CHN combustion furnace method.

Lake Taupo biannual nutrient database
Collection date:- 27 October 1994

		1994-1995																			
		Secchi Depth = 11.7 m																			
ID	Depth m	Temp C	DO g m ⁻³	BOD ₅ g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor. a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ mg m ⁻³	NO ₃ mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³	LEAD mg m ⁻³
MM1	1	11.7	10.5	0.30	0.93	0.55	1.16	1.6	0.7	2.5	4.8	1.1	0.2	61	0.1	16.6	78.9	0.67	193.3	20.3	0.22
MM2	10	11.5	10.6	0.35	0.86	0.49	0.97	1.5	0.4	2.5	4.4	2.2	0.1	50	<0.1	15.2	67.5	0.42	203.8	19.0	
MM3	20	11.5	10.8	0.70	0.87	0.58	0.92	1.2	1.1	2.8	5.1	5.1	<0.1	49	0.2	17.4	71.5	0.40	254.5	19.6	
MM4	30	11.3	10.7	0.30	0.86	0.54	0.99	1.2	0.0	2.3	3.5	<0.4	2.5	88	8.3	13.7	104.2	0.64	199.1	18.9	
MM5	40	10.9	10.5	0.05	0.83	0.49	0.97	1.0	1.4	2.1	4.5	0.4	<0.1	49	1.6	12.4	61.8	0.55	193.7	17.5	
MM6	50	10.9	10.4	0.15	0.85	0.48	0.83	1.0	0.9	2.2	4.1	<0.4	1.1	70	6.4	14.9	86.0	0.37	182.0	16.6	
MM7	60	10.8	10.4	0.00	1.04	0.53	0.88	1.1	0.9	2.1	4.1	<0.4	<0.1	47	1.0	13.6	60.6	0.46	184.6	20.0	
MM8	70	10.7	10.4	0.10	1.23	0.54	1.18	1.1	1.2	2.3	4.6	2.6	0.4	57	1.6	14.7	74.7	0.96	198.7	23.0	
MM9	80	10.6	10.4	0.35	1.07	0.45	1.37	1.0	1.4	2.4	4.8	1.2	0.1	47	1.0	15.3	63.6	0.51	154.4	22.6	
MM10	90	10.5	10.4	0.10	1.24	0.48	1.79	1.0	1.1	1.9	4.0	1.5	<0.1	43	1.3	15.6	60.1	0.48	152.0	22.0	
MM11	100	10.5	10.2	0.10	1.22	0.49	1.76	1.2	1.0	2.5	4.7	1.5	0.4	58	1.8	17.9	77.8	1.21	183.7	33.9	
MM12	110	10.5	10.3	0.45	1.15	0.48	1.78	1.4	0.4	3.0	4.8	1.4	0.4	52	1.9	16.8	70.6	0.65	105.8	28.4	
MM13	120	10.4	10.2	0.00	0.96	0.41	1.94	1.1	0.7	2.8	4.6	<0.4	0.6	61	1.6	16.7	78.4	1.00	106.7	29.8	
MM14	130	10.4	9.8	0.00	1.07	0.41	2.37	1.0	1.2	2.6	4.8	6.8	0.9	73	5.5	20.8	101.5	0.53	157.6	23.7	
MM15	140	10.4	9.8	0.00	1.63	0.57	2.32	1.1	1.1	2.3	4.5	3.7	0.9	61	1.9	20.6	86.2	0.44	176.0	19.2	0.36
MM16	150	10.3	9.9	0.25	1.73	0.75	2.49	1.8	0.8	2.3	4.9	4.2	1.9	60	12.1	39.6	105.7	0.57	303.6	44.0	1.09

MM17 Tube
Collection date:- 19 April 1995

		Secchi Depth = 16.1 m																			
ID	Depth m	Temp C	DO g m ⁻³	BOD ₅ g m ⁻³	SS g m ⁻³	VSS g m ⁻³	Chlor. a mg m ⁻³	DRP mg m ⁻³	DOP mg m ⁻³	PP mg m ⁻³	TP mg m ⁻³	NH ₄ mg m ⁻³	NO ₃ mg m ⁻³	DON mg m ⁻³	UREA mg m ⁻³	PN* mg m ⁻³	TN mg m ⁻³	DOC mg m ⁻³	PC mg m ⁻³	PN** mg m ⁻³	LEAD mg m ⁻³
SZ1	1	18.4	9.2	0.10	0.22	0.22	0.95	3.3	1.7	1.3	6.3	3.6	0.9	83	7.7	14.6	102.1	0.70	160.5	16.8	<0.5
SZ2	10	18.2	9.3	0.15	0.28	0.28	0.89	2.2	1.2	1.5	4.9	2.0	0.8	59	6.5	13.5	75.3	0.68	189.0	18.1	<0.5
SZ3	20	18.2	9.2	0.25	0.24	0.24	0.80	1.3	0.0	1.4	2.7	1.0	1.0	56	4.5	10.7	68.7	0.60	153.5	14.5	
SZ4	30	16.5	9.3	0.50	0.26	0.26	1.35	1.3	1.0	1.6	3.9	1.2	0.7	55	8.4	13.4	70.3	0.60	151.5	14.7	<0.5
SZ5	40	12.5	9.7	0.45	0.16	0.16	0.98	1.1	0.2	1.2	2.5	2.0	1.0	47	4.4	8.0	58.0	0.60	111.0	8.6	
SZ6	50	11.6	9.5	0.60	0.10	0.10	0.86	2.0	0.5	1.2	3.7	1.7	1.3	47	5.3	8.8	58.8	0.60	119.0	10.5	
SZ7	60	11.1	9.5	0.30	0.07	0.07	0.73	1.0	1.1	1.2	3.3	0.5	5.4	40	5.3	7.0	52.9	0.50	83.8	9.0	
SZ8	70	10.9	9.5	0.55	0.04	0.04	0.45	1.4	0.7	1.3	3.4	0.5	7.7	39	6.2	8.7	55.9	0.55	97.4	11.1	
SZ9	80	10.8	9.0	0.40	0.10	0.10	0.35	1.6	0.0	1.0	2.6	0.5	11.3	36	3.2	6.1	53.9	0.53	75.5	8.2	
SZ10	90	10.7	8.7	0.30	0.07	0.07	0.25	1.3	0.5	1.4	3.2	0.5	15.7	40	6.1	9.8	66.0	0.50	92.5	9.6	
SZ11	100	10.7	8.6	0.75	0.01	0.01	0.23	2.8	0.1	0.8	3.7	0.4	18.4	37	6.3	8.2	64.0	0.60	68.7	6.3	
SZ12	110	10.7	8.3	0.50	0.09	0.09	0.20	2.1	1.0	1.3	4.4	0.5	20.4	41	4.4	12.4	74.3	0.55	99.0	14.0	
SZ13	120	10.7	8.2	0.40	0.05	0.05	0.16	2.5	0.0	0.9	3.4	0.5	22.0	37	3.5	4.8	64.3	0.50	62.1	4.5	
SZ14	130	10.7	8.0	0.70	0.00	0.00	0.17	3.1	0.0	1.0	4.1	0.6	26.5	45	3.5	5.9	78.0	0.55	77.0	7.4	
SZ15	140	10.6	7.8	1.00	0.28	0.25	0.17	4.1	0.0	1.7	5.8	0.5	30.7	44	3.6	11.2	86.4	0.60	133.5	12.4	<0.5
SZ16	150	10.6	7.5	2.05	49.47	5.58	64.05	38.9	1.4	*	40.3	1.7	40.9	48	11.4	*	90.6	0.75	*	*	10.5

Surficial sediment

* = Sediment contamination, sample not filtered for analysis.

 Detection limits: DRP 0.5; NO₃-N 0.5; NH₄-N 1.0 mg m⁻³

* = analysed by wet digestion method, ** = analysed by CHN combustion furnace method.

8.5 Appendix 5: Phytoplankton data

In this report phytoplankton abundance is reported in cell counts per ml and biovolume per ml. In the previous system reporting only algal dominance, “Dominance” (rank 1 = dominant to rank 10 = rare), was calculated from algal biovolume. For continuance of the Dominance format, the species composition is ranked by biovolume.

Note: reporting counts as cells per ml rounded to a whole number may result in cell counts of “0” despite a large biovolume where the algal species is large or colonial e.g., *Botryococcus braunii*

The new algal data have been added to this report and the data from the previous year retained to accumulate, as has been done with temperature, DO, and nutrient data.

Note: *Leptolyngbya* sp. cells on 07/09/2009 are likely to have been washed off something rather than being local in 150 m of water.

Name changes:

Anabaena has changed to ***Dolichospermum*** as of August 2009. It will initially be referred to as follows: *Dolichospermum* sp. (formally; *Anabaena* sp.)

Lake Taupo phytoplankton enumeration (10-m tube) 2008-09

Cell counts and biovolume

Cells per ml numbers may be affected by rounding

	Sample code	RL4 16/09/2008	RL4 16/09/2008	SV2 14/10/2008	SV2 14/10/2008	UP4 26/11/2008	UP4 26/11/2008	XE2 22/12/2008	XE2 22/12/2008	XZ2 13/01/2009	XZ2 13/01/2009	XZ1 28/01/2009	XZ1 28/01/2009	AH2 11/02/2009	AH2 11/02/2009	AH4 25/02/2009	AH4 25/02/2009	DU1 26/03/2009	DU1 26/03/2009	EW2 15/04/2009	EW2 15/04/2009	GV2 7/05/2009	GV2 7/05/2009		
	Sampling date																								
	Species composition by class	Cell (per ml)	Biovolume (µm³)	Cell (per ml)	Biovolume (µm³)																				
Blue greens (Cyanophyceae)																									
<i>Anabaena lemmermannii</i>	0.0	0	0.0	0	46.5	1905	16.3	670	1.3	116	1.3	120	7.4	669	75.6	41	1.4	126	27.7	2495	13.6	1226	0		
<i>Pseudanabaena limnetica</i>	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.1	2	0.0	4.4	0	0.0	83	0.0	0.0	0.0	0		
<i>Anabaena planktonica</i>	0.0	0	0.0	0	0.0	0	0.0	0	0.8	299	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0.0	0		
<i>Anabaena</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0.0	0		
<i>Anabaena circinalis</i>	0.0	0	8.9	581	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0.0	0		
<i>Chroococcus</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0.3	4		
<i>Microcystis</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0		
<i>Leptolyngbya</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0.0	23		
<i>Snowella</i> sp.																							0		
Greens (Chlorophyceae)																									
<i>Monoraphidium</i> sp./ <i>Ankistrodesmus</i> <i>falcatus</i>	94	3956	4	172	4	172	16	688	53	2236	139	5848	56	2359	0	0	0	0	1	49	5	221			
<i>Stichococcus</i> <i>contortus</i>	12	211	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<i>Bryococcus</i> <i>braunii</i>	0.0	218	0.0	0	0.0	8877	0.0	127636	0.0	0	0.0	1908	0.0	0	0.0	543	0	0	0.0	4213	0.0	6058			
<i>Chlamydomonas</i> sp.	0	0	1	123	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<i>Elakothrix gelatinosa</i>	4	369	0	0	0	0	0	0	5	491	12	1229	16	1720	18	1843	0	0	1	114	0	0			
<i>Eudorina elegans</i>	0	0	0	0	0	0	0	0	6	1647	0	0	0	0	0	0	0	0	3	674	0	0			
<i>Nephrocystum lunatum</i>	0	0	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Oocystis</i> sp.	14	1994	8	1163	5	748	5	665	0	0	2	249	5	665	0	0	0	0	5	748	4	498			
<i>Tetradon</i> <i>gracile</i>	0	0	0	0	0	0	20	2252	9	1030	1	64	0	0	0	0	0	0	0	0	0	0			
<i>Paulschultzia</i> sp.	0	0	0	0	0	0	0	0	18	0	7	0	0	0	0	0	0	0	0	0	0	0			
<i>Dictyosphaeriace</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0	0	0			
<i>Crucigeniella</i> sp.	0	0	0	0	7	456	4	228	2	152	0	0	0	0	0	0	0	0	30	1969	53	3422			
Diatoms (Bacillariophyceae)																									
<i>Asterionella formosa</i>	64	18018	42	11794	29	8190	3	819	22	6061	35	9828	5	1310	1	328	4	1147	11	3112	19	5242			
<i>Aulacoseira granulata</i>	15	4534	0	0	0	0	0	0	0	0	0	0	0	0	0	8	2539	0	0	0	0	0	0		
<i>Aulacoseira</i> <i>granulata</i> var. <i>angustissima</i>	0	0	1	304	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Aulacoseira</i> sp.	12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Cyclotella stelligera</i>	15	2340	2	374	7	1123	0	0	1	187	1	187	1	187	0	0	1	187	1	187	4	655			
<i>Fragilaria crotonensis</i>	37	13194	33	11728	99	35603	66	23456	70	25132	21	7539	48	17173	16	5864	2	838	21	7539	8	2723			
<i>Nitzschia</i> sp.	0	0	0	0	0	0	4	1369	0	0	4	1597	2	913	2	913	0	0	0	0	0	0			
<i>Synechid</i> sp.	1	230	0	0	0	0	2	691	0	0	0	0	0	0	0	0	1	230	0	0	0	0			
<i>Amphora</i> sp.	0	0	0	0	0	0	1	306	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Cocconeis</i>	1	306	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Desmids (Mesotaeniaceae, Desmidiae)																									
<i>Closterium aciculare</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<i>Closterium acutum</i> var. <i>variable</i>	1	441	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Chrysophyta (Chrysophyceae)																									
<i>Dinobryon</i> sp.	0	0	53	3106	313	18466	23	1381	0	0	2	104	38	2243	53	3141	0	0	0	11	621	13	794		
<i>Cryptomonas</i> sp.	0	0	0	0	1	168	0	0	0	0	0	0	1	84	0	0	0	0	0	0	0	0			
Dinoflagellates (Dinophyceae)																									
<i>Ceratium hirundinella</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<i>Gymnodinium</i> sp. 1	0	0	0	0	1	205	1	205	1	4505	4	4505	3	3218	0	0	1	1287	1	644					
<i>Gymnodinium</i> sp. 2	0	0	0	0	1	14625	0	0	0	0	0	0	0	0	0	0	0	0	0	50	0	25			
<i>Peridinium</i> sp.	0	0	0	0	0	0	0	0	0	0	2	4680	1	2340	0	0	0	0	1	2340	0	0			
<i>Gonydinium</i> sp.																1	1170	1	1170	0	0	0			
Flagellates 5µm																									
Flagellates < 5µm/unicells	113	3972	68	2375	78	2723	249	8722	182	6368	57	2007	51	1781	83	2907	37	1290	51	1781	145	5078			

Lake Taupo phytoplankton enumeration (10-m tube) 2008-09 continued

Sample code Sampling date	GV4 27/05/2009	GV4 27/05/2009	JO1 18/06/2009	JO1 18/06/2009	KI1 6/07/2009	KI1 6/07/2009	NEW NAMES INTRODUCED August 2009	LT1 13/08/2009	LT1 13/08/2009	ND1 7/09/2009	ND1 7/09/2009
Species composition by class	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)		Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)
Blue greens (Cyanophyceae)											
<i>Anabaena lemmermannii</i>	9.4	849	5.8	41	0.3	28	<i>Dolichospermum c.f. lemmermannii</i> (formerly; <i>Anabaena c.f. lemmermannii</i>)	0.1	10	0.1	11
<i>Pseudanabaena limnetica</i>	0.0	0	0.0	0	1.0	19	<i>Pseudanabaena sp.</i> (formerly; <i>Dolichospermum planctonicum</i>)	0.0	0	0.0	0
<i>Anabaena planktonica</i>	0.2	88	0.0	0	0.0	0	<i>Anabaena planktonica</i> (formerly; <i>Anabaena</i> <i>sp.</i>)	0.0	0	0.0	0
<i>Anabaena</i> sp.	2.1	188	0.3	23	0.5	46	<i>Dolichospermum circinalis</i> (formerly; <i>Anabaena circinalis</i>)	0.0	0	0.0	0
<i>Anabaena circinalis</i>	0.0	0	0.0	0	0.0	0	<i>Chroococcus sp.</i>	0.0	0	0.0	0
<i>Chroococcus</i> sp.	0.1	1	0.0	0	0.0	0	<i>Microcystis sp.</i>	0.2	2	0.8	11
<i>Microcystis</i> sp.	0.0	0	0.0	0	0.0	0	<i>Leptolyngbya sp.</i>	0.0	0	2.5	53
<i>Leptolyngbya</i> sp.	0.6	6	0.1	2	0.0	0	<i>Snowella sp.</i>	0.0	120.0	1320	
<i>Snowella</i> sp.	0.1	3	0.0	0	0.0	0		3.3	83	222.9	5572
Greens (Chlorophyceae)											
<i>Monoraphidium</i> sp. / <i>Ankistrodesmus falcatus</i>	14	590	42	1744	42	1750	<i>Monoraphidium</i> sp. / <i>Ankistrodesmus falcatus</i>	24	1022	225	9459
<i>Stichococcus contortus</i>	0	0	3	53	0	0	<i>Stichococcus contortus</i>	19	351	63	1141
<i>Botryococcus braunii</i>	0.0	15954	0.0	14315	0.0	30946	<i>Botryococcus braunii</i> (colonies)	0.0	0	0.0	205716
<i>Chlamydomonas</i> sp.	0	0	0	0	0	0	<i>Chlamydomonas</i> sp.	0	0	0	0
<i>Elakothrix gelatinosa</i>	0	0	0	0	0	0	<i>Elakothrix gelatinosa</i>	1	114	8	819
<i>Eudorina elegans</i>	0	0	0	0	0	0	<i>Eudorina elegans</i>	0	0	0	0
<i>Nephrocystum lunatum</i>	0	0	0	0	0	0	<i>Nephrocystum lunatum</i>	0	0	0	0
<i>Oocystis</i> sp.	0	0	4	498	0	0	<i>Oocystis</i> sp.	15	2151	0	0
<i>Tetradon gracile</i>	0	0	0	0	0	0	<i>Tetradon gracile</i>	0	0	0	0
<i>Paulschulzia</i> sp.	0	0	0	0	0	0	<i>Paulschulzia</i> sp.	0	0	0	0
<i>Dictyosphaerium</i> sp.	0	0	0	0	0	0	<i>Dictyosphaerium</i> sp.	0	0	12	295
<i>Crucigeniella</i> sp.	36	2358	11	722	9	598	<i>Crucigeniella</i> sp.	2	141	0	0
Diatoms (Bacillariophyceae)											
<i>Asterionella formosa</i>	10	2785	22	6143	55	15299	<i>Asterionella formosa</i>	366	102400	215	60333
<i>Aulacoseira granulata</i>	7	2176	0	0	102	31529	<i>Aulacoseira granulata</i>	30	9392	18	5441
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	0	0	15	3955	0	0	<i>Aulacoseira granulata</i> var. <i>angustissima</i>	0	0	4	1014
<i>Aulacoseira</i> sp.	0	0	0	0	0	0	<i>Aulacoseira</i> sp.	0	0	0	0
<i>Cyclotella stelligera</i>	1	187	9	1404	2	346	<i>Cyclotella stelligera</i>	5	866	21	3432
<i>Fragilaria crotonensis</i>	18	6492	35	12566	24	8716	<i>Fragilaria crotonensis</i>	0	0	34	12217
<i>Nitzschia</i> sp.	1	456	2	913	2	844	<i>Nitzschia</i> sp.	5	2110	1	380
<i>Synedra</i> sp.	0	0	0	0	0	0	<i>Synedra</i> sp.	1	213	0	0
<i>Amphora</i> sp.	0	0	0	0	0	0	<i>Amphora</i> sp.	0	0	0	0
<i>Cocconeis</i>	0	0	1	306	0	0	<i>Cocconeis</i>	0	0	0	0
Desmids (Mesotaeniaceae, Desmidiaeae)											
<i>Cladophora aciculare</i>	0	0	1	350	0	0	<i>Cladophora aciculare</i>	0	0	0	0
<i>Cladophora acutum</i> var. <i>variable</i>	0	0	0	0	1	204	<i>Cladophora acutum</i> var. <i>variable</i>	0	0	1	368
Chrysophyta (Chrysophyceae)											
<i>Dinobryon</i> sp.	8	449	0	0	0	0	<i>Dinobryon</i> sp.	0	0	0	0
<i>Cryptomonas</i> sp.	0	0	1	84	1	78	<i>Cryptomonas</i> sp.	0	0	0	0
Dinoflagellates (Dinophyceae)											
<i>Ceratium hirundinella</i>	0	0	0	0	0	0	<i>Ceratium hirundinella</i>	0	0	0	0
<i>Gymnodinium</i> sp. 1	1	1287	1	644	2	1785	<i>Gymnodinium</i> sp. 1	0	0	0	0
<i>Gymnodinium</i> sp. 2	0	0	0	2925	0	0	<i>Gymnodinium</i> sp. 2	0	0	0	0
<i>Peridinium</i> sp.	0	0	0	0	0	0	<i>Peridinium</i> sp.	0	0	0	0
<i>Gonyaulax</i> sp.	1	2340	1	1170	0	0	<i>Gonyaulax</i> sp.	0	0	0	0
Flagellates 5μm											
Flagellates < 5μm/unicells	67	2334	51	1781	76	2651	Flagellates < 5μm/unicells	328	11494	193	6757

Lake Taupo phytoplankton enumeration (10-m tube) 2007-08

Cell counts and biovolume

Cells per ml numbers may be affected by rounding

	Sample code	TZ2 8/08/2007	TZ2 8/08/2007	TZ4 23/08/2007	TZ4 23/08/2007	WF2 11/09/2007	WF2 11/09/2007	XX1 9/10/2007	XX1 9/10/2007	XX4 30/10/2007	XX4 30/10/2007	AM1 15/11/2007	AM1 15/11/2007	BM1 4/12/2007	BM1 4/12/2007	BM3 20/12/2007	BM3 20/12/2007	DT1 17/01/2008	DT1 17/01/2008	EO1 31/01/2008	EO1 31/01/2008	EO3 14/02/2008	EO3 14/02/2008	EO5 27/02/2008	EO5 27/02/2008	
Species composition by class																										
	Sampling date	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	Cell (per ml)	Biovolume (μm^3)	
Blue greens (Cyanophyceae)																										
<i>Anabaena lemmermannii</i>	2	64	3	108	1	27	17	696	51	2100	18	725	1	27	29	1175	28.7	1175	21.3	875	25.0	1025	85.8	3518		
<i>Pseudanabaena limnetica</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.1	2	0.0	0	0.5	0	9		
<i>Chroococcus</i> sp.	0	0	1	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0	0.0	0	0.0	0	0		
<i>Microcystis</i> sp.	0	0	0	0	0	7	0	0	0	0	0	0	0	0	0	0	0	0.0	0	0.0	0	0.0	0	0		
<i>c.f. Rivularia</i> sp.	0	0	0	0	0	19	0	0	0	0	0	0	0	0	0	0	0	0.0	0	0.0	0	0.0	0	0		
<i>Aphaneotilice</i> sp.	0	0	1	15	0	1	0	0	0	0	0	0	0	0	0	0	0	0.0	0	0.0	0	0.0	0	0		
<i>Aphanizomenon</i> sp.	2	30	0	0	0	0	0	0	0	0	2	32	3	48	4	78	0.0	0	0.0	0	0.0	4.0	76			
<i>Leptolyngbya</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	1	8	0.0	0	0.0	0	0.0	0	0.0	0			
Greens (Chlorophyceae)																										
<i>Monoraphidium</i> sp./ <i>Ankistrodesmus</i> <i>falcatus</i>	20	839	17	695	3	123	6	247	10	418	28	1189	18	737	114	4785	66	2764	0	0	0	0	0	0	0	
<i>Stichococcus contortus</i>	175	0	97	1749	25	453	0	0	0	0	0	3	53	0	0	0	0	0	0	0	0	0	0	0		
<i>Kirchneriella contorta</i>	0	0	0	0	56	1853	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Botryococcus braunii</i>	0	0	0	4800	0	0	0	0	0	0	0	0	0	0	1	92840	0	0	0	0	0	0	0	259720		
<i>Chlamydomonas</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Elakothrix gelatinosa</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5	532	0	0	0	0	0	0	2	246		
<i>Eudorina elegans</i>	0	0	0	0	0	0	0	0	0	1	300	0	0	0	0	0	2	624	4	1108	0	0	3	749		
<i>Lagerheimia</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Oscytes</i> sp.	0	0	0	0	0	1	166	5	758	5	665	0	0	1	166	6	839	2	277	0	0	0	0	0		
<i>Planktothrix gelatinosa</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Quadrilateralis lacustris</i>	0	0	5	788	3	480	0	0	0	0	0	0	0	0	3	554	0	0	0	0	0	0	0			
<i>Westella botryoides</i>	10	634	29	1909	0	0	0	0	9	608	0	0	0	0	0	0	17	1077	0	0	0	0	0			
<i>Paulschulzia</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Diatoms (Bacillariophyceae)																										
<i>Astroionella formosa</i>	276	77123	292	81787	763	210074	124	34938	62	17263	15	4187	4	983	2	472	50	14060	11	3181	0	0	2	655		
<i>Aulacoseira granulata</i> var.	0	0	0	0	13	3990	0	0	16	5078	3	993	0	0	0	0	0	0	0	0	0	0	0			
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	52	13436	11	2777	0	0	0	0	3	761	0	0	0	0	0	2	507	0	0	0	0	0	0			
<i>Cyclotella stelligera</i>	14	2184	11	1709	8	1310	9	1452	11	1685	0	0	0	0	0	1	156	0	0	0	0	0	0			
<i>Fragilaria crotonensis</i>	57	20419	27	9750	0	0	0	0	0	0	2	574	1	209	9	3324	19	6806	5	1743	0	0	13	4607		
<i>Nitzschia</i> sp.	0	0	5	2083	1	228	0	0	0	0	0	0	0	1	456	14	5596	1	380	0	0	0	2	684		
<i>Synechra</i> sp.	1	0	0	0	0	1	1638	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0			
Small unknown diatom sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	60	0	0	0	0		
Desmids (Mesotaeniaceae, Desmidaceae)																										
<i>Closterium aciculare</i>	0	0	0	0	0	0	0	0	160	0	0	1	320	1	350	1	506	0	0	0	0	0	0	0		
<i>Closterium acutum</i> var. <i>variable</i>	1	551	1	201	1	221	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Chrysophyta (Chrysophyceae)																										
<i>Dinobryon</i> sp.	21	1266	2	126	0	0	146	8633	297	17534	81	4789	76	4487	8	448	7	431	6	383	32	1915	73	4314		
<i>Cryptomonas</i> sp.	0	0	1	77	0	0	1	77	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Mallomonas</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Dinoflagellates (Dinophyceae)																										
<i>Gymnodinium</i> sp. 1	0	1463	0	0	0	0	1	3204	1	1755	0	0	1	1755	1	2532	0	0	6	17853	4	10820	16	40140		
<i>Gymnodinium</i> sp. 2	0	12188	1	13350	0	0	0	6675	0	0	0	0	0	0	0	6094	0	0	0	0	0	0	3	73125		
Flagellates 5μm																										
Flagellates < 5μm/unicells	153	6582	296	10354	112	3911	129	4504	93	3256	78	2729	125	4382	526	18403	83	2901	99	3465	39	1373	60	2109		

	Sample code	HT1	HT1	HT3	HT3	KB1	KB1	LB1	LB1	LB3	LB3	MW1	MW3	MW3	OL1	OL1	OL3	OL3	QA2	QA2	QA4	QA4	RL2	RL2			
	Sampling date	13/03/2008	13/03/2008	26/03/2008	26/03/2008	17/04/2008	17/04/2008	7/05/2008	7/05/2008	22/05/2008	22/05/2008	5/06/2008	5/06/2008	18/06/2008	18/06/2008	1/07/2008	1/07/2008	15/07/2008	15/07/2008	7/08/2008	7/08/2008	20/08/2008	20/08/2008	4/09/2008	4/09/2008		
	Species composition by class	Cell (per ml)	Biovolume (μm^3)																								
Blue greens (Cyanophyceae)																											
<i>Anabaena lemmermannii</i>	92	3778	7.0	288	56.6	2319	120.6	4946	2.2	91	1.1	46	1.7	71	12.2	500	9.8	403	0.8	32	0.2	7	0.9	37			
<i>Pseudanabaena limnetica</i>	0.0	0	0.0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0	0	2.8	53	0.3	5	0.0	0	0.0	0.0	0		
<i>Chroococcus</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0.0	0.0	0.0	0		
<i>Microcystis</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0.0	0.0	0.0	0		
<i>c.f Rivularia</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
<i>Aphananthece</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0.0	0.0	0.0	0		
<i>Aphanizomenon</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0	0	0.0	0	0.0	0	0.0	0.0	0.0	0.0	0		
<i>Leptolyngbya</i> sp.	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0	0.0	0	0.0	0	0.0	1.4	16	0.0	0.0	0		
Greens (Chlorophyceae)																											
<i>Monoraphidium</i> sp./ <i>Ankistrodesmus</i> <i>falcatus</i>	0	0	0	0	0	0	5	197	0	0	0	0.0	0	0	188	7907	0	0	73	3047	73	3071	130	5479			
<i>Stichococcus contortus</i>	0	0	0	0	0	0	0	0	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	26	474			
<i>Kirchneriella contorta</i>	0	0	0	0	0	0	0	0	0	0	0	0.0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Bryococcus braunii</i>	0.1	469151	0	14435	0.04	259837	0	104870	0	28871	0	132806	0.0	3609	0	5774	0.1	226456	0.0	5413	0	0	0.0	0.0	17746		
<i>Chlamydomonas</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Elatostethris gelatinosa</i>	2	246	6	676	1	123	4	369	2	246	1	123	0	0	1	114	0	0	0	0	0	0	0	0			
<i>Eudorina elegans</i>	8	2097	0	0	0	0	0	0	0	0	0	11	2696	0	0	0	0	0	0	9	2246	0	0	0			
<i>Lagerheimia</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1797			
<i>Oocystis</i> sp.	0	0	0	0	0	1	166	5	665	2	332	0	0	0	0	6	914	0	0	0	5	665	7	997	0	0	
<i>Planktosphaeria gelatinosa</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	10	1412		
<i>Quadrigula lacustris</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Westella botryoides</i>	0	0	0	0	0	0	0	15	951	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Fauschulzia</i> sp.	0	0	0	0	0	0	0	9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Diatoms (Bacillariophyceae)																											
<i>Asterionella formosa</i>	19	5242	12	3276	5	1310	10	2785	28	7862	25	6880	22	6061	25	7043	102	28501	191	53399	79	22113	94	26208			
<i>Aulacoseira granulata</i>	0	0	0	0	0	0	0	2	725	12	3808	13	4171	2	725	0	0	35	10700	151	46788	0	0	18	5622		
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	0	0	0	0	0	0	0	4	913	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
<i>Cyclotella stelligera</i>	0	0	0	0	0	0	0	3	468	1	187	2	374	1	94	4	562	1	94	1	187	12	1872	18	2902		
<i>Fragilaria crotonensis</i>	0	0	0	15	5445	4	1466	0	0	57	20315	61	21781	84	29948	46	16545	30	10890	18	6283	49	17592	59	20943		
<i>Nitzschia</i> sp.	1	228	1	342	3	1141	2	684	2	913	0	0	1	228	4	1369	4	1597	1	456	0	0	2	684			
<i>Synedra</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Small unknown diatom sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Desmids (Mesotaeniaceae, Desmidiaeae)																											
<i>Cladotrichum aciculare</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1051		
<i>Cladotrichum acutum</i> var. <i>variable</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0			
Chrysophyta (Chrysophyceae)																											
<i>Dinobryon</i> sp.	26	1519	2	104	4	242	8	483	8	466	9	518	0	0	9	518	0	0	0	0	0	0	0	0	20	1208	
<i>Cryptomonas</i> sp.	1	84	0	0	1	84	1	168	1	84	1	84	2	337	0	0	2	337	0	0	0	0	0	0	0	0	
<i>Mallomonas</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	1053	0	0	0	0	0	0	0		
Dinoflagellates (Dinophyceae)																											
<i>Gymnodinium</i> sp. 1	6	19305	42	126360	12	36855	5	1843	35	12285	5	1638	4	1229	0	0	6	2048	0	0	0	0	0	0	0		
<i>Gymnodinium</i> sp. 2	0	0	0	0	0	0	0	0	0	0	1	29250	0	7313	0	0	1	14625	0	0	0	0	0	0	0		
Flagellates 5µm																											
Flagellates <5µm/unicells	57	1986	56	1945	73	2539	131	4586	47	1638	63	2191	111	3890	121	4238	115	4013	87	3030	207	7228	104	3645			

Lake Taupo phytoplankton dominance plus enumeration (10-m tube) 2006-07

Dominance by biovolume (rank 1 = dominant,...rank 10 = rare), plus cell counts and biovolume from May 2007

	Sample code	EM8	EM10	EM13	EM17	EM20	EM23	EM27	EM29	EM31	EM34	EM36	EM38	EM40	EM40	EM42	EM42	RY2	RY2	RY2	RY5	RY5	RY5		
	Sampling date	26/09/2006	18/10/2006	1/11/2006	5/12/2007	14/12/2007	9/01/2007	8/02/2007	21/02/2007	3/04/2007	19/04/2007	8/05/2007	22/05/07	22/05/07	14/06/07	14/06/07	27/06/07	27/06/07	27/06/07	18/07/2007	18/07/2007	18/07/2007			
Species composition by class																									
Blue greens (Cyanophyceae)	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Rank	Biovolume (μm^3)	cell (per ml)	Rank	Biovolume (μm^3)	cell (per ml)	Rank	Biovolume (μm^3)	cell (per ml)	Rank	Biovolume (μm^3)	cell (per ml)	
<i>Anabaena lemmermannii</i>	5	5	5	5	9	5	9	9	3	4	5	4	6	303	10	8	450	15	5	1091	36	4	3652	17	
<i>Anabaena</i> sp.														0	0		0	0	10	29	0	0	0	0	
<i>Aphanizomenon</i> sp.														8	8	7	7	9	9	10	5	0	0	0	1
<i>Phormidium</i> sp.														10	10	10	10	0	0	0	0	0	0	0	0
Greens (Chlorophyceae)																									
<i>Ankistrodesmus falcatus</i> / <i>Schroederia</i> sp.																									
<i>Botryococcus braunii</i>	7	2	2	3	3	1	1	1	1	1	5	3	1	1014600	0	1	38448	1	9	120	5	0	0	0	0
<i>Chlorosarcinopsis</i> sp.	10	10																	8	438	0				
<i>Elakothrix gelatinosa</i>																									
<i>Eudorina elegans</i>	9	9	10	10	10																				
<i>Kirchneriella contorta</i>																									
<i>Monoraphidium</i> sp/ <i>Ankistrodesmus falcatus</i>	10	10	10	10	10	10	10	10	8	8	9	7	5	561	19	2	20456	259	2	5061	46	5	2574	12	
<i>Oocystis</i> sp.	7	8	9	9	9	10	7	7	10	10	10	9	9	43	1	6	3210	11	4	1605	5	9	293	1	
<i>Quadrigula lacustris</i>	9																								
<i>Stichococcus contortus</i>																									
<i>Westella botryoidea</i>	9	9	9	10	10	10	10	10																	
Diatoms (Bacillariophyceae)																									
<i>Asterionella formosa</i>	2	2	6	4	4																				
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	3	1	1	1	2	9	6	2	2	2	1														
<i>Cyclotella stelligera</i>	5	5	9	7	6	6	5	6																	
<i>Fragilaria crotonensis</i>	1	4	7	6	6	6	7	6	6																
<i>Gomphonema</i> sp.																									
<i>Nitzschia</i> sp.	10	10	10	10	10	10	10	10	10	10	7	8	8	155	1	5	5559	14	5	1042	3	7	952	2	
unknown diatom sp.																									
Desmids (Mesotaeniaceae, Desmidaceae)																									
<i>Closterium acutum</i>	9	10	10	9	9	7	8	8	10	10															
<i>Closterium acutum</i> var. <i>variable</i>	10	10	10	9	8	8	8	8																	
<i>Mougeotia</i> sp.																									
<i>Staurastrum</i> sp.	10	10																							
Chrysophyta (Chrysophyceae)																									
<i>Cryptomonas</i> sp.	10	10	10	2	1	2	6	8	3	10	10	10	10	0	0	9	267	1	9	196	1	9	293	1	
<i>Dinobryon</i> sp.	9	3	3	2	1	2	6	8	3	5	2	1	7	256	1	0	0	0	0	0	0	0	0	0	0
Dinoflagellates (Dinophyceae)																									
<i>Ceratium hirundinella</i>																									
<i>Gymnodinium</i> sp.	5	7	4	3	5	7	3	3	4	6	4	8	2	0	0	1	11748	1	0	0	0	0	0	0	0
<i>Gymnodinium</i> sp. 2																									
Flagellates 5μm																									
Flagellates < 5μm/unicells	3	6	8	6	6	6	2	4	5	4	3	4	4	2138	50	3	16227	381	1	7521	177	3	4133	97	

Lake Taupo phytoplankton species composition and biovolume (μm^3) 2008-2009

From Site A (Mid Lake) 14/10/2008

	Sample code	SZ1	SZ2	SZ3	SZ6	SZ11	SZ16	SZ1	SZ2	SZ3	SZ6	SZ11	SZ16
	Depth	Surface Cell (per ml)	10m Cell (per ml)	20m Cell (per ml)	50m Cell (per ml)	100m Cell (per ml)	150m Cell (per ml)	Surface Biovolume (μm^3)	10m Biovolume (μm^3)	20m Biovolume (μm^3)	50m Biovolume (μm^3)	100m Biovolume (μm^3)	150m Biovolume (μm^3)
Species composition by class													
Blue greens (Cyanophyceae)													
<i>Dolichospermum c.f. lemmermannii</i> (formally; <i>Anabaena c.f. lemmermannii</i>)	0.0	1.3	0.0	0.8	0.0	0.0	0.0	0	51	0	31	0	0
<i>Aphanothec sp.</i>	0.0	0.0	0.0	0.0	7.3	0.0	0.0	0	0	0	0	66	0
<i>Pseudanabaena sp.</i>	0.0	0	0.0	0.0	22.2	5.3	0.0	0	0	0	0	422	100
Greens (Chlorophyceae)													
<i>Monoraphidium sp. / Ankistrodesmus falcatus</i>	68	71	0.5	55	13	6	2875	2998	22	2318	545	273	
<i>Stichococcus contortus</i>	0	0	0.0	0	17	15	0	0	0	0	0	302	263
<i>Kirchneriella contorta</i>	0	0	0.0	1	0	0	0	0	0	0	36	0	0
<i>Botryococcus braunii</i> (colonies)	0.0	0	0.0	0.0	0	0	0	0	21653	16240	76507.95	0	
<i>Elakothrix gelatinosa</i>	5	10	2	2	0	0	491	1044	227	227	0	0	
<i>Nephrocytium agardhianum</i>	2	0	0	0	0	0	0	0	0	0	0	0	0
<i>Oocystis sp.</i>	6	1	4	1	4	1	831	166	581	166	498	166	
<i>Quadrigula lacustris</i>	2	0	0	0	0	0	384	0	0	0	0	0	
Diatoms (Bacillariophyceae)													
<i>Asterionella formosa</i>	94	71	102	71	6	2	26372	19820	28501	19984	1802	655	
<i>Aulacoseira granulata</i>	0	0	0	1	3	1	0	0	0	0	363	907	363
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	1	22	8	8	0	0	304	5628	2129	1977	0	0	
<i>Cyclotella stelligera</i>	5	4	11	4	2	2	842	562	1685	562	374	281	
<i>Fragilaria crotonensis</i>	151	42	9	183	15	7	54033	14870	3141	65552	5236	2513	
<i>Synedra sp.</i>	1	0	0	0	0	0	0	0	0	0	0	0	
<i>Eunotia sp.</i>	0	1	0	0	0	0	0	0	0	0	0	0	
Desmids (Mesotaeniaceae, Desmidiaeae)													
<i>Closterium aciculare</i>	1	1	0	1	0	0	701	701	0	701	0	0	
<i>Closterium acutum</i> var. <i>variable</i>	0	0	1	0	1	1	0	0	221	0	221	221	
Chrysophyta (Chrysophyceae)													
<i>Dinobryon sp.</i>	1	0	32	3	0	0	69	0	1898	173	0	0	
<i>Cryptomonas sp.</i>	0	1	0	1	0	0	0	84	0	84	0	0	
Dinoflagellates (Dinophyceae)													
<i>Gymnodinium sp. 2</i>	1	0	1	0	0	0	14625	0	14625	0	0	0	
Flagellates 5 μm													
Flagellates < 5 μm /unicells	132	201	111	140	24	13	4607	7023	3870	4914	839	450	

Lake Taupo phytoplankton species composition and biovolume (μm^3) 2008-2009

From Site A (Mid Lake) 14/10/2008

	Sample code	EU1	EU2	EU6	EU8	EU11	EU16	EU1	EU2	EU6	EU8	EU11	EU16
	Depth	Surface Cell (per ml)	10m Cell (per ml)	50m Cell (per ml)	70m Cell (per ml)	100m Cell (per ml)	150m Cell (per ml)	Surface Biovolume (μm^3)	10m Biovolume (μm^3)	50m Biovolume (μm^3)	70m Biovolume (μm^3)	100m Biovolume (μm^3)	150m Biovolume (μm^3)
Species composition by class													
Blue greens (Cyanophyceae)													
<i>Dolichospermum c.f. lemmermannii</i> (formally; <i>Anabaena c.f. lemmermannii</i>)	1.2	8.5	1.6	0.0	0.0	0.0	104	767	143	4	0	0	0
<i>Dolichospermum sp.</i> (formally; <i>Anabaena sp.</i>)	0.5	0.9	0.0	0.0	0.0	0.0	49	83	0	0	0	0	0
<i>Pseudanabaena sp.</i>	0.0	0.0	0.0	1.7	0.3	0.6	0	0	0	33	5	11	
Greens (Chlorophyceae)													
<i>Monoraphidium sp. / Ankistrodesmus falcatus</i>	0	0	54	2	19	2	0	0	2260	66	786	82	
<i>Botryococcus braunii</i> (colonies)	0.0	1	0	0	1	0	123784	1111500	370500	0	741000	0	
<i>Crucigeniella sp</i>	52	53	5	3	0	0	3399	3448	304	203	0	0	
<i>Elakothrix gelatinosa</i>	1	0	0	0	0	0	76	0	0	0	0	0	
<i>Eudorina elegans</i>	0	11	2	0	0	0	0	2796	599	0	0	0	
<i>Oocystis sp.</i>	3	0	2	0	1	0	410	0	222	0	111	0	
<i>Westella botryoides</i>	0	5	3	2	0	0	0	304	203	152	0	0	
<i>Paulschulzia sp.</i>	2	0	0	0	0	0	0	0	0	0	0	0	
Diatoms (Bacillariophyceae)													
<i>Asterionella formosa</i>	3	6	4	4	1	1	707	1638	1201	1092	218	218	
<i>Aulacoseira granulata</i>	0	2	4	9	5	1	0	605	1209	2660	1693	242	
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	0	2	6	0	0	2	0	507	1622	0	0	406	
<i>Cyclotella stelligera</i>	1	1	4	1	0	0	115	187	686	125	62	62	
<i>Fragilaria crotonensis</i>	6	10	0	0	0	1	2066	3630	0	0	0	419	
<i>Nitzschia sp.</i>	0	0	0	0	0	0	70	152	0	0	0	152	
Desmids (Mesotaeniaceae, Desmidiaeae)													
<i>Closterium aciculare</i>	0	0	0	0	0	0	0	0	117	0	0	0	
<i>Closterium acutum</i> var. <i>variable</i>	0	0	0	2	0	0	0	0	147	735	0	0	
Chrysophyta (Chrysophyceae)													
<i>Dinobryon sp.</i>	7	2	0	0	0	0	426	138	0	0	0	0	
<i>Cryptomonas sp.</i>	0	0	1	0	0	0	0	0	168	0	0	0	
Dinoflagellates (Dinophyceae)													
<i>Gymnodinium sp. 1</i>	0	2	0	0	0	0	0	2145	0	0	0	0	
<i>Gymnodinium sp. 2</i>	0	1	0	0	0	0	0	19500	0	0	0	0	
<i>Gonyaulax sp.</i>	1	1	0	0	0	0	2164	1560	0	0	0	0	
Flagellates 5μm													
Flagellates < 5 μm /unicells	34	46	27	22	10	9	1174	1611	956	778	355	300	

Lake Taupo phytoplankton species composition and biovolume (μm^3) 2007-2008

From Site A (Mid Lake) 30/10/2007

	Sample code	ZA1	ZA2	ZA3	ZA6	ZA8	ZA11	ZA16		ZA1	ZA2	ZA3	ZA6	ZA8	ZA11	ZA16
	Depth	Surface	10m	20m	50m	70m	100m	150m	Biovolume	Surface	10m	20m	Biovolume	Biovolume	Biovolume	Biovolume
		cell (per ml)	μm^3		cell (per ml)	cell (per ml)	μm^3	μm^3	μm^3	μm^3						
Blue greens (Cyanophyceae)																
<i>Anabaena lemmermannii</i>		18.7	22.0	2.9	0.4	0.0	0.0	1.6	1683	1976	257	33	0	0	0	140
<i>Chroococcus</i> sp.		0.0	0.0	0.0	0.1	0.0	0.0	0.0	0	0	0	1	0	0	0	0
<i>Aphanocapsa</i> sp.		0.0	0.0	0.0	6.9	0.0	5.8	6.6	0	0	0	62	0	52	59	
<i>Planktolyngbya</i> sp.		21.3	0.0	0.0	0.0	0.0	0.0	0.0	192	0	0	0	0	0	0	0
<i>Pseudanabaena</i> sp.		0.0	0.0	0.0	0.0	0.0	4.9	0.3	0	0	0	0	0	0	94	6
Greens (Chlorophyceae)																
<i>Monoraphidium</i> sp. / <i>Ankistrodesmus falcatus</i>		52	21	29	15	6	0	0	2187	885	1229	614	270	0	0	0
<i>Stichococcus contortus</i>		39	6	13	15	6	2	4	706	116	242	274	116	42	63	
<i>Botryococcus braunii</i> (colonies)		0	0	0	1	0	0	0	0	0	0	0	235139	0	804	0
<i>Eudorina elegans</i>		13	3	7	0	0	0	0	3295	749	1797	0	0	0	0	
<i>Crucigeniella</i> sp.		0	2	8	5	5	0	0	0	152	532	304	304	0	0	
<i>Nephrocystium agardhianum</i>		0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Oocysts</i> sp.		9	4	0	1	0	9	1	1246	498	0	166	0	1246	166	
Diatoms (Bacillariophyceae)																
<i>Asterionella formosa</i>		33	73	102	62	34	4	14	9173	20311	28665	17363	9500	983	3931	
<i>Aulacoseira granulata</i>		15	37	91	25	9	25	13	4715	11606	28109	7617	2902	7617	4171	
<i>Aulacoseira granulata</i> var. <i>angustissima</i>		0	0	0	0	0	3	0	0	0	0	0	0	761	0	
<i>Cyclotella stelligera</i>		6	8	22	9	5	9	10	1030	1217	3557	1404	842	1404	1591	
<i>Fragilaria crotonensis</i>		11	14	22	7	7	20	2	3770	5026	7958	2513	2513	7330	838	
<i>Nitzschia</i> sp.		0	0	0	0	0	1	0	0	0	0	0	0	228	0	
Desmids (Mesotaeniaceae, Desmidiaeae)																
<i>Cladophora aciculare</i>		1	1	0	1	1	1	1	701	350	0	526	526	350	350	
<i>Cladophora acutum</i> var. <i>variable</i>		1	1	0	0	0	0	0	221	265	0	44	0	0	0	
Chrysophyta (Chrysophyceae)																
<i>Dinobryon</i> sp.		275	182	227	135	108	1	0	16222	10734	13392	7938	6351	69	0	
<i>Cryptomonas</i> sp.		0	0	1	1	0	0	0	0	0	168	168	0	0	0	
Dinoflagellates (Dinophyceae)																
<i>Gymnodinium</i> sp. 1		0	1	1	1	1	0	0	0	3510	3510	1755	1755	0	0	
<i>Gymnodinium</i> sp. 2		0	1	0	1	0	0	0	0	14044	26750	1463	0	0	0	
Flagellates 5μm																
Flagellates < 5 μm /unicells		139	404	406	243	144	25	13	4853	14148	14210	8497	5037	860	450	

Lake Taupo phytoplankton species composition and biovolume (μm^3) 2007-2008

From Site A (Mid Lake) 17/04/2008

Species composition by class	Sample code	KA1	KA2	KA3	KA6	KA11	KA16	KA1	KA2	KA3	KA6	KA11	KA16
	Depth	Surface	10m	20m	50m	100m	150m	Surface	10m	20m	50m	100m	150m
		(per ml)	Biovolume (μm^3)										
Blue greens (Cyanophyceae)													
<i>Anabaena lemmermannii</i>	44.8	46.9	24.3	0.0	6.5	1.4	4031	4220	2183	0	584	16	
<i>Pseudanabaena</i> sp.	0.0	0.0	0.0	0.0	0.0	17.4	0	0	0	0	0	331	
Greens (Chlorophyceae)													
<i>Monoraphidium</i> sp. / <i>Ankistrodesmus falcatus</i>	14	3	8	8	0	1	590	123	344	344	0	49	
<i>Stichococcus contortus</i>	6	26	6	0	0	0	116	463	116	0	0	0	
<i>Botryococcus braunii</i> (colonies)	0	0	0	0	0	1	54	31352	6431	26908	1608	156759	
<i>Elakothrix gelatinosa</i>	0	1	1	0	1	0	0	154	123	0	123	0	
<i>Eudorina elegans</i>	0	6	0	0	0	0	75	1498	75	0	0	0	
<i>Crucigeniella</i> sp.	0	0	0	1	0	0	0	0	0	76	0	0	
<i>Oocystis</i> sp.	2	10	2	0	2	1	332	1412	332	0	332	83	
<i>Westella botryooides</i>	0	0	0	0	0	0	0	0	0	8	0	0	
Diatoms (Bacillariophyceae)													
<i>Asterionella formosa</i>	12	23	32	12	3	4	3276	6552	8935	3276	819	983	
<i>Aulacoseira granulata</i>	5	16	5	12	5	9	1484	4946	1484	3808	1632	2720	
<i>Cyclotella stelligera</i>	2	6	2	5	1	1	340	936	340	749	94	94	
<i>Fragilaria crotonensis</i>	4	10	39	1	1	1	1523	3427	14089	419	419	209	
<i>Nitzschia</i> sp.	0	0	22	0	0	0	0	0	8442	0	0	0	
Small unknown diatom sp.	0	0	0	0	1	0	0	0	0	0	64	0	
Desmids (Mesotaeniaceae, Desmidiaeae)													
<i>Closterium aciculare</i>	0	1	0	0	1	0	105	701	105	0	350	4	
<i>Closterium acutum</i> var. <i>variable</i>	0	1	2	2	0	0	0	221	662	662	0	22	
Chrysophyta (Chrysophyceae)													
<i>Dinobryon</i> sp.	64	164	101	0	0	0	3797	9664	5971	0	0	0	
<i>Cryptomonas</i> sp.	1	1	1	3	0	0	84	84	84	421	0	0	
Dinoflagellates (Dinophyceae)													
<i>Gymnodinium</i> sp. 1	1	1	1	0	0	0	3191	3191	3191	0	0	0	
<i>Gymnodinium</i> sp. 2	0	0	0	0	0	0	0	0	0	146	134	0	
Flagellates 5μm													
Flagellates < 5 μm /unicells	46	126	196	37	7	3	1619	4411	6850	1290	246	102	

Lake Taupo phytoplankton species composition and biovolume (μm^3) 2006-2007
 From Site A (Mid Lake) 1/11/2006

	Sample code	HW1	HW3	HW6	HW11	HW16	HW1	HW3	HW6	HW11	HW16
	Depth	surface	20 m	50 m	100 m	150 m	surface	20 m	50 m	100 m	150 m
		cell (per ml)	Biovolume (μm^3)								
Species composition by class											
Blue greens (Cyanophyceae)											
<i>Anabaena lemmermannii</i>	63	25	0	0	0	0	3488.1	1367	25	15	0
<i>Aphanocapsa</i> sp.	0	0	2	3	0	0	0	0	14	31	0
Greens (Chlorophyceae)											
<i>Botryococcus braunii</i> (colonies)	0	0	0	0	0	0	5151	5901	7321	0	0
<i>Chlorosarcinopsis</i> sp.	3	0	2	2	0	0	259	0	182	208	0
<i>Eudorina elegans</i>	2	5	6	0	0	0	621	1198	1498	0	0
<i>Kirchneriella contorta</i>	5	4	0	0	0	0	176	116	0	0	0
<i>Lagerheimia</i> sp.	0	1	1	0	0	0	0	125	166	0	0
<i>Monoraphidium</i> sp. / <i>Ankistrodesmus falcatus</i>	3	0	0	0	0	0	143	0	0	0	0
<i>Oocystis</i> sp.	7	6	6	6	3	0	1034	872	831	831	415
<i>Westella botryooides</i>	0	0	7	0	0	0	0	0	0	0	0
Diatoms (Bacillariophyceae)											
<i>Asterionella formosa</i>	14	8	7	8	2	0	3806	2129	1884	2211	573
<i>Aulacoseira granulata</i>	63	54	49	47	54	0	19413	16866	15052	14689	16594
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	0	0	2	3	0	0	0	0	456	837	0
<i>Cyclotella stelligera</i>	46	8	4	7	4	0	7301	1264	562	1123	655
<i>Fragilaria crotonensis</i>	5	0	2	8	3	0	1912	0	628	2723	1047
<i>Nitzschia</i> sp.	2	1	1	0	0	0	947	342	342	0	0
Desmids (Mesotaeniaceae, Desmidiaceae)											
<i>Closterium aciculare</i>	0	0	0	0	0	0	0	35	175	0	0
<i>Closterium acutum</i> var. <i>variable</i>	0	0	0	0	0	0	0	0	110	0	0
Chrysophyta (Chrysophyceae)											
<i>Dinobryon</i> sp.	8	4	6	0	0	0	458	242	362	0	0
Dinoflagellates (Dinophyceae)											
<i>Gymnodinium</i> sp. 1	0	1	0	0	0	0	0	2633	1316	0	88
<i>Gymnodinium</i> sp. 2	0	0	0	0	0	0	6068	0	0	0	0
Flagellates 5μm											
Flagellates < 5 μm /unicells	50	19	31	23	4	0	1750	676	1085	788	143

Lake Taupo phytoplankton species composition and biovolume (μm^3) 2006-2007
 From Site A (Mid Lake) 2/04/2007

	Sample code Depth	HW17 surface cell (per ml)	HW18 10 m cell (per ml)	HW19 20 m cell (per ml)	HW22 50 m cell (per ml)	HW27 100 m cell (per ml)	HW32 150 m cell (per ml)	HW17 surface Biovolume (μm^3)	HW18 10 m Biovolume (μm^3)	HW19 20 m Biovolume (μm^3)	HW22 50 m Biovolume (μm^3)	HW27 100 m Biovolume (μm^3)	HW32 150 m Biovolume (μm^3)	
Species composition by class														
Blue greens (Cyanophyceae)														
<i>Anabaena lemmermannii</i>	36	65	56	0	2	0	1493	2655	2286	5	86	10		
Greens (Chlorophyceae)														
<i>Botryococcus braunii</i> (colonies)	1	0	0	0	0	0	27630	0	0	41446	0	0		
<i>Monoraphidium</i> sp./ <i>Ankistrodesmus falcatus</i>	49	17	17	0	1	0	2064	725	725	0	25	0		
<i>Oocystis</i> sp.	2	1	1	0	1	0	332	166	125	0	166	0		
<i>Stichococcus contortus</i>	0	0	0	0	0	1	0	0	0	0	0	0		21
Diatoms (Bacillariophyceae)														
<i>Asterionella formosa</i>	0	0	1	0	0	1	0	82	246	0	0	164		
<i>Aulacoseira granulata</i>	2	0	0	5	11	8	544	0	0	1541	3264	2630		
<i>Aulacoseira granulata</i> var. <i>angustissima</i>	0	0	0	0	7	2	0	0	0	76	1901	608		
<i>Cyclotella stelligera</i>	1	1	1	1	2	1	168	94	94	234	374	140		
<i>Eunotia</i> sp.	0	0	0	0	4	0	0	0	0	0	0	0		
<i>Fragilaria crotonensis</i>	0	0	0	0	0	1	0	0	0	0	0	0		209
<i>Nitzschia</i> sp.	2	0	1	0	0	0	799	114	228	0	0	0		
Small unknown diatom sp.	0	0	0	0	1	0	0	0	0	0	0	64		0
Desmids (Mesotaeniaceae, Desmidiaeae)														
<i>Closterium aciculare</i>	0	0	0	1	4	0	0	0	0	350	2453	0		
<i>Closterium acutum</i> var. <i>variable</i>	0	0	0	1	0	0	0	0	0	331	0	0		
Chrysophyta (Chrysophyceae)														
<i>Cryptomonas</i> sp.	0	1	1	4	0	0	0	211	126	590	0	0		
<i>Dinobryon</i> sp.	0	0	0	1	0	0	0	0	0	86	0	0		
Dinoflagellates (Dinophyceae)														
<i>Gymnodinium</i> sp. 1	1	0	0	0	0	0	2106	878	878	176	0	0		
<i>Gymnodinium</i> sp. 2	1	1	1	0	0	0	14625	21938	14625	0	0	0		
Flagellates 5 μm														
Flagellates < 5 μm /unicells	185	97	84	127	16	10	6470	3389	2928	4433	573	338		

8.6 Appendix 6: Historical data

Historical data held by NIWA has frequently been referred to and included in some analysis or comparison of the data from the long-term monitoring programme. To ensure that these data are always readily available, copies of the relevant historical data are included in this report. These data are the spring and autumn profiles of NO₃-N and DRP from 1974 to 1990 extracted from archived data books. The nitrate data for 27 September 1979 was taken from Vincent (1983). Subsequent data can be found in the previous appendices.

Note that the profiles given are aligned with the spring data above the corresponding autumn data, by date. Note also that the early profiles were to a depth of 110 m rather than 150 m. Also, as there was no March or April data collected in 1976, for completeness, the last valid profile in that series (12 January 1976) has been included.

The elapsed time given is the number of days between the spring profile in about October and the autumn profile in March/April of the following year. The average elapsed time between the two samplings across all data from 1974 to 2006 is 165 days.

The historical data also include an un-paired profile from July 1987. As there were no data for April 1987 and the lake was still stratified in July, when the next period of monitoring began, the July 1987 may be used as an indication of the total mass of nutrient accumulation in that year. Because these data are for an un-paired profile in July and not April, if the data are converted to rate estimates the assumption must be made that there was no spring carryover and the elapsed time is longer, being estimated as the average elapsed time plus three months.

Because the 1976 and 1987 data are for periods other than spring (October/November) to autumn (March/April), these data points have been excluded from any regression analysis of time-series data although the data points have been plotted as an indication of levels/rates for those years.

Historical data from Site A in Lake Taupo
Nitrate concentrations (mg m⁻³)
Spring

Date	18/11/1974	16/10/1975	4/10/1977	10/10/1978	27/09/1979	5/10/1987	17/10/1988	6/10/1989
Depth (m)								
0	0.8	0.3	1.1	0.0	0.0	0.3	2.6	1.2
10	0.3	0.4	1.2	1.4	0.0	0.4	2.7	1.8
20	0.0	0.0	0.6	0.8	0.5	0.5	2.8	1.0
30	0.3	0.4	0.0	0.7	0.5	0.4	2.8	1.4
40	0.8	0.0	0.1	0.6	1.0	0.6	3.0	1.3
50	2.1	0.3	0.6	0.7	1.0	0.8	2.9	1.0
60	4.9	0.0	1.0	0.8	0.5	1.2	2.5	0.8
70	4.1	0.4	1.1	0.8	1.0	1.0	2.9	1.6
80	5.3	0.0	3.2	1.2	1.5	1.4	2.9	1.6
90	5.4	0.0	1.3	1.2	1.0	1.5	2.5	1.7
100	8.4	1.8	3.3	1.4	1.5	1.2	2.6	1.7
110	12.0	4.1	2.8	1.4	1.5	6.0	2.4	0.8
120			2.8	1.7	2.5	0.7	2.7	1.6
130			2.7	2.1	5.0	1.2	2.7	1.1
140			1.7	2.1	6.0	1.2	3.1	1.1
150			1.4	2.5	7.0	1.1	2.4	0.3

Autumn

Date	14/04/1975	12/01/1976	14/03/1978	10/04/1979	10/03/1980	7/07/1987	5/04/1988	4/04/1989	10/04/1990
Depth (m)									
0	0.8	0.5	0.0	0.3	0.0	2.0	1.1	2.1	0.1
10	0.4	1	0.0	0.0	0.3	1.6	1.3	2.5	0.6
20	0.2	0.2	0.0	0.0	0.0	1.0	1.3	2.4	1.3
30	0.1	0	0.0	0.0	0.0	0.2	1.1	2.5	1.2
40	0.3	0.2	0.0	0.3	0.2	0.9	2.2	2.4	1.7
50	0.5	0.3	0.0	1.0	0.8	1.1	4.0	4.9	4.9
60	4.2	1.3	0.0	7.3	4.9	14.5	12.3	5.2	3.4
70	5.6	1.5	2.2	11.1	6.2	16.4	14.6	5.1	12.0
80	9.2	8.3	4.9	12.7	9.4	16.1	16.9	10.9	11.2
90	11.2	11.1	5.8	13.5	13.5	18.5	19.0	13.5	12.4
100	12.4	14	7.4	15.0	14.4	19.8	20.7	17.1	17.1
110	16.0		9.2	14.8	15.7	20.2	19.1	20.4	16.2
120			10.1	15.0	16.7	20.9	18.6	23.3	18.2
130			8.0	16.6	18.9	21.9	21.5	24.2	17.9
140			11.0	17.3	19.4	22.1	25.4	27.1	22.4
150			14.2	19.7	19.9	21.5	27.0	28.6	24.2

DRP concentrations (mg m⁻³)
Spring

Date	18/11/1974	16/10/1975	4/10/1977	10/10/1978	5/10/1987	17/10/1988	6/10/1989
Depth (m)	???						
0	8.7	1.1	0.3	0.6	0.2	0.2	0.0
10	8.0	1.2	0.0	0.6	0.1	0.1	0.2
20	8.3	1.1	0.1	0.5	0.2	0.0	0.1
30	7.5	0.9	0.0	0.3	0.3	0.1	0.0
40	8.4	0.8	0.3	0.2	0.2	0.1	0.0
50	7.6	0.8	0.2	0.3	0.4	0.1	0.0
60	8.3	0.7	0.0	0.3	0.3	0.2	0.0
70	7.7	0.7	1.1	0.4	0.3	0.2	0.0
80	8.1	0.8	0.7	0.5	0.3	0.2	0.3
90	7.9	1.0	0.8	0.4	0.2	0.3	0.1
100	8.5	1.7	0.4	0.4	0.2	0.3	0.1
110	9.8	1.6	0.4	0.4	0.4	0.5	0.1
120		0.5	0.4		0.4	0.4	0.0
130		0.4	0.3		0.4	0.4	0.2
140		0.6	0.3		0.4	0.5	0.3
150		0.5	0.4		0.3	0.5	0.2

Autumn

Date	14/04/1975	12/01/1976	14/03/1978	10/04/1979	10/03/1980	7/07/1987	5/04/1988	4/04/1989	10/04/1990
Depth (m)									
0	0.8	1.4	0.2	0.1	0.7	1.9	0.1	0.0	0.2
10	0.5	1.4	0.2	0.1	0.4	2.2	0.1	0.0	0.0
20	0.5	7.0	0.2	0.1	0.3	0.9	0.2	0.0	0.1
30	0.5	2.5	0.2	0.1	0.2	1.0	0.2	0.0	0.2
40	0.5	0.2	0.2	0.4	0.5	0.9	0.6	0.2	0.5
50	0.5	0.9	0.7	1.0	0.7	0.7	1.1	0.5	1.1
60	1.0	0.1	0.7	1.6	1.0	3.4	2.0	0.6	0.9
70	1.0	0.8	1.0	2.0	1.1	3.7	2.2	0.9	1.9
80	1.7	1.2	1.5	2.2	1.6	3.6	2.7	1.1	1.7
90	2.0	2.0	1.8	2.4	2.2	4.1	2.9	1.3	1.8
100	2.2	3.3	1.9	2.7	2.4	4.6	3.1	1.9	2.6
110	2.9		2.4	2.8	2.6	4.5	2.9	2.7	2.1
120		2.7	2.9	2.7	4.7	3.0	3.4	2.5	
130		2.1	3.0	3.7	5.1	3.4	3.8	2.4	
140		2.8	3.6	3.6	5.3	4.4	4.5	3.5	
150		0.9	3.8	3.8	5.0	4.6	4.8	4.0	

Elapsed period (days)

147	88	161	182	165	270*	183	169	186
-----	----	-----	-----	-----	------	-----	-----	-----

??? = possible analytical problem (e.g., Si interference)

* = average period of 165 days plus 3 months